

VII. IRODALOMJEGYZÉK

Abramov E, Dolev I, Fogel H, Ciccotosto GD, Ruff E, Slutsky I.(2009): Amyloid-beta as a positive endogenous regulator of release probability at hippocampal synapses. *Nat Neurosci.* 2009 Dec;12(12):1567-76. doi: 10.1038/nn.2433.

Alberdi E, Sánchez-Gómez MV, Cavaliere F, Pérez-Samartín A, Zugaza JL, Trullas R, Domercq M, Matute C. (2010): Amyloid beta oligomers induce Ca²⁺ dysregulation and neuronal death through activation of ionotropic glutamate receptors. *Cell Calcium.* 2010 Mar; 47(3):264-72. doi: 10.1016/j.ceca.2009.12.010. Epub 2010 Jan 12.

Alle H, Roth A, Geiger JR.(2009): Energy-efficient action potentials in hippocampal mossy fibers. *Science* 2009; 325:1405–1408.

Ambrosio, A.F., Malva, J.O., Carvalho, A.P., Carvalho, A.M. (1997): Inhibition of N-, P/Q- and other types of Ca²⁺ channels in rat hippocampal nerve terminals by adenosine A1 receptor. *Eur. J. Pharmacol.* 340, 301±310.

Angelatou F, Pagonopoulou O, Maraziotis T, Olivier A, Villemeure JG, Avoli M, Kostopoulos G. (1993): Upregulation of A1 adenosine receptors in human temporal lobe epilepsy: a quantitative autoradiographic study. *Neurosci Lett.* 1993 Nov 26;163(1):11-4.

Angulo E1, Casadó V, Mallol J, Canela EI, Viñals F, Ferrer I, Lluís C, Franco R. (2003): A1 adenosine receptors accumulate in neurodegenerative structures in Alzheimer disease and mediate both amyloid precursor protein processing and tau phosphorylation and translocation. *Brain Pathol.* 2003 Oct;13(4):440-51.

Aronica E, Sandau US, Iyer A, Boison D.(2013): Glial adenosine kinase--a neuropathological marker of the epileptic brain. *Neurochem Int.* 2013 Dec;63(7):688-95. doi: 10.1016/j.neuint.2013.01.028. Epub 2013 Feb 4.

Baloyannis SJ. (2009): Dendritic pathology in Alzheimer's disease. *J Neurol Sci.* 2009 Aug 15;283(1-2):153-7. doi: 10.1016/j.jns.2009.02.370. Epub 2009 Mar 17.

Bayer TA, Wiestler OD, Wolf HK. (1995): Hippocampal loss of N-methyl-D-aspartate receptor subunit 1 mRNA in chronic temporal lobe epilepsy.

Berg, AT (2008). "Risk of recurrence after a first unprovoked seizure. *Epilepsia.* 49 Suppl 1: 13–8.

Bettler B¹, Mulle C (1995) Review: neurotransmitter receptors. II. AMPA and kainate receptors. *Neuropharmacology.* 1995 Feb;34(2):123-39.

Bliss TV, Collingridge GL (1993) A synaptic model of memory: long-term potentiation in the hippocampus. *Nature.* 1993 Jan 7;361(6407):31-9.

Boison D. (2005): Adenosine and epilepsy: from therapeutic rationale to new therapeutic strategies. *Neuroscientist*. 2005 Feb;11(1):25-36.

Boison D. (2006): Adenosine kinase, epilepsy and stroke: mechanisms and therapies. *Trends Pharmacol Sci*. 2006 Dec;27(12):652-8. Epub 2006 Oct 23. Review.

Boison D. (2008): The adenosine kinase hypothesis of epileptogenesis. *Prog Neurobiol*. 2008 Mar; 84(3):249-62. doi: 10.1016/j.pneurobio.2007.12.002. Epub 2007 Dec 23. Review.

Boison D. (2009): Engineered adenosine-releasing cells for epilepsy therapy: human mesenchymal stem cells and human embryonic stem cells. *Neurotherapeutics*. 2009 Apr; 6(2):278-83. doi: 10.1016/j.nurt.2008.12.001. Review

Boison D.(2012): Adenosine dysfunction in epilepsy. *Glia*. 2012 Aug;60(8):1234-43. doi: 10.1002/glia.22285. Epub 2011 Dec 22.

Borges K, Gearing M, McDermott DL, Smith AB, Almonte AG, Wainer BH, Dingledine R. (2003): Neuronal and glial pathological changes during epileptogenesis in the mouse pilocarpine model. *Exp Neurol*. 2003 Jul;182(1):21-34.

Bowser DN1, Khakh BS. (2007): Vesicular ATP is the predominant cause of intercellular calcium waves in astrocytes. *J Gen Physiol*. 2007 Jun;129(6):485-91. Epub 2007 May 15.

Brookmeyer R, Johnson E, Ziegler-Graham K, MH Arrighi. (2007): Forecasting the global burden of Alzheimer's disease. *Alzheimer's and Dementia*. [Retrieved 18 June 2008];3(3):186–91. doi:10.1016/j.jalz.2007.04.381

Chen Y, Dubé CM, Rice CJ, Baram TZ . (2008): Rapid loss of dendritic spines after stress involves derangement of spine dynamics by corticotropin-releasing hormone. *J Neurosci*. 2008 Mar 12;28(11):2903-11. doi: 10.1523/JNEUROSCI.0225-08.2008

Cheng , W. Logge, J. K. Low, B. Garner , T. Karl (2013): Novel behavioural characteristics of the *APP_{Swe}/PS1 Δ E9* transgenic mouse model of Alzheimer's disease *Behavioural Brain Research* Volume 245, 15 May 2013, Pages 120–127

Chourbaji S, Vogt MA, Fumagalli F, Sohr R, Frasca A, Brandwein C, Hörtnagl H, Riva MA, Sprengel R, Gass P. (2008): AMPA receptor subunit 1 (GluR-A) knockout mice model the glutamate hypothesis of depression. *FASEB J*. 2008 Sep;22(9):3129-34. doi: 10.1096/fj.08-106450. Epub 2008 May 20.

Ciruela F, Ferré S, Casadó V, Cortés A, Cunha RA, Lluís C, Franco R. (2006): Heterodimeric adenosine receptors: a device to regulate neurotransmitter release. *Cell Mol Life Sci*. 2006 Nov;63(21):2427-31.

Cohen-Gadol AA1, Pan JW, Kim JH, Spencer DD, Hetherington HH. (2004): Mesial temporal lobe epilepsy: a proton magnetic resonance spectroscopy study and a histopathological analysis. *J Neurosurg*. 2004 Oct;101(4):613-20.

Culebras A, Miller M, Bertram L, Koch J. (1987): Differential response of growth hormone, cortisol, and prolactin to seizures and to stress. *Epilepsia*. 1987 Sep-Oct;28(5):564-70.

Cull-Candy S, Brickley S, Farrant M. (2001): NMDA receptor subunits: diversity, development and disease. *Curr Opin Neurobiol*. 2001 Jun; 11(3):327-35.

Cunha C, Brambilla R, Thomas KL. (2010): A simple role for BDNF in learning and memory? *Front Mol Neurosci*. 2010 Feb 9;3:1. doi: 10.3389/neuro.02.001.2010. eCollection 2010.

Cunha RA. (2005): Neuroprotection by adenosine in the brain: From A(1) receptor activation to A (2A) receptor blockade. *Purinergic Signal*. 2005 Jun;1(2):111-34. doi: 10.1007/s11302-005-0649-1. Epub 2005 Mar 17.

Cunha RA.(2001): Adenosine as a neuromodulator and as a homeostatic regulator in the nervous system: different roles, different sources and different receptors. *Neurochem Int*. 2001 Feb; 38(2):107-25. Review.

de Quervain DJ, Aerni A, Schelling G, Roozendaal B. (2009): Glucocorticoids and the regulation of memory in health and disease. *Front Neuroendocrinol*. 2009 Aug;30(3):358-70. doi: 10.1016/j.yfrne.2009.03.002. Epub 2009 Mar 31.

Devinsky O, Vezzani A, Najjar S, De Lanerolle NC, Rogawski MA. (2013): Glia and epilepsy: excitability and inflammation. *Trends Neurosci*. 2013 Mar;36(3):174-84. doi: 10.1016/j.tins.2012.11.008. Epub 2013 Jan 5.

DeWitt DA, Perry G, Cohen M, Doller C, Silver J.(1998): Astrocytes regulate microglial phagocytosis of senile plaque cores of Alzheimer's disease. *Exp Neurol* 1998;149:329–340.

Dunwiddie, T.V., Fredholm, B.B.(1989): Adenosine A1 receptors inhibit adenylate cyclase activity and neurotransmitter release and hyperpolarize pyramidal neurons in rat hippocampus. *J. Pharmacol. Exp. Ther.* 249, 31±37.

Dunwiddie, T.V., Haas, H.L. (1985): Adenosine increases synaptic facilitation in the in vitro rat hippocampus: evidence for a presynaptic site of action. *J. Physiol.* 369, 365±377

During M. J., Spencer D. D. MD (2004): Adenosine: A potential mediator of seizure arrest and postictal refractoriness; *Annals of Neurology*; Volume 32, Issue 5, pages 618–624, November 1992

E Y Yuen, W Liu, I N Karatsoreos, Y Ren, J Feng, B S McEwen and Z Yan (2011): Mechanisms for acute stress-induced enhancement of glutamatergic transmission and working memory. *Molecular Psychiatry* (2011) 16, 156–170; doi:10.1038/mp.2010.50; published online 11 May 2010

Eadie MJ. (2012): Shortcomings in the current treatment of epilepsy. *Expert Rev Neurother*. 2012 Dec;12(12):1419-27. doi: 10.1586/ern.12.129.

Ekonomou A, Sperk G, Kostopoulos G, Angelatou F. (2000): Reduction of A1 adenosine receptors in rat hippocampus after kainic acid-induced limbic seizures. *Neurosci Lett*. 2000 Apr 21; 284(1-2):49-52.

Fedele DE, Gouder N, Güttinger M, Gabernet L, Scheurer L, Rüllicke T, Crestani F, Boison D. (2005): Astroglialosis in epilepsy leads to overexpression of adenosine kinase, resulting in seizure aggravation. *Brain*. 2005 Oct; 128(Pt 10):2383-95. Epub 2005 Jun 1.

Fedele DE, Li T, Lan JQ, Fredholm BB, Boison D. (2006): Adenosine A1 receptors are crucial in keeping an epileptic focus localized. *Exp Neurol*. 2006 Jul; 200(1):184-90. Epub 2006 Jun 5.

Ferreira, I.L., Bajouco, L.M., Mota, S.I., Auberson, Y.P., Oliveira, C.R., Rego, A.C., (2012): Amyloid beta peptide 1-42 disturbs intracellular calcium homeostasis through activation of GluN2B-containing N-methyl-d-aspartate receptors in cortical cultures. *Cell Calcium* 51, 95e106.

Fiacco TA1, McCarthy KD. (2004): Intracellular astrocyte calcium waves in situ increase the frequency of spontaneous AMPA receptor currents in CA1 pyramidal neurons. *J Neurosci*. 2004 Jan 21; 24(3):722-32.

Forrest D, Yuzaki M, Soares HD, Ng L, Luk DC, Sheng M, Stewart CL, Morgan JI, Connor JA, Curran T. (1994): Targeted disruption of NMDA receptor 1 gene abolishes NMDA response and results in neonatal death. *Neuron*. 1994 Aug;13(2):325-38.

Fredholm BB, Chen JF, Masino SA, Vaugeois JM. (2005): Actions of adenosine at its receptors in the CNS: insights from knockouts and drugs. *Annu Rev Pharmacol Toxicol*. 2005;45:385-412. Review.

Fredholm BB, Irenius E, Kull B, Schulte G. (2001): Comparison of the potency of adenosine as an agonist at human adenosine receptors expressed in Chinese hamster ovary cells. *Biochem Pharmacol*. 2001 Feb 15;61(4):443-8.

Garcia-Alloza M, Robbins EM, Zhang-Nunes SX, Purcell SM, Betensky RA, Raju S, Prada C, Greenberg SM, Bacskai BJ, Frosch MP. (2006): Characterization of amyloid deposition in the APP^{swe}/PS1^{dE9} mouse model of Alzheimer disease. *Neurobiol Dis*. 2006 Dec;24(3):516-24. Epub 2006 Oct 5.

Goebel DJ, Poosch MS.(1999): NMDA receptor subunit gene expression in the rat brain: a quantitative analysis of endogenous mRNA levels of NR1Com, NR2A, NR2B, NR2C, NR2D and NR3A. *Brain Res Mol Brain Res*. 1999 Jun 8;69(2):164-70.

Gomes CV, Kaster MP, Tomé AR, Agostinho PM, Cunha RA. (2011): Adenosine receptors and brain diseases: neuroprotection and neurodegeneration. *Biochim Biophys Acta*. 2011 May;1808(5):1380-99. doi: 10.1016/j.bbame.2010.12.001. Epub 2010 Dec 9. Review.

Gouder N, Scheurer L, Fritschy JM, Boison D.(2004): Overexpression of adenosine kinase in epileptic hippocampus contributes to epileptogenesis. *J Neurosci*. 2004 Jan 21; 24(3):692-701.

Groc L, Choquet D, Chaouloff F (2008): The stress hormone corticosterone conditions AMPAR surface trafficking and synaptic potentiation. *Nat Neurosci.* 2008 Aug;11(8):868-70. doi: 10.1038/nn.2150. Epub 2008 Jul 11.

Gulya Károly (2000): *Neurocitológia*. Szegedi Tudományegyetem, Természettudományi és Informatikai Kar. pp. 103-105.

Guthrie PB, Knappenberger J, Segal M, Bennett MV, Charles AC, Kater SB.(1999): ATP released from astrocytes mediates glial calcium waves. *J Neurosci.* 1999 Jan 15;19(2):520-8.

Halassa MM, Haydon PG. (2010): Integrated brain circuits: astrocytic networks modulate neuronal activity and behavior. *Annu Rev Physiol.* 2010;72:335-55. doi: 10.1146/annurev-physiol-021909-135843. Review.

Hanks AN, Dlugolenski K, Hughes ZA, Seymour PA, Majchrzak MJ. (2013): Pharmacological disruption of mouse social approach behavior: relevance to negative symptoms of schizophrenia.

Haydon PG, Carmignoto G. (2006): Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol Rev.* 2006 Jul;86(3):1009-31. Review.

Hynd, M.R., Scott, H.L., Dodd, P.R., (2004b): Selective loss of NMDA receptor NR1 subunit isoforms in Alzheimer's disease. *J. Neurochem.* 89, 240e247.

Iadecola C1, Nedergaard M. (2007): Glial regulation of the cerebral microvasculature. *Nat Neurosci.* 2007 Nov;10(11):1369-76.

Jacob, C.P., Koutsilieri, E., Bartl, J., Neuen-Jacob, E., Arzberger, T., Zander, N., Ravid, R., Roggendorf, W., Riederer, P., Grunblatt, E., (2007): Alterations in expression of glutamatergic transporters and receptors in sporadic Alzheimer's disease. *J. Alzheimers Dis.* 11, 97e116.

Jourdain P, Bergersen LH, Bhaukaurally K, Bezzi P, Santello M, Domercq M, Matute C, Tonello F, Gundersen V, Volterra A. (2007): Glutamate exocytosis from astrocytes controls synaptic strength. *Nat Neurosci.* 2007 Mar;10(3):331-9. Epub 2007 Feb 18.

Karst H, Joëls M. (2005): Corticosterone slowly enhances miniature excitatory postsynaptic current amplitude in mice CA1 hippocampal cells. *J Neurophysiol.* 2005 Nov;94(5):3479-86. Epub 2005 Jul 20.

Kennedy MB. (1994): The biochemistry of synaptic regulation in the central nervous system. *Annu Rev Biochem.* 1994;63:571-600.

Kito S, Miyoshi R, Nomoto T.(1990): Influence of age on NMDA receptor complex in rat brain studied by in vitro autoradiography. *J Histochem Cytochem.* 1990 Dec;38(12):1725-31.

Kowaluk EA, Jarvis MF. (2000): Therapeutic potential of adenosine kinase inhibitors. *Expert Opin Investig Drugs.* 2000 Mar;9(3):551-64. Review.

Kristel Sleepers, Nathalie Brouwers, Ilse Gijssels, Jessie Theuns, Dirk Goossens, Jan Wauters, Jurgen Del-Favero, Marc Cruts, Cornelia M. van Duijn, Christine Van Broeckhoven (2006): APP duplication is sufficient to cause early onset Alzheimer's dementia with cerebral amyloid angiopathy. *Brain*. 2006 Nov;129(Pt 11):2977-83. Epub 2006 Aug 18.

Kron MM1, Zhang H, Parent JM. (2010): The developmental stage of dentate granule cells dictates their contribution to seizure-induced plasticity. *J Neurosci*. 2010 Feb 10;30(6):2051-9. doi: 10.1523/JNEUROSCI.5655-09.2010.

Lai, CW; Trimble, MR (1997): Stress and epilepsy. *Journal of Epilepsy*. 10 177 - 186.

Lakhan SE, Caro M, Hadzimichalis N.(2013): NMDA Receptor Activity in Neuropsychiatric Disorders. *Front Psychiatry*. 2013 Jun 10; 4:52. doi: 10.3389/fpsyt.2013.00052. eCollection 2013.

Lan Xiong, MD, PhD, Claudia Gaspar, PhD, Guy A. Rouleau, MD, PhD, FRCPC (2005): Genetics of Alzheimer's Disease and Research Frontiers in Dementia. *Disclosures Geriatrics and Aging*. 2005;8(4):31-35.

Latini S, Pedata F. (2001): Adenosine in the central nervous system: release mechanisms and extracellular concentrations. *J Neurochem*. 2001 Nov;79(3):463-84. Review.

Lee, M.C., Ting, K.K., Adams, S., Brew, B.J., Chung, R., Guillemin, G.J. (2010): Characterisation of the expression of NMDA receptors in human astrocytes. *PLoS One*, e14123.

Li T, Lytle N, Lan JQ, Sandau US, Boison D.(2012): Local disruption of glial adenosine homeostasis in mice associates with focal electrographic seizures: a first step in epileptogenesis? *Glia*. 2012 Jan;60(1):83-95. doi: 10.1002/glia.21250. Epub 2011 Sep 30.

Li T, Ren G, Lusardi T, Wilz A, Lan JQ, Iwasato T, Itohara S, Simon RP, Boison D. (2008): Adenosine kinase is a target for the prediction and prevention of epileptogenesis in mice. *J Clin Invest*. 2008 Feb;118(2):571-82. doi: 10.1172/JCI33737

Lopes LV, Cunha RA, Ribeiro JA.(1999): Cross talk between A(1) and A(2A) adenosine receptors in the hippocampus and cortex of young adult and old rats. *J Neurophysiol*. 1999 Dec;82(6):3196-203.

Magistretti PJ.(2009): Low-cost travel in neurons. *Neuroscience*. 2009;325:1349–1351.

Maguire J, Salpekar JA. (2013): Stress, seizures, and hypothalamic-pituitary-adrenal axis targets for the treatment of epilepsy. *Epilepsy Behav*. 2013 Mar; 26 (3):352-62. doi: 10.1016/j.yebeh.2012.09.040. Epub 2012 Nov 29.

McEwen BS.(1999): Stress and hippocampal plasticity. *Annu Rev Neurosci*. 1999;22:105-22. Review.

Minkeviciene R1, Rheims S, Dobszay MB, Zilberter M, Hartikainen J, Fülöp L, Penke B, Zilberter Y, Harkany T, Pitkänen A, Tanila H. (2009): Amyloid beta-induced neuronal hyperexcitability triggers progressive epilepsy. *J Neurosci*. 2009 Mar 18;29(11):3453-62.

Mishizen-Eberz AJ, Rissman RA, Carter TL, Ikonovic MD, Wolfe BB, Armstrong DM. (2004): Biochemical and molecular studies of NMDA receptor subunits NR1/2A/2B in hippocampal subregions throughout progression of Alzheimer's disease pathology. *Neurobiol Dis*. 2004 Feb;15(1):80-92.

Mohamed A1, Posse de Chaves E. (2011): A β internalization by neurons and glia. *Int J Alzheimers Dis*. 2011 Feb 15;2011:127984. doi: 10.4061/2011/127984.a

Monaghan DT, Bridges RJ, Cotman CW. (1989): The excitatory amino acid receptors: their classes, pharmacology, and distinct properties in the function of the central nervous system. *Annu Rev Pharmacol Toxicol*. 1989;29:365-402. Review. No abstract available.

Mori F¹, Ribolsi M, Kusayanagi H, Siracusano A, Mantovani V, Marasco E, Bernardi G, Centonze D.(2011): Genetic variants of the NMDA receptor influence cortical excitability and plasticity in humans. *J Neurophysiol*. 2011 Oct;106(4):1637-43. doi: 10.1152/jn.00318.2011. Epub 2011 Jul 13.

Mota SI, Ferreira IL, Rego AC (2014): Dysfunctional synapse in Alzheimer's disease - A focus on NMDA receptors. *Neuropharmacology*. 2014 Jan;76 Pt A:16-26. doi:10.1016/j.neuropharm.2013.08.013. Epub 2013 Aug 22.

Mozhui K, Karlsson RM, Kash TL, Ihne J, Norcross M, Patel S, Farrell MR, Hill EE, Graybeal C, Martin KP, Camp M, Fitzgerald PJ, Ciobanu DC, Sprengel R, Mishina M, Wellman CL, Winder DG, Williams RW, Holmes A.(2010): Strain differences in stress responsivity are associated with divergent amygdala gene expression and glutamate-mediated neuronal excitability. *J Neurosci*. 2010 Apr 14;30(15):5357-67. doi: 10.1523/JNEUROSCI.5017-09.2010.

Nagele RG, D' Andrea MR, Lee H, Venkataraman V. Wang HY.(2003): Astrocytes accumulate A β 42 and give rise to astrocytic amyloid plaques in Alzheimer disease brains. *Brain Res* 2003;971:197-209

Nagele RG, Wegiel J, Venkataraman V, Imaki H, Wang KC, Wegiel J. (2004): Contribution of glial cells to the development of amyloid plaques in Alzheimer's disease. *Neurobiol Aging*. 2004 May-Jun;25(5):663-74. Review.

Neves G, Cooke SF, Bliss TV. (2012): Synaptic plasticity, memory and the hippocampus: a neural network approach to causality. *Nat Rev Neurosci*. 2008 Jan;9(1):65-75.

Neves G, Cooke SF, Bliss TV.(2008): Synaptic plasticity, memory and the hippocampus: a neural network approach to causality. *Nat Rev Neurosci*. 2008 Jan;9(1):65-75.

Newman EA. (2003): New roles for astrocytes: regulation of synaptic transmission. *Trends Neurosci*. 2003 Oct;26(10):536-42. Review.

Nicoll JA, Weller RO.(2003): A new role for astrocytes: beta-amyloid homeostasis and degradation. *Trends Mol Med.* 2003 Jul;9(7):281-2.

Olabarria M, Noristani HN, Verkhratsky A, Rodriguez JJ. (2010): Concomitant astroglial atrophy and astrogliosis in a triple transgenic animal model of Alzheimer's disease. *Glia* 2010;58:831–838

O'Toole KK, Hooper A, Wakefield S, Maguire J (2014): Seizure-induced disinhibition of the HPA axis increases seizure susceptibility. *Epilepsy Res.* 2014 Jan;108(1):29-43. doi: 10.1016/j.eplepsyres.2013.10.013. Epub 2013 Oct 28.

Palop JJ, Mucke L. (2009): Epilepsy and cognitive impairments in Alzheimer disease. *Arch Neurol.* 2009 Apr;66(4):435-40. doi: 10.1001/archneurol.2009.15. Epub 2009 Feb 9. Review.

Pandis D, Scarneas N. (2012): Seizures in Alzheimer disease: clinical and epidemiological data. *Epilepsy Curr.* 2012 Sep;12(5):184-7. doi: 10.5698/1535-7511-12.5.184.

Papa M, De Luca C, Petta F, Alberghina L, Cirillo G. (2014): Astrocyte-neuron interplay in maladaptive plasticity. *Neurosci Biobehav Rev.* 2014 Feb 6;42C:35-54. doi: 10.1016/j.neubiorev.2014.01.010. [Epub ahead of print] Review.

Parameshwaran K¹, Dhanasekaran M, Suppiramaniam V. (2008): Amyloid beta peptides and glutamatergic synaptic dysregulation. *Exp Neurol.* 2008 Mar;210 (1):7-13. Epub 2007 Oct 24.

Park, J., Jang,M., Chang, S., (2013): Deleterious effects of soluble amyloid-beta oligomers

Park, T.J., Chung, S., Han, M.K., Kim, U.H., Kim, K.T., 1997. Inhibition of voltage-sensitive calcium channels by the A2A adenosine receptor in PC12 cells. *J. Neurochem.* 71, 1251±1260.

Parodi, J., Sepulveda, F.J., Roa, J., Opazo, C., Inestrosa, N.C., Aguayo, L.G. (2010): Beta-amyloid causes depletion of synaptic vesicles leading to neurotransmission failure. *J. Biol. Chem.* 285, 2506e2514.

Parpura V, Verkhratsky A.(2012): Homeostatic function of astrocytes: Ca(2+) and Na(+) signalling. *Transl Neurosci.* 2012 Dec;3(4):334-344.

Pascual O, Casper KB, Kubera C, Zhang J, Revilla-Sanchez R, Sul JY, Takano H, Moss SJ, McCarthy K, Haydon PG. (2005): Astrocytic purinergic signaling coordinates synaptic networks. *Science.* 2005 Oct 7;310(5745):113-6.

Paula-Lima AC, Brito-Moreira J, Ferreira ST. (2013): Deregulation of excitatory neurotransmission underlying synapse failure in Alzheimer's disease. *J Neurochem.* 2013 Jul;126(2):191-202. doi: 10.1111/jnc.12304. Epub 2013 May 28.

Pearson T, Damian K, Lynas RE, Frenguelli BG. (2006): Sustained elevation of extracellular adenosine and activation of A1 receptors underlie the post-ischaemic inhibition of neuronal function in rat hippocampus in vitro. *J Neurochem*. 2006 Jun;97(5):1357-68.

Petroff OA, Errante LD, Rothman DL, Kim JH, Spencer DD. (2002): Glutamate-glutamine cycling in the epileptic human hippocampus. *Epilepsia*. 2002 Jul;43(7):703-10.

Plamondon H, Blondeau N, Heurteaux C, Lazdunski M. (1999): Mutually protective actions of kainic acid epileptic preconditioning and sublethal global ischemia on hippocampal neuronal death: involvement of adenosine A1 receptors and K(ATP) channels. *J Cereb Blood Flow Metab*. 1999 Dec;19(12):1296-308.

Puzzo D, Privitera L, Fa' M, Staniszewski A, Hashimoto G, Aziz F, Sakurai M, Ribe EM, Troy CM, Mercken M, Jung SS, Palmeri A, Arancio O.(2011): Endogenous amyloid- β is necessary for hippocampal synaptic plasticity and memory. *Ann Neurol*. 2011 May;69(5):819-30. doi: 10.1002/ana.22313. Epub 2011 Apr 6.

Querfurth HW, LaFerla FM. (2010): Alzheimer's disease. *N Engl J Med* 362:329–344. CrossRef Medline

Rebola N, Pinheiro PC, Oliveira CR, Malva JO, Cunha RA. (2003): Subcellular localization of adenosine A(1) receptors in nerve terminals and synapses of the rat hippocampus. *Brain Res*. 2003 Oct 10;987(1):49-58.

Rebola N, Rodrigues RJ, Oliveira CR, Cunha RA. (2005): Different roles of adenosine A1, A2A and A3 receptors in controlling kainate-induced toxicity in cortical cultured neurons. *Neurochem Int*. 2005 Oct;47(5):317-25.

Ribeiro JA.(2005): What can adenosine neuromodulation do for neuroprotection? *Curr Drug Targets CNS Neurol Disord*. 2005 Aug;4(4):325-9. Review.

Rodríguez JJ, Olabarria M, Chvatal A, Verkhratsky A. (2009): Astroglia in dementia and Alzheimer's disease. *Cell Death Differ*. 2009 Mar;16(3):378-85. doi: 10.1038/cdd.2008.172. Epub 2008 Dec 5. Review.

Rossner S, Lange-Dohna C, Zeitschel U, Perez-Polo JR. (2005): Alzheimer's disease beta-secretase BACE1 is not a neuron-specific enzyme. *J Neurochem* 2005;92:226–234.

Sántha P., Pákási M. , Fazekas Ö. , Fodor E., Kálmán S., Kálmán J. Jr., Janka Z. , Szabó Gy., Kálmán J. (2012): Restraint stress in rats alters gene transcription and protein translation in the hippocampus. *Neurochemical Research* .May 2012, Volume 37, Issue 5, pp 958-964

Scemes E, Giaume C. (2006): Astrocyte calcium waves: what they are and what they do. *Glia*. 2006 Nov 15;54(7):716-25. Review.

Scemes E, Giaume C.(2006): Astrocyte calcium waves: what they are and what they do. *Glia*. 2006 Nov 15;54(7):716-25.

Schaeffer EL, Gattaz WF. (2008): Cholinergic and glutamatergic alterations beginning at the early stages of Alzheimer disease: participation of the phospholipase A2 enzyme. *Psychopharmacology (Berl)*. 2008 May;198(1):1-27. doi: 10.1007/s00213-008-1092-0. Epub 2008 Feb 19.

Schell MJ1, Molliver ME, Snyder SH. (1995): D-serine, an endogenous synaptic modulator: localization to astrocytes and glutamate-stimulated release. *Proc Natl Acad Sci U S A*. 1995 Apr 25;92(9):3948-52.

Sebastião AM, Ribeiro JA.(1996): Adenosine A2 receptor-mediated excitatory actions on the nervous system. *Prog Neurobiol*. 1996 Feb;48(3):167-89.

Segal, M., (1982): Intracellular analysis of a post-synaptic action of adenosine in the rat hippocampus. *Eur. J. Pharmacol*. 49, 193±199.

Seifert G, Schilling K, Steinhäuser C. (2006): Astrocyte dysfunction in neurological disorders: a molecular perspective. *Nat Rev Neurosci*. 2006 Mar;7(3):194-206. Review.

Selkoe DJ.(2002): Alzheimer's disease is a synaptic failure. *Science*. 2002 Oct 25;298(5594):789-91.

Serrano A, Haddjeri N, Lacaille JC, Robitaille R. (2006): GABAergic network activation of glial cells underlies hippocampal heterosynaptic depression. *J Neurosci*. 2006 May 17;26(20):5370-82.

Shapiro LA, Ribak CE, Jessberger S. (2008): Structural changes for adult-born dentate granule cells after status epilepticus. *Epilepsia*. 2008 Jun;49 Suppl 5:13-8. doi: 10.1111/j.1528-1167.2008.01633.x. Review.

Shen HY, Lusardi TA, Williams-Karnesky RL, Lan JQ, Poulsen DJ, Boison D. (2011): Adenosine kinase determines the degree of brain injury after ischemic stroke in mice. *J Cereb Blood Flow Metab*. 2011 Jul;31(7):1648-59. doi: 10.1038/jcbfm.2011.30. Epub 2011 Mar 23.

Simpson JE, Ince PG, Lace G, et al.(2010): Astrocyte phenotype in relation to Alzheimer-type pathology in the ageing brain. *Neurobiol Aging* 2010;31:578–590.

Snyder EM, Nong Y, Almeida CG, Paul S, Moran T, Choi EY, Nairn AC, Salter MW, Lombroso PJ, Gouras GK, Greengard P.(2005): Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci*. 2005 Aug;8(8):1051-8. Epub 2005 Jul 17.

Stephen G. Waxman (2005): Összehasonlító neuroanatómia; Medicina kiadó

Temkin NR, Davis GR (1984): Stress as a risk factor for seizures among adults with epilepsy. *Epilepsia*. 1984 Aug;25(4):450-6.

Thompson K, Holm AM, Schousboe A, Popper P, Micevych P, Wasterlain C. (1998): Hippocampal stimulation produces neuronal death in the immature brain. *Neuroscience*. 1998 Jan;82(2):337-48.

Thompson, S.M., Hass, H.L., Gahwiller, B.H. (1992): Comparison of the actions of adenosine at pre- and post-synaptic receptors in the rat hippocampus in vitro. *J. Physiol.* 451, 347±363.

Thurman, DJ; Beghi, E; Begley, CE; Berg, AT; Buchhalter, JR; Ding, D; Hesdorffer, DC; Hauser, WA; Kazis, L; Kobau, R; Kroner, B; Labiner, D; Liow, K; Logroscino, G; Medina, MT; Newton, CR; Parko, K; Paschal, A; Preux, PM; Sander, JW; Selassie, A; Theodore, W; Tomson, T; Wiebe, S; (2011): "Standards for epidemiologic studies and surveillance of epilepsy." *Epilepsia*. 52 Suppl 7: 2–26.

Tsai G, Coyle JT.(2002): Glutamatergic mechanisms in schizophrenia. *Annu Rev Pharmacol Toxicol*. 2002;42:165-79. Review.

Tsien JZ, Huerta PT, Tonegawa S.(1996): The essential role of hippocampal CA1 NMDA receptor-dependent synaptic plasticity in spatial memory. *Cell*. 1996 Dec 27;87(7):1327-38.

Umemori J, Takao K, Koshimizu H, Hattori S, Furuse T, Wakana S, Miyakawa T.(2013): ENU-mutagenesis mice with a non-synonymous mutation in *Grin1* exhibit abnormal anxiety-like behaviors, impaired fear memory, and decreased acoustic startle response. *BMC Res Notes*. 2013 May 21;6:203. doi: 10.1186/1756-0500-6-203.

van Groen T, Kiliaan AJ, Kadish I.(2006): Deposition of mouse amyloid beta in human APP/PS1 double and single AD model transgenic mice. *Neurobiol Dis*. 2006 Sep;23(3):653-62. Epub 2006 Jul 10.

Verkhratsky A, Orkand RK, Kettenmann H. (1998): Glial calcium: homeostasis and signaling function. *Physiol Rev*. 1998 Jan;78(1):99-141.

Walter F. Boron and Emile L. Boulpaep (2011) *Medical Physiology* 2nd edition. Elsevier.

Wang HW, Pasternak JF, Kuo H et al. (2002): Soluble oligomers of beta amyloid (1-42) inhibit long-term potentiation but not long-term depression in rat dentate gyrus. *Brain Res*. 924:133-140, 2002.

Wetherington J, Serrano G, Dingledine R. (2008): Astrocytes in the epileptic brain. *Neuron*. 2008 Apr 24;58(2):168-78. doi: 10.1016/j.neuron.2008.04.002. Review.

Wilden, JA; Cohen-Gadol, AA (2012): Evaluation of first nonfebrile seizures. *American family physician* 86 (4): 334–40.

Wu, L.G., Saggau, P., (1994): Adenosine inhibits evoked synaptic transmission primarily by reducing presynaptic calcium influx in area CA1 of hippocampus. *Neuron*. 12, 1139±1148.

Yawo, H., Chuhma, N., (1993): Preferential inhibition of ω -conotoxin-sensitive presynaptic Ca^{2+} channels by adenosine autoreceptors. *Nature*. 365, 256±258.

Zarate C Jr, Machado-Vieira R, Henter I, Ibrahim L, Diazgranados N, Salvadore G.(2010): Glutamatergic modulators: the future of treating mood disorders? *Harv Rev Psychiatry*. 2010 Sep-Oct;18(5):293-303. doi: 10.3109/10673229.2010.511059. Review.

Zhao MG, Toyoda H, Lee YS, Wu LJ, Ko SW, Zhang XH, Jia Y, Shum F, Xu H, Li BM, Kaang BK, Zhuo M. (2005): Roles of NMDA NR2B subtype receptor in prefrontal long-term potentiation and contextual fear memory. *Neuron*. 2005 Sep 15;47(6):859-72.

Zimmermann H, Braun N.(1996): Extracellular metabolism of nucleotides in the nervous system. *J Auton Pharmacol*. 1996 Dec;16(6):397-400.

Ziyatdinova S1, Gurevicius K, Kutchiashvili N, Bolkvadze T, Nissinen J, Tanila H, Pitkänen A. (2011): Spontaneous epileptiform discharges in a mouse model of Alzheimer's disease are suppressed by antiepileptic drugs that block sodium channels. *Epilepsy Res*. 2011 Mar;94(1-2):75-85. doi: 10.1016/j.epilepsyres.2011.01.003.

<http://jaxmice.jax.org/strain/005864.html> (Utolsó letöltés: 2014. 04.10.)

http://www.frontiersin.org/files/Articles/30507/fendo-03-00113-HTML/image_m/fendo-03-00113-g002.jpg (Utolsó letöltés: 2014. 04.10.)

<http://www.ncbi.nlm.nih.gov/gene> ((Utolsó letöltés: 2014. 04.11.)

<http://www.ncbi.nlm.nih.gov/tools/primer-blast/> (Utolsó letöltés: 2014. 04.11.)

VIII. TÉTELES SZERZŐI HOZZÁJÁRULÁS

Alulírott*Ivitz Eszter Virág*..... a dolgozatomban leírt munkafolyamatok közül a:

- Teljes RNS izolálást
- mRNS-ről történő cDNS átírást
- Real-time PCR-t
- Kiértékelést

végeztem el.

.....
aláírás

Szeged, 2013. május 14.