

Hivatkozások

A dolgozat alapját képező publikáció:

Tőkési, N., Oláh, J., Hlavanda, E., Szunyogh, S., **Szabó, A.**, Babos, F., Magyar, A., Lehotzky, A., Vass, E. and Ovádi, J. (2014) Identification of motives mediating alternative functions of the neomorphic moonlighting TPPP/p25, *Biochim Biophys Acta*, 1842, 547-557.

További hivatkozások:

Abeliovich, A., Schmitz, Y., Farinas, I., Choi-Lundberg, D., Castillo, W.H. Ho, P.E., Shinsky, N. , Verdugo, J.M., Armanini, M., Ryan, A., Hynes, M., Phillips, H., Sulzer, D., Rosenthal, A. (2000) Mice lacking alpha-synuclein display functional deficits in the nigrostriatal dopamine system, *Neuron*, 25, 239–252

Bellucci, A., Navarria, L., Zaltieri, M., Missale, C. and Spano, P. (2012) alpha-Synuclein synaptic pathology and its implications in the development of novel therapeutic approaches to cure Parkinson's disease, *Brain Res*, 1432, 95-113.

Dunker, A.K., Obradovic, Z., Romero, P., Garner, E.C. and Brown, C.J. (2000) Intrinsic protein disorder in complete genomes, *Genome Inform Ser Workshop Genome Inform*, 11, 161-171.

Dunker, A.K., Silman, I., Uversky, V.N. and Sussman, J.L. (2008) Function and structure of inherently disordered proteins, *Curr Opin Struct Biol*, 18, 756-764.

Fink, A.L. (2005) Natively unfolded proteins, *Curr Opin Struct Biol*, 15, 35-41.

Hlavanda, E., Klement, E., Kokai, E., Kovacs, J., Vincze, O., Tokesi, N., Orosz, F., Medzihradszky, K.F., Dombradi, V. and Ovadi, J. (2007) Phosphorylation blocks the activity of tubulin polymerization-promoting protein (TPPP): identification of sites targeted by different kinases, *J Biol Chem*, 282, 29531-29539.

Hlavanda, E., Kovacs, J., Olah, J., Orosz, F., Medzihradszky, K.F. and Ovadi, J. (2002) Brain-specific p25 protein binds to tubulin and microtubules and induces aberrant microtubule assemblies at substoichiometric concentrations, *Biochemistry*, 41, 8657-8664.

Hoftberger, R., Fink, S., Aboul-Enein, F., Botond, G., Olah, J., Berki, T., Ovadi, J., Lassmann, H., Budka, H. and Kovacs, G.G. (2010) Tubulin polymerization promoting protein (TPPP/p25) as a marker for oligodendroglial changes in multiple sclerosis, *Glia*, 58, 1847-1857.

Hollósi, M., Laczkó, I., Majer, Zs. (2004) A sztereokémia és kiroptikai spektroszkópia alapjai. In: Nemzeti Tankönyvkiadó, Budapest, 168-195.

Irvine, G.B., El-Agnaf, O.M., Shankar, G.M. and Walsh, D.M. (2008) Protein aggregation in the brain: the molecular basis for Alzheimer's and Parkinson's diseases, *Mol Med*, 14, 451-464.

Iwai, A., Masliah, E., Yoshimoto, M., Ge, N., Flanagan, L., de Silva, H.A., Kittel, A. and Saitoh, T. (1995) The precursor protein of non-A beta component of Alzheimer's disease amyloid is a presynaptic protein of the central nervous system, *Neuron*, 14, 467-475.

Jeffery, C.J. (1999) Moonlighting proteins, *Trends Biochem Sci*, 24, 8-11.

Jeffery, C.J. (2011) Proteins with neomorphic moonlighting functions in disease, *IUBMB Life*, 63, 489-494.

Jenco, J. M., Rawlingson, A., Daniels, B., Morris, A. J. (1998) Regulation of phospholipase D2: selective inhibition of mammalian phospholipase D isoenzymes by alpha- and beta-synucleins, *Biochemistry*, 37, 4901-4909.

Kovacs, G.G., Laszlo, L., Kovacs, J., Jensen, P.H., Lindersson, E., Botond, G., Molnar, T., Perczel, A., Hudecz, F., Mezo, G., Erdei, A., Tirian, L., Lehotzky, A., Gelpi, E., Budka, H., Ovadi J. (2004) Natively unfolded tubulin polymerization promoting protein TPPP/p25 is a common marker of alpha-synucleinopathies, *Neurobiol Dis.*, 17, 155-162.

Leal, S.S. and Gomes, C.M. (2007) Studies of the molten globule state of ferredoxin: structural characterization and implications on protein folding and iron-sulfur center assembly, *Proteins*, 68, 606-616.

Lehotzky, A., Lau, P., Tokesi, N., Muja, N., Hudson, L.D. and Ovadi, J. (2010) Tubulin polymerization-promoting protein (TPPP/p25) is critical for oligodendrocyte differentiation, *Glia*, 58, 157-168.

Liliom, K., Wágner, G., Kovács, J., Comin, B., Cascante, M., Orosz, F. and Ovádi, J. (1999) Combined enhancement of microtubule assembly and glucose metabolism in neuronal systems *in vitro*. Decreased sensitivity to copper toxicity, *Biochem. Biophys. Res. Commun.*, 264, 605-610.

Lindersson, E., Lundvig, D., Petersen, C., Madsen, P., Nyengaard, J.R., Hojrup, P., Moos, T., Otzen, D., Gai, W.P., Blumbergs, P.C. and Jensen, P.H. (2005) p25alpha Stimulates alpha-synuclein aggregation and is co-localized with aggregated alpha-synuclein in alpha-synucleinopathies, *J Biol Chem*, 280, 5703-5715.

Na, C.N. and Timasheff, S.N. (1986) Interaction of vinblastine with calf brain tubulin: multiple equilibria, *Biochemistry*, 25, 6214-6222.

Oláh, J., Tokesi, N., Lehotzky, A., Orosz, F., Ovádi, J. (2013) Moonlighting microtubule-associated proteins: Regulatory functions by day and pathological functions at night, *Cytoskeleton (Hoboken)*, 70(11), 677-85.

Oláh, J., Vincze, O., Virok, D., Simon, D., Bozso, Z., Tokesi, N., Horvath, I., Hlavanda, E., Kovacs, J., Magyar, A., Szucs, M., Orosz, F., Penke, B. and Ovadi, J. (2011) Interactions of pathological hallmark proteins: tubulin polymerization promoting protein/p25, beta-amyloid, and alpha-synuclein, *J Biol Chem*, 286, 34088-34100.

Ovádi, J. (2011) Moonlighting proteins in neurological disorders, *IUBMB Life*, 63, 453-456.

Ovádi, J., Orosz, F. and Lehotzky, A. (2005) What is the biological significance of the brain-specific tubulin-polymerization promoting protein (TPPP/p25)?, *IUBMB Life*, 57, 765-768.

Piatigorsky, J. and Wistow, G.J. (1989) Enzyme/crystallins: gene sharing as an evolutionary strategy, *Cell*, 57, 197-199.

Preusser, M., Lehotzky, A., Budka, H., Ovadi, J. and Kovacs, G.G. (2007) TPPP/p25 in brain tumours: expression in non-neoplastic oligodendrocytes but not in oligodendroglioma cells, *Acta Neuropathol*, 113, 213-215.

Schägger, H. and von Jagow, G. (1987) Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa, *Anal. Biochem.*, 166, 368.

Seki, N., Hattori, A., Sugano, S., Suzuki, Y., Nakagawara, A., Muramatsu, M., Hori, T. and Saito, T. (1999) A novel human gene whose product shares significant homology with the bovine brain-specific protein p25 on chromosome 5p15.3, *J. Hum. Genet.*, 44, 121-122.

Semisotnov, G.V., Rodionova, N.A., Razgulyaev, O.I., Uversky, V.N., Gripas, A.F. and Gilmanshin, R.I. (1991) Study of the "molten globule" intermediate state in protein folding by a hydrophobic fluorescent probe, *Biopolymers*, 31, 119-128.

Shiratsuchi, A., Sato, S., Oomori, A., Ishiguro, K., Uchida, T. and Imahori, K. (1995) cDNA cloning of a novel brain-specific protein p25, *Biochem. Biophys. Acta*, 1251, 66-68.

Sickmeier, M., Hamilton, J.A., LeGall, T., Vacic, V., Cortese, M.S., Tantos, A., Szabo, B., Tompa, P., Chen, J., Uversky, V.N., Obradovic, Z. and Dunker, A.K. (2007) DisProt: the Database of Disordered Proteins, *Nucleic Acids Res*, 35, D786-793.

Spillantini, M.G., Crowther, R.A., Jakes, R., Hasegawa, M. and Goedert, M. (1998) alpha-Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with lewy bodies, *Proc Natl Acad Sci U S A*, 95, 6469-6473.

Takahashi, M., Tomizawa, K., Fujita, S. C., Sato, K., Uchida, T. and Imahori, K. (1993) A brain-specific protein p25 is localized and associated with oligodendrocytes, neuropil, and fiber-like structures of the CA3 hippocampal region in the rat brain, *J Neurochem* 60, 228-35.

Takahashi, M., Tomizawa, K., Ishiguro, K., Sato, K., Omori, A., Sato, S., Shiratsuchi, A., Uchida, T. and Imahori, K. (1991) A novel brain-specific 25 kDa protein (p25) is phosphorylated by a Ser/Thr-Pro kinase (TPK II) from tau protein kinase fractions, *FEBS Lettr.*, 289, 37-43.

Tirián, L., Hlavanda, E., Oláh, J., Horváth, I., Orosz, F., Szabó, B., Kovács, J., Szabad, J. and Ovádi, J. (2003) TPPP/p25 promotes tubulin assemblies and blocks mitotic spindle formation, *Proc Natl Acad Sci U S A*, 100, 13976-13981.

Tofaris, G. K., Spillantini, M. G. (2007) Physiological and pathological properties of alpha-synuclein, *Cell. Mol. Life Sci.*, 64, 2194-2201

Tőkesi, N., Lehotzky, A., Horváth, I., Szabó, B., Oláh, J., Lau, P., Ovádi, J. (2010) TPPP/p25 promotes tubulin acetylation by inhibiting histone deacetylase 6, *J Biol Chem.*, 285(23):17896-906.

Uversky, V.N. (2002) What does it mean to be natively unfolded?, *Eur. J Biochem.* , 2– 12., 269, 2– 12.

Uversky, V.N. (2003) A protein-chameleon: conformational plasticity of alpha-synuclein, a disordered protein involved in neurodegenerative disorders, *J Biomol Struct Dyn*, 21, 211-234.

Uversky, V.N. and Dunker, A.K. (2010) Understanding protein non-folding, *Biochim Biophys Acta*, 1804, 1231-1264.

Vincze, O., Olah, J., Zadori, D., Klivenyi, P., Vecsei, L. and Ovadi, J. (2011) A new myelin protein, TPPP/p25, reduced in demyelinated lesions is enriched in cerebrospinal fluid of multiple sclerosis, *Biochem Biophys Res Commun*, 409, 137-141.

Vincze, O., Tőkési, N., Oláh, J., Hlavanda, E., Zotter, Á., Horváth, I., Lehotzky, A., Tirián, L., Medzihradszky, K.F., Kovács, J., Orosz, F. and Ovádi, J. (2006) Tubulin polymerization promoting proteins (TPPPs): members of a new family with distinct structures and functions, *Biochemistry*, 45, 13818-13826.

Wistow, G. and Piatigorsky, J. (1987) Recruitment of enzymes as lens structural proteins, *Science*, 236, 1554-1556.

Zotter, A., Bodor, A., Olah, J., Hlavanda, E., Orosz, F., Perczel, A. and Ovadi, J. (2011a) Disordered TPPP/p25 binds GTP and displays Mg(2+)-dependent GTPase activity, *FEBS Lett*, 585, 803-808.

Zotter, A., Olah, J., Hlavanda, E., Bodor, A., Perczel, A., Szigeti, K., Fidy, J. and Ovadi, J. (2011b) Zn(2+)-induced rearrangement of the disordered TPPP/p25 affects its microtubule assembly and GTPase activity, *Biochemistry*, 50, 9568-9578.

Rövidítésjegyzék

AEBSF	4-(2-aminoetil)-benzénszulfonil-fluorid-hidroklorid
ANS	8-anilino-1-naftalin-szulfonsav
BLAST	„Basic Local Alignment Search Tool”
BSA	„bovine serum albumin” (szarvasmarha szérum albumin)
CD	cirkuláris dikroizmus
CHO	„chinese hamster ovary” (kínai hörcsög petefészek)
DAPI	4',6-diamidino-2-fenilindol
DTE	ditioeritol
EDTA	etilén-diamin-tetraecetsav
EGTA	etilén-glikol-bisz (b-aminoetil éter)-N, N, N', N'-tetraecetsav
ELISA	„enzyme-linked immunosorbent assay”
GTP	guanozin-trifoszfát
HDAC6	hiszton deacetiláz 6
IMAC	„immobilized metal affinity chromatography”
IPTG	izopropil- β -D-1-tiogalaktopiranozid
LB	„Luria-Bertani Broth”
MAP	Mikrotubulushoz Asszociált Protein
MES	2-(N-morfolino)-etánszulfonsav
NAC	non-amyloid- β component
NMR	„Nuclear Magnetic Resonance” (mágneses magrezonancia)
PAGE	poliakrilamid gélelektroforézis
PBS	„Phosphate Buffered Saline” (foszfát pufferelt fiziológiás sóoldat)
pEGFP	emlős expressziós vektor, ami EGFP-vel („enhanced green fluorescent protein”) fuzionált fehérjét képes expresszálni
SDS	nátrium-dodecil-szulfát
SEM	„standard error of the mean” (standard hiba)
SNCA	synuclein, alpha (non A4 component of amyloid precursor)
tPBS	tween-tartalmú PBS
TPPP	Tubulin Polymerization Promoting Protein
Tricin	N-(trisz-(hidroximetil)-metil)-glicin
Tris	trisz-(hidroximetil)-aminometán

Köszönetnyilvánítás

Ezúton szeretnék köszönetet mondani mindenekelőtt témavezetőimnek: Dr. Ovádi Juditnak, aki lehetővé tette, hogy munkacsoportjában dolgozhassak és szakdolgozatomat az általa vezetett laborban készítsem el, valamint irányította munkámat, és sok hasznos információval és útmutatással látott el, valamint Dr. Oláh Juditnak, aki mindig türelemmel, figyelemmel kísérte munkámat. Sok mérési technika, laboratóriumi vizsgálati módszer elsajátítását neki köszönhetem.

Köszönettel tartozom a csoport további tagjainak: Szunyogh Sándornak, Lehotzky Attilának és Orosz Ferencnek, hogy kérdéseimre mindig készségesen válaszoltak, problémáim megoldásában segítettek.

Hozzájárulásuk nélkül nem sikerült volna elkészítenem munkámat.