

KÖSZÖNETNYILVÁNÍTÁS

Szeretném megköszönni témavezetőmnek, Dr. Nyiri Gábornak a munkámhoz nyújtott rengeteg segítséget és biztatást. Szintén köszönettel tartozom munkatársaimnak, elsősorban Dr. Szőnyi Andrásnak, valamint Dr. Cserép Csabának, Dr. Takács Virágnak, Sós Katalinnak, Pósfai Balázsnak és Hegedüs Pannának. Köszönöm a technikai segítséget Lengyel Katalinnak, Szépné Simon Emőkének, Iványi Katalinnak és Goda Győzőnek. Nem utolsó sorban köszönöm Prof. Freund Tamás Igazgató Úrnak a megteremtett körülményeket és lehetőséget, hogy az általa vezetett Agykéreg Kutatócsoportban végezhetem diákköri munkám.

IRODALOMJEGYZÉK

- Abercrombie M. (1946) Estimation of nuclear population from microtome sections. *Anat Rec* **94**, 239-47.
- Acsády L., Arabadzisz D., Katona I. & Freund T.F. (1996) Topographic distribution of dorsal and median raphe neurons with hippocampal, septal and dual projection. *Acta Biol Hung* **47**, 9-19.
- Acsády L., Halasy K. & Freund T.F. (1993) Calretinin is present in non-pyramidal cells of the rat hippocampus--III. Their inputs from the median raphe and medial septal nuclei. *Neuroscience* **52**, 829-41.
- Affaticati P., Mignen O., Jambou F., Potier M.C., Klingel-Schmitt I., Degrouard J., Peineau S., Gouadon E., Collingridge G.L., Liblau R., Capiod T. & Cohen-Kaminsky S. (2011) Sustained calcium signalling and caspase-3 activation involve NMDA receptors in thymocytes in contact with dendritic cells. *Cell Death Differ* **18**, 99-108.
- Amilhon B., Lepicard E., Renoir T., Mongeau R., Popa D., Poirel O., Miot S., Gras C., Gardier A.M., Gallego J., Hamon M., Lanfumey L., Gasnier B., Giros B. & El Mestikawy S. (2010) VGLUT3 (vesicular glutamate transporter type 3) contribution to the regulation of serotonergic transmission and anxiety. *J Neurosci* **30**, 2198-210.
- Aznar S., Qian Z.X. & Knudsen G.M. (2004) Non-serotonergic dorsal and median raphe projection onto parvalbumin- and calbindin-containing neurons in hippocampus and septum. *Neuroscience* **124**, 573-81.
- Boyce R.W., Dorph-Petersen K.A., Lyck L. & Gundersen H.J. (2010) Design-based stereology: introduction to basic concepts and practical approaches for estimation of cell number. *Toxicol Pathol* **38**, 1011-25.
- Colom L.V. (2006) Septal networks: relevance to theta rhythm, epilepsy and Alzheimer's disease. *J Neurochem* **96**, 609-23.
- Colom L.V., Castaneda M.T., Reyna T., Hernandez S. & Garrido-Sanabria E. (2005) Characterization of medial septal glutamatergic neurons and their projection to the hippocampus. *Synapse* **58**, 151-64.
- Dahlström A. & Fuxe K. (1964) Localization of monoamines in the lower brain stem. *Experientia* **20**, 398-9.
- Euston D.R., Gruber A.J. & McNaughton B.L. (2012) The role of medial prefrontal cortex in memory and decision making. *Neuron* **76**, 1057-70.

- Freneau R.T., Burman J., Qureshi T., Tran C.H., Proctor J., Johnson J., Zhang H., Sulzer D., Copenhagen D.R., Storm-Mathisen J., Reimer R.J., Chaudhry F.A. & Edwards R.H. (2002) The identification of vesicular glutamate transporter 3 suggests novel modes of signaling by glutamate. *Proc Natl Acad Sci U S A* **99**, 14488-93.
- Freund T.F. (1989) GABAergic septohippocampal neurons contain parvalbumin. *Brain Res* **478**, 375-81.
- Freund T.F. & Antal M. (1988) GABA-containing neurons in the septum control inhibitory interneurons in the hippocampus. *Nature* **336**, 170-3.
- Freund T.F. & Buzsáki G. (1996) Interneurons of the hippocampus. *Hippocampus* **6**, 347-470.
- Freund T.F., Gulyás A.I., Acsády L., Görcs T. & Tóth K. (1990) Serotonergic control of the hippocampus via local inhibitory interneurons. *Proc Natl Acad Sci U S A* **87**, 8501-5.
- Gras C., Herzog E., Bellenchi G.C., Bernard V., Ravassard P., Pohl M., Gasnier B., Giros B. & El Mestikawy S. (2002) A third vesicular glutamate transporter expressed by cholinergic and serotonergic neurons. *J Neurosci* **22**, 5442-51.
- Halasy K., Miettinen R., Szabat E. & Freund T.F. (1992) GABAergic Interneurons are the Major Postsynaptic Targets of Median Raphe Afferents in the Rat Dentate Gyrus. *Eur J Neurosci* **4**, 144-53.
- Heidbreder C.A. & Groenewegen H.J. (2003) The medial prefrontal cortex in the rat: evidence for a dorso-ventral distinction based upon functional and anatomical characteristics. *Neurosci Biobehav Rev* **27**, 555-79.
- Hensler J.G. (2006) Serotonergic modulation of the limbic system. *Neurosci Biobehav Rev* **30**, 203-14.
- Jackson J., Bland B.H. & Antle M.C. (2009) Nonserotonergic projection neurons in the midbrain raphe nuclei contain the vesicular glutamate transporter VGLUT3. *Synapse* **63**, 31-41.
- Klausberger T. & Somogyi P. (2008) Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science* **321**, 53-7.
- Kosofsky B.E. & Molliver M.E. (1987) The serotonergic innervation of cerebral cortex: different classes of axon terminals arise from dorsal and median raphe nuclei. *Synapse* **1**, 153-68.
- Köhler C., Chan-Palay V., Haglund L. & Steinbusch H. (1980) Immunohistochemical localization of serotonin nerve terminals in the lateral entorhinal cortex of the rat: demonstration of two separate patterns of innervation from the midbrain raphe. *Anat Embryol (Berl)* **160**, 121-9.

- Köhler C., Chan-Palay V. & Steinbusch H. (1982) The distribution and origin of serotonin-containing fibers in the septal area: a combined immunohistochemical and fluorescent retrograde tracing study in the rat. *J Comp Neurol* **209**, 91-111.
- Köhler C. & Steinbusch H. (1982) Identification of serotonin and non-serotonin-containing neurons of the mid-brain raphe projecting to the entorhinal area and the hippocampal formation. A combined immunohistochemical and fluorescent retrograde tracing study in the rat brain. *Neuroscience* **7**, 951-75.
- Leranth C. & Vertes R.P. (1999) Median raphe serotonergic innervation of medial septum/diagonal band of Broca (MSDB) parvalbumin-containing neurons: possible involvement of the MSDB in the desynchronization of the hippocampal EEG. *J Comp Neurol* **410**, 586-98.
- Mamounas L.A., Mullen C.A., O'Hearn E. & Molliver M.E. (1991) Dual serotonergic projections to forebrain in the rat: morphologically distinct 5-HT axon terminals exhibit differential vulnerability to neurotoxic amphetamine derivatives. *J Comp Neurol* **314**, 558-86.
- McKenna J.T. & Vertes R.P. (2001) Collateral projections from the median raphe nucleus to the medial septum and hippocampus. *Brain Res Bull* **54**, 619-30.
- Mesulam M.M., Mufson E.J., Wainer B.H. & Levey A.I. (1983) Central cholinergic pathways in the rat: an overview based on an alternative nomenclature (Ch1-Ch6). *Neuroscience* **10**, 1185-201.
- Morales M. & Bloom F.E. (1997) The 5-HT₃ receptor is present in different subpopulations of GABAergic neurons in the rat telencephalon. *J Neurosci* **17**, 3157-67.
- Paxinos G. & Franklin Keith B. J. The mouse brain in stereotaxic coordinates. (Academic press, 2004).
- Ropert N. & Guy N. (1991) Serotonin facilitates GABAergic transmission in the CA1 region of rat hippocampus in vitro. *J Physiol* **441**, 121-36.
- Schmitz C. & Hof P.R. (2005) Design-based stereology in neuroscience. *Neuroscience* **130**, 813-31.
- Shutoh F., Ina A., Yoshida S., Konno J. & Hisano S. (2008) Two distinct subtypes of serotonergic fibers classified by co-expression with vesicular glutamate transporter 3 in rat forebrain. *Neurosci Lett* **432**, 132-6.
- Somogyi J., Baude A., Omori Y., Shimizu H., El Mestikawy S., Fukaya M., Shigemoto R., Watanabe M. & Somogyi P. (2004) GABAergic basket cells expressing

- cholecystokinin contain vesicular glutamate transporter type 3 (VGLUT3) in their synaptic terminals in hippocampus and isocortex of the rat. *Eur J Neurosci* **19**, 552-69.
- Swanson L.W. & Cowan W.M. (1979) The connections of the septal region in the rat. *J Comp Neurol* **186**, 621-55.
- Szabadics J., Varga C., Molnar G., Olah S., Barzo P. & Tamas G. (2006) Excitatory effect of GABAergic axo-axonic cells in cortical microcircuits. *Science* **311**, 233-5.
- Szabadits E., Cserép C., Szonyi A., Fukazawa Y., Shigemoto R., Watanabe M., Itohara S., Freund T.F. & Nyiri G. (2011) NMDA receptors in hippocampal GABAergic synapses and their role in nitric oxide signaling. *J Neurosci* **31**, 5893-904.
- Szentágothai J. (1975) The 'module-concept' in cerebral cortex architecture. *Brain Res* **95**, 475-96.
- Szonyi A., Mayer M.I., Cserép C., Takács V.T., Watanabe M., Freund T.F. & Nyiri G. (2014) The ascending median raphe projections are mainly glutamatergic in the mouse forebrain. *Brain Structure and Function* 2014 Nov 9 [Epub ahead of print].
- Tschanz S., Schneider J.P. & Knudsen L. (2014) Design-based stereology: Planning, volumetry and sampling are crucial steps for a successful study. *Ann Anat* **196**, 3-11.
- Varga V., Losonczy A., Zemelman B.V., Borhegyi Z., Nyiri G., Domonkos A., Hangya B., Holderith N., Magee J.C. & Freund T.F. (2009) Fast synaptic subcortical control of hippocampal circuits. *Science* **326**, 449-53.
- Vertes R.P., Fortin W.J. & Crane A.M. (1999) Projections of the median raphe nucleus in the rat. *J Comp Neurol* **407**, 555-82.
- Vertes R.P. & Kocsis B. (1997) Brainstem-diencephalo-septohippocampal systems controlling the theta rhythm of the hippocampus. *Neuroscience* **81**, 893-926.
- West M.J., Slomianka L. & Gundersen H.J. (1991) Unbiased stereological estimation of the total number of neurons in the subdivisions of the rat hippocampus using the optical fractionator. *Anat Rec* **231**, 482-97.
- Zhao Z.Q., Chiechio S., Sun Y.G., Zhang K.H., Zhao C.S., Scott M., Johnson R.L., Deneris E.S., Renner K.J., Gereau R.W. & Chen Z.F. (2007) Mice lacking central serotonergic neurons show enhanced inflammatory pain and an impaired analgesic response to antidepressant drugs. *J Neurosci* **27**, 6045-53.

Tételes szerzői hozzájárulás

Alulírott **Mayer Márton István** a dolgozatban leírt munkafolyamatok közül a:

- sztereotaxiás műtétek és
- retrográd jelölőanyag beadások
- állatok perfúziója
- az agyak vibrotómos felmetszése
- immunfluoreszcens reakciók
- képkötés konfokális mikroszkóppal
- sztereológiai sejtszámolások és
- retrográd pályajelölés kiértékelése
- adatelemzés

végeztem el.

Budapest, 2014. december 11.



aláírás