8. *Irodalomjegyzék*

Adams F. (ed.). The Exant Works of Aretaeus, the Cappadocian. The Sydenham Society, London, pp 52-53. (1856)

Akerman S., Williamson D. J., Kaube H., Goadsby P. J. Nitric oxide synthase inhibitors can antagonize neurogenic and calcitonin gene-related peptide induced dilation of dural meningeal vessels. British Journal of Pharmacology; 137:62-68. (2002)

Andó R. D., Méhész B., Gyires K., Illes P., Sperlágh B. A comparative analysis of the activity of ligands acting at P2X and P2Y receptor subtypes in models of neuropathic, acute and inflammatory pain. Br J Pharmacol; 159(5):1106-17. (2010)

Armstrong J. N., Brust T. B., Lewis R. G., MacVicar B. A. Activation of presynaptic P2X7-like receptors depresses mossy fiber-CA3 synaptic transmission through p38 mitogen-activated protein kinase. J Neurosci.; 22(14):5938-45. (2002)

Ashina M., Brendtsen L., Jensen R., Schifter S., Olesen J. Evidence for increased plasma levels of calcitonin gene-related peptide in migraine outside of attacks. Pain; 86:133-138. (2000)

Bánk J., Márta M. Hungarian migraine epidemiology. Headache; 40:164-169. (2000)

Bianchi B. R., Lynch K. J., Touma E., Niforatos W., Burgard E. C., Alexander K. M., Park H. S., Yu H., Metzger R., Kowaluk E., Jarvis M. F., van Biesen T. Pharmacological characterization of recombinant human and rat P2X receptor subtypes. Eur J Pharmacol; 376 (1-2): 127-138. (1999)

Bigal M. E., Lipton R. B., Krymchantowski A. V. The medical management of migraine. Am J Ther.; 11:130-40. (2004)

Braughler J. M., Mittal C. K., Murad F. Effects of thiols, sugars, and proteins on nitric oxide activation of guanylate cyclase. J Biol Chem.; 254(24):12450-4. (1979)

Bulanova E., Budagian V., Orinska Z., Hein M., Peteresen F., Thon L., Adam D., Bulfone-Paus S. Extracellular ATP induces cytokine expression and apoptosis through P2X7 receptor in murine mast cells. J Immunol; 174: 3880-3890. (2005)

Burstein R., Collins B., Jakubowski M. Defeating migraine pain with triptans: a race against the development of cutaneous allodynia. Ann Neurol; 55 (1):19-26. (2004)

Burstein R., Jakubowski M. Clinical implications of periferal and central sensitization during migraine. Pain. An Updated Review: Refresher Course Syllabus, ed. Giamberardino M. A., IASP, Press, Seattle. (2002)

Buzzi M. G., Bonamini M., Moskowitz M. A. Neurogenic model of migraine. Cephalalgia.; 15(4):277-80. (1995)

Buzzi M. G., Carter W. B., Shimizu T., Heath H. 3rd, Moskowitz M. A. Dihydergotamine and sumatriptan attenuate concentrations of CGRP in plasma in rat superior sagittal sinus during electrical stimulation of trigeminal ganglion. Neuropharmacology; 30:1193-1200. (1991)

Chen Y., Zhang X., Wang C., Li G., Gu Y., Huang L. Y. Activation of P2X7 receptors in glial satellite cells reduces pain through downregulation of P2X3 receptors in nociceptive neurons. Proc Natl Acad Sci; (USA) 105(43):16773-8. (2008)

Chessell I. P., Hatcher J. P., Bountra C., Michel A. D., Hughes J.P., Green P., Egerton J., Murfin M., Richardson J., Peck W. L., Grahames C. B., Casula M. A., Yiangou Y., Birch R., Anand P., Buell G. N. Disruption of the P2X7 purinoceptor gene abolishes chronic inflammatiory and neuropathic pain. Pain; 114: 386-396. (2005)

Classification and diagnostic criteria for headache disorders, cranial neuralgias and facial pain. 2nd ed. Headache Classification Committee of the International Headache Society. Cephalalgia; Suppl.1. 160 p. (2004)

Codocedo J. F., Godoy J. A., Poblete M. I., Inestrosa N. C., Huidobro-Toro J. P. ATP induces NO production in hippocampal neurons by P2X(7) receptor activation independent of glutamate signaling. PLoS One.; 8(3):e57626. (2013)

Costa C., Tozzi A., Rainero I., Cupini L. M., Calabresi P., Ayata C., Sarchielli P. Cortical spreading depression as a target for anti-migraine agents. J Headache Pain.; 23;14:62. (2013)

Deli T. Purinerg jelátvitel a sejtek differenciálódásában és malignus transzformációjában: a P2X7 receptor kitüntetett szerepe. (Purinergic signalling in the differentiation and malignant transformation of cells: the the special role of the P2X7 receptor.). Ph.D. dissertation, University of Debrecen, Hungary. (2006)

Dell'Antonio G., Quattrini A., Dal Cin E., Fulgenzi A., Ferrero M. E. Antinociceptive effect of a new P(2Z)/P2X7 antagonist, oxidized ATP, in arthritic rats. Neurosci Lett.; 327(2):87-90. (2002)

Denlinger L. C., Sommer J. A., Parker K., Gudipaty L., Fisette P. L., Watters J. W., Proctor R. A., Dubyak G. R., Bertics P. J. Mutation of a dibasic amino acid motif within the C terminus of the P2X7 nucleotide receptor results in trafficking defects and impaired function. J Immunol; 171 (3): 1304-1311. (2003)

Donnelly-Roberts D. L., Jarvis M.F. Discovery of P2X7 receptor-selective antagonists offers new insights into P2X7 receptor function and indicates a role in chronic pain states. Br J Pharmacol.; 151(5):571-9. (2007)

Fozard J. R., Kalkman H. O. Hydroxytryptamine (5-HT) and the initiation of migraine: new perspectives. Naunyn-Schmiedebergs Archives of Pharmacology; 350:225-229. (1994)

Franka G. Bradford Assay Performed on BMG LABTECH´s FLUOstar Omega with new Evaluation Software. Application Note; (E. J. Dell, BMG LABTECH, Durham, USA) 158 Rev. (2007)

Franke H., Grosche J., Schadlich H., Krügel U., Allgaier C., Illes P. P2X receptor expression on astrocytes in the nucleus accumbens of rats. Neuroscience; 108 (3): 421-429. (2001)

Friberg L., Olesen J., Iversen H. K., Sperling B. Migraine pain associadted with middle cerebral artery dilatation: reversal by sumatriptan. Lancet; 338:13-17. (1991)

Friedle S. A., Curet M. A., Watters J. J.. Recent Patents on Novel $P2X_7$ Receptor Antagonists and Their Potential for Reducing Central Nervous System Inflammation. Recent Pat CNS Drug Discov.; 5(1): 35–45. (2010)

Gartland A., Skarratt K. K., Hocking L. J., Parsons C., Stokes L., Jørgensen N. R., Fraser W. D., Reid D. M., Gallagher J. A., Wiley J. S. Polymorphisms in the P2X7 receptor gene are

associated with low lumbar spine bone mineral density and accelerated bone loss in post-menopausal women. Eur J Hum Genet; 20(5):559-64. (2012)

Goadsby P. J. Migraine, allodynia, sensitisation and all of that ... Eur Neurol.; 53 Suppl 1:10-6. Epub. (2005)

Goadsby P. J., Edvinsson L. The trigeminovascular system and migraine: studies characterizing cerebrovascular and neuropeptide changes seen in humans and cats. Ann Neurol; 33:48-56. (1993)

Goadsby P. J., Hoskin K. L. The distribution of trigeminovascular afferents in the nonhuman primate brain Macaca nemestrina: a c-fos immunocytochemical study. Journal of Anatomy; 190:367-375, 1997

Gölöncsér F., Sperlágh B. Effect of genetic deletion and pharmacological antagonism of P2X7 receptors in a mouse animal model of migraine. J Headache Pain. 15:24. (2014)

Guseo A., Giczi J. Epidemiology of migraine in urban and rural communities in central Transdanubia, Hungary. In: Headache Classification and Epidemiology. Ed.: Olesen J., New York, Raven Press; pp. 263-265. (1994)

Hadjikhani N., Sanchez Del Rio M., Wu O., Schwartz D., Bakker D., Fischl B., Kwong K. K., Cutrer F. M., Rosen B. R., Tootell R. B., Sorensen A. G., Moskowitz M. A. Mechanisms of migraine aura revealed by functional MRI in human visual cortex. Proc Natl Acad Sci, (USA); 98 (8):4687-92. (2001)

Hargreaves R. J., Shepheard S. L. Pathophysiology of migraine--new insights. Can J Neurol Sci. Suppl 3:S12-9. (1999)

He W. J., Cui J., Du L., Zhao Y. D., Burnstock G., Zhou H. D., Ruan H. Z. Spinal P2X(7) receptor mediates microglia activation-induced neuropathic pain in the sciatic nerve injury rat model. Behav Brain Res.; 226(1):163-70. (2012)

Hide I., Tanaka M., Inoue A., Nakajima K., Kohsaka S., Inoue K., Nakata Y. Extracellular ATP triggers tumor necrosis factor-alpha release from rat microglia. J Neurochem; 75 (3): 965-972. (2000)

Itoh K., Chiang C. Y. Central sensitization of nociceptive neurons in rat medullary dorsal horn involves purinergic P2X7 receptors. Neuroscience; 192, 721–731. (2011)

Iversen H. K. Human migraine models. Cephalalgia; 21:781-5. (2001)

Iversen H. K., Olesen J. Headache induced by a nitric oxide donor (nitroglycerin) responds to sumatriptan. A human model for development of migraine drugs. Cephalalgia; 16(6):412-8. (1996)

Jane P. Hughes, Jonathan P. Hatcher, Iain P. Chessell. The role of $P2X_7$ in pain and inflammation. Purinergic Signal.; 3(1-2): 163–169. (2007)

Jiang L. H., Mackenzie A. B., North R. A., Surprenant A. Brilliant blue G selectively blocks ATP-gated rat P2X(7) receptors. Mol Pharmacol; 58(1):82-8. (2000)

Keidel M. A migrén. Okok, fajták, terápiák. Corvina Kiadó Kft. ISBN 978 963 13 5776 9. (2009)

Kim M., Jiang L. H., Wilson H. L., North R. A., Surprenant A. Proteomic and functional evidence for a P2X7 receptor signalling complex. EMBO J 20 (22): 6347-58. (2001)

Knyihár-Csillik E., Tajti J. Electrical stimulation of the Gasserian ganglion induces structural alterations of calcitonin generelated peptide-immunoreactive perivascular sensory nerve terminals in the rat cerebral duramater: a possible model of migraine headache. Neurosci. Lett; 184, 189–192. (1995)

Knyihár-Csillik E., Tajti J., Csillik A. E., Chadaide Z., Mihály A., Vécsei L. Effects of eletriptan on the peptidergic innervation of the cerebral dura mater and trigeminal ganglion, and on the expression of c-fos and c-jun in the trigeminal complex of the rat in an experimental migraine model. Eur J Neurosci; 12:3991-4002. (2000)

Knyihár-Csillik E., Tajti J., Samsam M., Sáry G., Slezak S., Vécsei L. Effect of a serotonin agonist (sumatriptan) on the peptidergic innervation of the rat cerebral dura mater and on the expression of c-fos in the caudal trigeminal nucleus in an experimental migraine model. J Neurosci; Res 48:449-464. (1997)

Knyihár-Csillik E., Toldi J., Mihály A., Krisztin-Peva B., Chadaide Z., Németh H., Fenyő R., Vécsei L. Kynurenine in combination with probenecid mitigates the stimulation-induced increase of c-Fos immunoreactivity of the rat caudal trigeminal nucleus in an experimental migraine model. J Neural Transm; 114(4): 417-21. (2007)

Kors E., Haan J., Ferrari M. Migraine genetics. Curr Pain Headache; Rep 7:212-217. (2003)

Lam H. H., Hanley D.F., Trapp B. D., Saito S., Raja S., Dawson T. M., Yamaguchi H. Induction of spinal cord neuronal nitric oxide synthase (NOS) after formalin injection in the rat hind paw. Neurosci Lett; 210(3):201-4. (1996)

Larmande P., Ligeard P., Maillot F., Belin C. Migraine attacks triggered by food. Med. Nut.; 29 (6): 263-267. (1993)

Larmande P., Ligeard P., Maillot F., Belin C. Wine: migraine triggering factor. Med. Nut.; 30 (4): 205-206. (1994)

Lassen L. H., Ashina M., Christiansen I., Ulrich V., Olesen J. Nitric oxide synthase inhibition in migraine. Lancet; 349(9049):401-2. (1997)

Leao A. A. P. Spreading depression of activity in cerebral cortex. J Neurophysiol; 7:359-390. (1944)

Lord B., Aluisio L., Shoblock J. R., Neff R. A., Varlinskaya E. I., Ceusters M., Lovenberg T. W., Carruthers N., Bonaventure P., Letavic M. A., Deak T., Drinkenburg W., Bhattacharya A. Pharmacology of a Novel Central Nervous System-Penetrant P2X7 Antagonist JNJ-42253432. J Pharmacol Exp Ther.; 351(3):628-41. (2014)

Lyons A., Petrucelli R. J. Medicine: An Illustrated History. New York, Abrams Inc. ISBN 10: 0810980800. (1987)

Markowitz S., Saito K., Moskowitz M. A. Neurogenically mediated leakage of plasma protein occurs from blood vessels in dura mater but not brain. J Neurosci; 7 (12):4129-4136. (1987)

Moskowitz M. A. Defining a pathway to discovery from bench to beside: the trigeminovascular system and sensitization. Headache; 48 (5): 688-90. (2008)

Moskowitz M. A. Genes, proteases, cortical spreading depression and migraine: impact on pathophysiology and treatment. Funct Neurol; 22(3):133-6. (2007)

Moskowitz M. A. Neurogenic inflammation in the pathophysiology and treatment of migraine. Neurology; 43:16-20. (1993)

Murad F. The nitric oxide-cyclic GMP signal transduction system for intracellular and intercellular communication. Recent Progress in Hormone Research; 43:109-142. (1994)

Nathan C., Xie Q. W. Regulation of biosynthesis of nitric oxide. J Biol Chem.; 269(19):13725-8. (1994)

Nelson D. W., Gregg R. J., Kort M. E., Perez-Medrano A., Voight E. A., Wang Y., Grayson G., Namovic M. T., Donnelly-Roberts D. L., Niforatos W., Honore P., Jarvis M. F., Faltynek C. R., Carroll W. A. Structure-activity relationship studies on a series of novel, substituted 1-benzyl-5-phenyltetrazole P2X7 antagonists. J Med Chem.; 49(12):3659-66. (2006)

North R. A. Molecular physiology of P2X receptors. Physiol; Rev 82: 1013–1067. (2002)

Olesen J., Friberg L., Olsen T. S., Iversen H. K., Lassen N. A., Andersen A. R., Karle A. Timing and topography of cerebral blood flow, aura, and headache during migraine attacks. Ann Neurol.;(6):791-8. (1990)

Olesen J., Iversen H. K., Thomsen L. L. Nitric oxide supersensitivity. A possible mechanism of migraine pain. Neuroreport; 4:1027-103. (1993)

Párdutz A., Krizbai I., Multon S., Vécsei L., Schoenen J. Systemic nitroglycerin increases nNOS levels in rat trigeminal nucleus caudalis. Neuroreport; 11(14):3071-5. (2000)

Parvathenani L. K., Tertyshnikova S., Greco C. R., Roberts S. B., Robertson B., Posmantur R. P2X7 mediates superoxide production in primary microglia and is up-regulated in a transgenic mouse model of Alzheimer's disease. J Biol Chem; 278: 13309–13317. (2003)

Peng W., Cotrina M. L., Han X., Yu H., Bekar L., Blum L., Takano T., Tian G. F., Goldman S. A., Nedergaard M. Systemic administration of an antagonist of the ATP-sensitive

receptor P2X7 improves recovery after spinal cord injury. Proc Natl Acad Sci (USA); 106(30):12489-93. (2009)

Ralevic V., Burnstock G. Receptors for purines and pyrimidines. Pharmacol; Rev.; 50(3):413-92. (1998)

Read S. J., Parsons A. A. Cortical spreading depression and migraine. In: Migraine and headache pathophysiology. Ed.: Edvinsson L., London, Martin Dunitz; pp. 81-92. (1999)

Robert W. P., Cutler M. D.: Fejfájás a Neurológia fejezetben. Scientific American Medicine. (1990)

Russell M. B. The genetics of migraine. Eur J Neurol; 4:2-7. (1997)

Samsam M., Coveñas R., Ahangari R., Yajeya J., Narváez J. A., Tramu G. Alterations in neurokinin A-, substance P- and calcitonin gene-related peptide immunoreactivities in the caudal trigeminal nucleus of the rat following electrical stimulation of the trigeminal ganglion. Neurosci Lett; 261(3):179-82. (1999)

Sandkühler J. Models and mechanisms of hyperalgesia and allodynia. Physiol Rev.; 89(2):707-58. (2009)

Saxena P. R. The pathogenesis and pharmacology of migraine. Reviews of Contemp Pharmacotherapeutic; 5: 259-269. (1994)

Schuh-Hofer S., Tayefeh M., Reuter U., Dirnagl U., Arnold G. Effects of Parecoxib on Plasma Protein Extravasation and c-Fos Expression in the Rat. Headache; 46(2):276-85. (2006)

Silberstein S. D. A nők és a migrén. Orvostovábbképző Szemle; 2 (4): 123-129. (1995)

Sperlágh B., Kőfalvi A., Deuchars J., Atkinson L., Milligan C. J., Buckley N. J., Vizi E. S. Involvement of P2X7 receptors in the regulation of neurotransmitter release in the rat hippocampus. J Neurochem; 81: 1196–1211. (2002)

Sperlágh B., Szabó G., Erdélyi F., Baranyi M., Vizi E. S. Homo- and hetero-P2X Receptors Regulate Hippocampal Noradrenaline Release exchange of adenine nucleotides and

nucleosides in rat hippocampal slices by the nucleoside transport system. Br J Pharmacol; 139: 623–633. (2003)

Tajti J., Szok D., Párdutz Á., Tuka B., Csáti A., Kuris A., Toldi J., Vécsei L. Where does a migraine attack originate? In the brainstem. J Neural Transm.; 119(5):557-68. (2012)

Takemura M., Shimada T., Sugiyo S., Nokubi T., Shigenaga Y. Mapping of c-Fos in the trigeminal sensory nucleus following high- and low-intensity afferent stimulation in the rat. Exp Brain; Res.; 130(2):113-23. (2000)

Tassorelli C., Joseph S. A. Systemic nitroglicerin induces Fos immunreactivity in brainstem and forebrain structures of the rat. Brain; Res.; 682(1-2): 325-8. (1995)

Taylor S. Anxiety sensitivity: theoretical perspectives and recent findings. Behav Res Ther; 33:243-58. (1995)

Ter Horst G. J., Meijler W. J., Korf J., Kemper R. H. Trigeminal nociception-induced cerebral Fos expression in the conscious rat. Cephalalgia.; 21(10):963-75. (2001)

The World health report : 2001 : Mental health : new understanding, new hope. WHO Library Cataloguing in Publication Data. ISBN 92 4 156201 3. (2001)

Waeber C., Moskowitz M. A. Therapeutic implications of central and peripheral neuroligic mechanisms in migraine. Neurology 61 (Suppl. 4), S. S9 – S20. (2003)

Weber R. W., Vaughan T. R. Food and migraine headache. Immun. Allergy Clin. North Am.; 11: 4, 831-841. (1991)

Weiller C., May A., Limmroth V., Juptner M., Kaube H., Schayck R. V., Coenen H. H., Diener H. C. Brain stem activation in spontaneous human migraine attacks. Nature Medicine; 1(7): 658-60. (1995)

Worthington R. A., Smart M. L., Gu B. J., Williams D. A., Petron S., Wiley J. S., Barden J. A. Point mutations confer loss of ATP-induced human P2X (7) receptor function. FEBS Lett; 512 (1-3): 43-46. (2002)

9. ***Köszönetnyilvánítás***

Ezúton szeretnék köszönetet mondani Prof. Dr. Vécsei Lászlónak, hogy kísérleti munkámhoz lehetőséget és a szükséges feltételeket biztosította.

Köszönettel tartozom témavezetőimnek Dr. Párdutz Árpádnak és Bohár Zsuzsannának a szakmai irányításért, hasznos tanácsokért és a kitartó türelemért.

Belső konzulensemnek, Dr. Kis Zsoltnak köszönöm a segítségét.

Köszönet illeti Nagy-Grócz Gábort, Fejes-Szabó Annamáriát és Vékonyné Széll Valériát, akik a kísérleti eljárások elsajátításában hasznos útmutatással láttak el és esetleges kérdéseimmel is mindig fordulhattam hozzájuk.

Végül köszönettel és hálával tartozom páromnak, családomnak és barátaimnak a türelmükért és bíztatásukért.

10. Tételes szakmai hozzájárulás

Alulírott Szabadi Nikolett a dolgozatban leírt munkafolyamatok közül a:

- patkány intravénás kezelését
- anesztéziát
- koponya feltárást, majd Gasser-dúc ingerlést
- transzkardiális perfundálást követő agy, cervicalis gerincvelő és Gasser-dúc preparálását
- TNC metszetek készítését
- c-Fos-ra és nNOS-ra immunhisztokémiai festést

végeztem el.

.....

aláírás

Szeged, 2014. november 12.