

Irodalomjegyzék

- [1] R. C. Lee, R. L. Feinbaum, and V. Ambros, “The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*,” *Cell*, vol. 75 5, pp. 843–854, 1993.
- [2] M. Ghildiyal and P. D. Zamore, “Small silencing RNAs: an expanding universe,” *Nature Reviews Genetics*, vol. 10, no. 2, pp. 94–108, 2009.
- [3] R. C. Lee and V. Ambros, “An extensive class of small RNAs in *Caenorhabditis elegans*,” *Science*, vol. 294, no. 5543, pp. 862–864, 2001.
- [4] miRBase, Release 21: <http://www.mirbase.org>
- [5] Stefan L. Ameres and Phillip D. Zamore, „Diversifying microRNA sequence and function,” *Nature Reviews Molecular Cell Biology*, 14, 475–488, 2013.
- [6] K. Miyoshi, T. Miyoshi, and H. Siomi, “Many ways to generate microRNA-like small RNAs: non-canonical pathways for microRNA production,” *Molecular genetics and genomics MGG*, vol. 284, no. 2, pp. 95–103, 2010.
- [7] Y. Lee, M. Kim, J. Han, K.-H. Yeom, S. Lee, S. H. Baek, and V. N. Kim, “MicroRNA genes are transcribed by RNA polymerase II,” *the The European Molecular Biology Organization Journal*, vol. 23, no. 20, pp. 4051–4060, 2004.
- [8] J. Han, Y. Lee, K. Yeom, Y. Kim, H. Jin, and V. N. Kim, “The Drosha – DGCR8 complex in primary microRNA processing,” *Genes & Development*, pp. 3016–3027, 2004.
- [9] M. T. Bohnsack, K. Czapinski, and D. Gorlich, “Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs,” *Rna New York Ny*, vol. 10, no. 2, pp. 185–191, 2004.
- [10] V. N. Kim, J. Han, and M. C. Siomi, “Biogenesis of small RNAs in animals,” *Nature Reviews Molecular Cell Biology*, vol. 10, no. 2, pp. 126–139, 2009.
- [11] K. Okamura, M. D. Phillips, D. M. Tyler, H. Duan, Y. Chou, and E. C. Lai, “The regulatory activity of microRNA* species has substantial influence on microRNA and 3’ UTR evolution,” *Nature Structural & Molecular Biology*, vol. 15, no. 4, pp. 354–363, 2008.
- [12] A. Kozomara and S. Griffiths-Jones, “miRBase: integrating microRNA annotation and deep-sequencing data,” *Nucleic Acids Research*, vol. 39, no. Database issue, pp. D152–

7, 2011.

- [13] J.-S. Yang and Lai, "Alternative miRNA Biogenesis Pathways and the Interpretation of Core miRNA Pathway Mutants," *Molecular Cell*, vol. 43, no. 6, pp. 892–903, 2011.
- [14] J. G. Ruby, C. H. Jan, and D. P. Bartel, "Intronic microRNA precursors that bypass Drosha processing.," *Nature*, vol. 448, no. 7149, pp. 83–86, 2007.
- [15] K. Okamura, J. W. Hagen, H. Duan, D. M. Tyler, and E. C. Lai, "The mirtron pathway generates microRNA-class regulatory RNAs in Drosophila.," *Cell*, vol. 130, no. 1, pp. 89–100, 2007.
- [16] J. E. Babiarz, J. G. Ruby, Y. Wang, D. P. Bartel, and R. Blelloch, "Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, Dicer-dependent small RNAs," *Genes & Development*, vol. 22, no. 20, pp. 2773–2785, 2008.
- [17] J. O. Westholm and E. C. Lai, "Mirtrons: microRNA biogenesis via splicing.," *Biochimie*, vol. 93, no. 11, pp. 1897–904, 2011.
- [18] E. Berezhikov, W.-J. Chung, J. Willis, E. Cuppen, and E. C. Lai, "Mammalian mirtron genes.," *Molecular Cell*, vol. 28, no. 2, pp. 328–336, 2007.
- [19] C. R. Sibley, Y. Seow, S. Saayman, K. K. Dijkstra, S. El Andaloussi, M. S. Weinberg, and M. J. A. Wood, "The biogenesis and characterization of mammalian microRNAs of mirtron origin," *Nucleic Acids Research*, vol. 40, no. 1, pp. 1–11, 2011.
- [20] A. Schamberger, B. Sarkadi, and T. I. Orban, "Human mirtrons can express functional microRNAs simultaneously from both arms in a flanking exon-independent manner," *RNA Biology*, vol. 9, no. 9, pp. 1177–1185, Sep. 2012.
- [21] A. S. Flynt, J. C. Greimann, W.-J. Chung, C. D. Lima, and E. C. Lai, "MicroRNA biogenesis via splicing and exosome-mediated trimming in Drosophila.," *Molecular Cell*, vol. 38, no. 6, pp. 900–907, 2010.
- [22] J. Liu, M. A. Carmell, F. V Rivas, C. G. Marsden, J. M. Thomson, J.-J. Song, S. M. Hammond, L. Joshua-Tor, and G. J. Hannon, "Argonaute2 is the catalytic engine of mammalian RNAi.," *Science*, vol. 305, no. 5689, pp. 1437–1441, 2004.
- [23] E. C. Lai, "Micro RNAs are complementary to 3' UTR sequence motifs that mediate negative post-transcriptional regulation.," *Nature Genetics*, vol. 30, no. 4, pp. 363–364, 2002.
- [24] S. Yekta, I.-H. Shih, and D. P. Bartel, "MicroRNA-directed cleavage of HOXB8 mRNA.," *Science*, vol. 304, no. 5670, pp. 594–596, 2004.
- [25] Megraw,M., Sethupathy,P., Corda,B. and Hatzigeorgiou,A.G., „miRGen: a database for

the study of animal microRNA genomic organization and function.,” *Nucleic Acids Res.*, 35, D149–D155., 2007.

- [26] Bentwich I, Avniel A, Karov Y, et al., „Identification of hundreds of conserved and nonconserved human microRNAs.,” *Nat Genet*, 37: 766–770., 2005
- [27] I. Flor and J. Bullerdiek, “The dark side of a success story: microRNAs of the C19MC cluster in human tumours.,” *J. Pathol.*, vol. 227, no. 3, pp. 270–4, Jul. 2012.
- [28] Zhang R, Wang YQ, Su B, „Molecular evolution of a primate- specific microRNA family.,” *Mol Biol Evol*, vol. 25, pp. 1493–1502, 2008.
- [29] Vergoulis, T. I. Vlachos, P. Alexiou, G. Georgakilas, M. Maragkakis, M. Reczko, S. Gerangelos, N. Koziris, T. Dalamagas, AG Hatzigeorgiou; „Tarbase 6.0: Capturing the Exponential Growth of miRNA Targets with Experimental Support.,” *Nucl. Acids Res.* 40 (D1): D222-D229., 2012.
- [30] Cao H, Yang CS, Rana TM., „Evolutionary emergence of microRNAs in human embryonic stem cells.,” *PLoS One*, 3: e2820., 2008.
- [31] M.-L. Bortolin-Cavaillé, M. Dance, M. Weber, and J. Cavaillé, “C19MC microRNAs are processed from introns of large Pol-II, non-protein-coding transcripts.,” *Nucleic Acids Res.*, vol. 37, no. 10, pp. 3464–73, Jun. 2009.
- [32] C. Bellemer, M.-L. Bortolin-Cavaillé, U. Schmidt, S. M. R. Jensen, J. Kjems, E. Bertrand, and J. Cavaillé, “Microprocessor dynamics and interactions at endogenous imprinted C19MC microRNA genes.,” *J. Cell Sci.*, vol. 125, no. Pt 11, pp. 2709–20, Jun. 2012.
- [33] M. Noguer-Dance, S. Abu-Amero, M. Al-Khtib, A. Lefèvre, P. Coullin, G. E. Moore, and J. Cavaillé, “The primate-specific microRNA gene cluster (C19MC) is imprinted in the placenta.,” *Hum. Mol. Genet.*, vol. 19, no. 18, pp. 3566–82, Sep. 2010.
- [34] Tsai KW, Kao HW, Chen HC, et al, „Epigenetic control of the expression of a primate-specific microRNA cluster in human cancer cells.,” *Epigenetics*, vol. 4, pp. 587–592, 2009.
- [35] Saito Y, Jones PA, „Epigenetic activation of tumor suppressor microRNAs in human cancer cells.,” *Cell Cycle*, vol. 5, pp. 2220–2222, 2006.
- [36] Y. Saito, H. Suzuki, H. Tsugawa, I. Nakagawa, J. Matsuzaki, Y. Kanai, and T. Hibi, “Chromatin remodeling at Alu repeats by epigenetic treatment activates silenced microRNA-512-5p with downregulation of Mcl-1 in human gastric cancer cells.,” *Oncogene*, vol. 28, no. 30, pp. 2738–44, Jul. 2009.

- [37] Wu, S., Huang, S., Ding, J., Zhao, Y., Liang, L., Liu, T., Zhan, R., and He, X, "Multiple microRNAs modulate p21Cip1/Waf1 expression by directly targeting its 3'untranslated region.," *Oncogene*, vol. 29, pp. 2302–2308, 2010.
- [38] S. M. Tan, R. Kirchner, J. Jin, O. Hofmann, L. McReynolds, W. Hide, and J. Lieberman, "Sequencing of Captive Target Transcripts Identifies the Network of Regulated Genes and Functions of Primate-Specific miR-522.," *Cell Rep.*, vol. 8, no. 4, pp. 1225–1239, Aug. 2014.
- [39] Keklikoglou I, Koerner C, Schmidt C, et al. MicroRNA-520/373 family functions as a tumor suppressor in estrogen receptor negative breast cancer by targeting NF-κB and TGFβ signaling pathways. *Oncogene* 2011; [Epub ahead of print 12 Dec 2011. DOI: 10.1038/onc.2011.571]
- [40] Lv SQ, Kim YH, Giulio F, et al, „Genetic alterations in microRNAs in medulloblastomas.," *Brain Pathology*, vol. 22, pp. 230–239, 2012.
- [41] Li M, Lee KF, Lu Y, et al, „Frequent amplification of a chr19q13.41 microRNA polycistron in aggressive primitive neuroectodermal brain tumors.," *Cancer Cell*, vol. 16, pp. 533–546, 2009.
- [42] Butz H, Liko I, Czirjak S, et al, "Down-regulation of Wee1 kinase by a specific subset of microRNA in human sporadic pituitary adenomas.," *J Clin Endocrinol Metab*, vol. 95, pp. 181–191, 2010.
- [43] Schepeler T, Reinert JT, Ostensfeld MS, et al, "Diagnostic and prognostic microRNAs in stage II colon cancer.," *Cancer Res*, vol. 68, pp. 6416–6424, 2008.
- [44] Fornari F, Milazzo M, Chieco P, et al, „In hepatocellular carcinoma miR-519d is upregulated by p53 and DNA hypomethylation and targets CDKN1A/p21, PTEN, AKT3 and TIMP2.," *J Pathol*, 2012.
- [45] Kim TH, Kim YK, Kwon Y, et al, "Deregulation of miR-519a, 153, and 485-5p and its clinicopathological relevance in ovarian epithelial tumours.," *Histopathology*, vol. 57, pp. 734–743, 2010.
- [46] Liu P, Wilson MJ, "miR-520c and miR-373 upregulate MMP9 expression by targeting mTOR and SIRT1, and activate the Ras/Raf/MEK/Erk signaling pathway and NF-κB factor in human fibrosarcoma cells.," *J Cell Physiol*, vol. 227, pp. 867–876, 2012.
- [47] Li X, Pan YZ, Seigel GM, et al, "Breast cancer resistance protein BCRP/ABCG2 regulatory microRNAs (hsa-miR-328, -519c and -520h) and their differential expression in stem-like ABCG2+ cancer cells.," *Biochem Pharmacol*, vol. 81, pp. 783–792, 2011.

- [48] Wang F, Xue X, Wei J, et al, “hsa-miR-520h downregulates ABCG2 in pancreatic cancer cells to inhibit migration, invasion, and side populations.,” *Br J Cancer*, vol. 103, pp. 567–574. 2010.
- [49] A. L. Kasinski and F. J. Slack, “Epigenetics and genetics. MicroRNAs en route to the clinic: progress in validating and targeting microRNAs for cancer therapy.,” *Nat. Rev. Cancer*, vol. 11, no. 12, pp. 849–64, Dec. 2011.
- [50] S. S. Jeffrey, “Cancer biomarker profiling with microRNAs.,” *Nat. Biotechnol.*, vol. 26, no. 4, pp. 400–1, Apr. 2008.
- [51] F. Chen, H. Zhu, L. Zhou, S. Wu, J. Wang, and Z. H. I. Chen, “Inhibition of c-FLIP expression by miR-512-3p contributes to Taxol-induced apoptosis in hepatocellular carcinoma cells,” pp. 1457–1462, 2010.
- [52] M. Petersen, C. B. Nielsen, K. E. Nielsen, G. A. Jensen, K. Bondensgaard, S. K. Singh, V. K. Rajwanshi, A. A. Koshkin, B. M. Dahl, J. Wengel, and J. P. Jacobsen, “The conformations of locked nucleic acids (LNA).,” *Journal of molecular recognition JMR*, vol. 13, no. 1, pp. 44–53, 2000.
- [53] C. R. Sibley, Y. Seow, H. Curtis, M. S. Weinberg, and M. J. a Wood, “Silencing of Parkinson’s disease-associated genes with artificial mirtron mimics of miR-1224.,” *Nucleic acids research*, vol. 40, no. 19, pp. 9863–75, Oct. 2012.
- [54] Anna Git, Heidi Dvinge, Mali Salmon-Divon, Michelle Osborne, Claudia Kutter, James Hadfield, Paul Bertone, Carlos Caldas, „Systematic comparison of microarray profiling, real-time PCR, and next-generation sequencing technologies for measuring differential microRNA expression.,” *RNA*, 16, 5:991–1006., 2010.
- [55] „R: A Language and Environment for Statistical Computing”, <http://www.R-project.org>
- [56] Rosenbloom KR, Sloan CA, Malladi VS, Dreszer TR, Learned K, Kirkup VM, Wong MC, Maddren M, Fang R, Heitner SG, Lee BT, Barber GP, Harte RA, Diekhans M, Long JC, Wilder SP, Zweig AS, Karolchik D, Kuhn RM, Haussler D, Kent WJ., „ENCODE data in the UCSC Genome Browser: year 5 update.,” *Nucleic Acids Res.*, 41(Database issue):D56-63, 2013 Jan.

Köszönetnyilvánítások

Köszönöm témavezetőmnek, Orbán Tamásnak a szakmai segítséget és a lehetőséget, hogy a laborjában dolgozhatom. Köszönöm a segítséget Schamberger Anitának és a labor többi dolgozójának. Továbbá köszönöm Erdei Zsuzsának és Apáti Ágotának a HuES9 mintákat, melyeken a vizsgálataimat végeztem.