

VI. ÉVFOLYAM 3-4. SZÁM 2014. OKTÓBER

IMMUNOLÓGIAI SZEMLE

NEGYEDÉVENTE MEGJELENŐ
ORVOSTUDOMÁNYI FOLYÓIRAT

IMMUNOLOGY QUARTERLY

ALAPÍTVÁ 2009-BEN

MEDICINA KÖNYVKIADÓ

WWW.MEDICINA-KIADO.HU

Köszöntő

A Magyar Immunológiai Társaság 43. Vándorgyűlésének összefoglalói

Nem genomikus glukokortikoid-hatások

A Marfan-szindróma klinikai jellemzői

Epesavak, immunrendszer

Spondyloarthritis: célértékre törekvő kezelés

Methotrexat rheumatoid arthritisben

Methotrexatezelés finomhangolása



V. Károly császár utolsó hónapjai

In memoriam Gomez Roberto

Tudta-e Ön ...?

Kongresszusi beszámoló: EULAR

Kongresszusi naptár



ISSN 2061-0203

mice contained few PNAd⁺ HEVs. Treatment of neonatal Nkx2-3 mice with LTβR-Ig fusion protein led to the complete absence of PNAd and the disappearance of residual MADCAM-1. Cell extravasation through PNAd⁺ vascular segments in mutant PPs was significantly slower compared to wild type samples. Adult PPs in Nkx2-3 mutant mice contained higher number of CD3⁺ T cells and regulatory T cells and significantly less B cells.

Conclusions: Our results indicate that appearance of PNAd⁺ HEVs requires activation of LTβR. However, this signal can be delivered by non-T/non-B cells as Nkx2-3^{-/-}xRag1^{-/-} mice lacking mature B and T cells expressed PNAd⁺ vessels. The switch in addressin profile influences the lymphocyte composition and also the rate of cell homing to mutant PPs.

Supported by OTKA K108429

GLOBAL ANALYSIS OF RXR-REGULATED GENES IN MACROPHAGES REVEALS A GENE NETWORK POTENTIALLY INVOLVED IN PROMOTING ANGIOGENESIS AND METASTASIS

KISS MÁTÉ¹, DÁNIEL BENCE¹, CZIMMERER ZSOLT¹, PAP ATTILA¹, PÓLISKA SZILÁRD², NAGY LÁSZLÓ^{1,3}

¹Department of Biochemistry and Molecular Biology, Research Center for Molecular Medicine, University of Debrecen, Debrecen, Hungary

²Center for Clinical Genomics and Personalized Medicine, University of Debrecen, Debrecen, Hungary

³MTA-DE „Lendület” Immunogenomics Research Group, University of Debrecen, Debrecen, Hungary

Introduction: Retinoid X receptor (RXR), a member of the family of nuclear receptors, is activated by vitamin A metabolite 9-cis-retinoic acid and fatty acids and acts as the obligate heterodimeric partner of several nuclear receptors, like PPAR, LXR, or RAR. Thus, RXR can mediate diverse signaling pathways simultaneously as a component of various heterodimers. Nevertheless, the global pattern of RXR-regulated gene expression changes in macrophages has not yet been analyzed. Our aim was to examine the genome-wide effect of the absence of RXR (loss of function) as well as specific activation of the receptor (gain of function) in macrophages, and identify novel RXR-dependent gene networks potentially involved in the regulation of macrophage metabolism and immune function.

Methods: Bone-marrow-derived macrophages (BMDMs) from Lys-M-Cre RXRα^{+/+}β^{-/-} (control) and Lys-M-Cre RXRα^{flox/flox}β^{-/-} (macrophage RXR KO) animals were treated with pan-RXR agonist LG268. Changes in mRNA expression induced by specific RXR activation or the absence of RXR were detected by microarray analysis. The angiogenic

potential of RXR-activated BMDMs was assessed by human umbilical vein endothelial cell (HUVEC)-based tube formation assay. Primary and metastatic tumor development in animals transplanted with macrophage RXR KO bone marrow was studied in B16-F10 mouse melanoma xenograft model.

Results: The absence of RXR in BMDMs resulted in the altered expression of 141 genes (fold change ≥ 1.3; p ≤ 0.05), which comprised a gene network linked to macrophage adhesion and tissue accumulation. In addition, several of these genes (e.g. S100a8, S100a9, Camp, Sell) have been previously shown to regulate the pro-metastatic action of macrophages. We found 103 genes, which were regulated by specific RXR activation in BMDMs. Among the upregulated genes we found Vegfa, one of the crucial mediators of tumor angiogenesis. Using Ingenuity Pathway Analysis, we identified several additional RXR-induced genes (e.g. Hbegf, Nus1, Serpine1) also implicated in angiogenesis. Accordingly, RXR-activated macrophages showed enhanced angiogenic potential in HUVEC tube formation assay *in vitro*. However, primary B16-F10 melanoma tumor growth was unimpaired in macrophage RXR KO animals, suggesting that RXR expressed by tumor-infiltrating macrophages is dispensable for tumor vascularization. In contrast, mice lacking macrophage RXR had a significantly higher number of B16-F10 lung metastases.

Conclusion: Our results suggest that *in vitro* RXR activation generates a macrophage phenotype possessing pro-angiogenic capacity. Based on our *in vivo* observations we hypothesize that this signaling pathway is presumably either inactive or silenced in the primary tumor microenvironment, but has a potential role altering pro-metastatic activity of macrophages.

INVESTIGATION OF ZAP-70 EXPRESSION IN THE PERIPHERAL BLOOD OF PATIENTS WITH CHRONIC LYMPHOCYTIC LEUKEMIA USING FLOW CYTOMETRY

KOHL ZOLTÁN¹, HAJNA LOSONCZY², TIMEA BERKI¹, FERENC BOLDIZSÁR¹

¹UP, CC, Department of Immunology and Biotechnology

²UP, CC, 1st Department of Internal Medicine

Chronic lymphocytic leukemia (CLL) is the most common form of adult leukemia in the Western world. Two main subtypes of the disease had been previously defined, an indolent form and a more progressive variant.

ZAP-70 is a tyrosine kinase that plays a major role during the early phases of TCR signaling. Interestingly, ZAP-70 is expressed by the leukemic B-cells in a subgroup of CLL patients and has been shown to correlate with a worse clinical prognosis. Although the flow cytometric analysis of ZAP-70 expression in CLL was described more than a decade ago, it

is still Pécs d

Our cible fl into di betwee see if t

We of CLL control anti-CD subgro perme: ALEXA intracel then at which fluoresc tive. Th definec ture.

Our 2014. I among results. positive The me experin in the i regardir vs. 40% median (7.16 y tients cl lent clin 30% of treatme the neg. still rem Besi results c progress with the