



ACTA MINERALOGICA-PETROGRAPHICA

ABSTRACT SERIES

Volume 7

Miskolc, 2012

JOINT MSCC+CEMC 2012

5th Mineral Sciences in the Carpathians Conference

3rd Central-European Mineralogical Conference



Co-organiser



Scientific
sponsors:



ACTA MINERALOGICA-PETROGRAPHICA

established in 1922

ABSTRACT SERIES

HU ISSN 1589-4835

HU ISSN 0324-6523

Editor-In-Chief

Elemér Pál-Molnár

University of Szeged, Szeged, Hungary

E-mail: palm@geo.u-szeged.hu

EDITORIAL BOARD

Péter Árkai, György Buda, István Dódoný, Tamás Fancsik, János Földessy,
Szabolcs Harangi, Magdolna Hetényi, Balázs Koroknai, Tivadar M. Tóth, Gábor Papp,
Mihály Pósfai, Péter Rózsa, Péter Sipos, Csaba Szabó, Sándor Szakáll,
Tibor Szederkényi, István Viczián, Tibor Zelenka

Editorial Office Manager

Anikó Batki

University of Szeged, Szeged, Hungary

E-mail: batki@geo.u-szeged.hu

Editorial Address

H-6701 Szeged, Hungary

P.O. Box 651

E-mail: asviroda@geo.u-szeged.hu

Abbreviated title:

Acta Mineral. Petrogr. Abstr. Ser., Szeged

The Acta Mineralogica-Petrographica is published by the Department of Mineralogy, Geochemistry and Petrology, University of Szeged, Szeged, Hungary

© Department of Mineralogy, Geochemistry and Petrology, University of Szeged

On the cover: The main entrance of the University of Miskolc.

On the logo of the conference: Euchroite from Lubietová, Slovakia (after Grailich & Lang, 1857).

JOINT
5th MINERAL SCIENCES IN THE CARPATHIANS CONFERENCE (MSCC)
and
3rd CENTRAL-EUROPEAN MINERALOGICAL CONFERENCE (CEMC)
Miskolc, Hungary, 19–21 April 2012



ABSTRACTS

Edited by

Béla Fehér

Abstracts revised by

Ferenc Máдай, Csaba Szabó, Sándor Szakáll, Pavel Uher, Tamás G. Weiszbürg and
Norbert Zajzon

Sponsored by

CBGA Commission on Mineralogy and Geochemistry
Czech Geological Society
Herman Ottó Museum

Mineralogical and Geochemical Branch of the Hungarian Geological Society
Mineralogical Society of Poland
Mineralogical Society of Romania
Slovak Geological Society
Ukrainian Mineralogical Society

Organizer

University of Miskolc

Co-organizer

Slovak Mineralogical Society

International Scientific Board

Igor Broska

Geological Institute, Slovak Academy of Sciences, Bratislava, Slovakia

Corina Ionescu

Babeş-Bolyai University, Cluj-Napoca, Romania

Friedrich Koller

University of Vienna, Vienna, Austria

Victor Kvasnytsya

Institute of Geochemistry, Mineralogy and Ore Formation, National Academy of Sciences, Kyiv, Ukraine

Milan Novák

Masaryk University, Brno, Czech Republic

Adam Pieczka

AGH – University of Science and Technology, Cracow, Poland

Mihály Pósfai

University of Pannonia, Veszprém, Hungary

Local Organizing Committee

Sándor Szakáll (chairman)

János Földessy, Ferenc Mádai, Viktor Mádai and Norbert Zajzon

University of Miskolc, Miskolc, Hungary

Béla Fehér

Herman Ottó Museum, Miskolc, Hungary

MODAL COMPOSITIONS AND TEXTURES OF ALKALI BASALT HOSTED UPPER MANTLE XENOLITHS FROM THE NORTHERN PART OF THE CARPATHIAN-PANNONIAN REGION (NÓGRÁD-GÖMÖR VOLCANIC FIELD, HUNGARY-SLOVAKIA)

LIPTAI, N.*; PATKÓ, L. & ARADI, L.

Lithosphere Fluid Research Lab, Department of Petrology and Geochemistry, Eötvös University, Pázmány Péter sétány 1/C, H-1117 Budapest, Hungary

* E-mail: n.liptai.elte@gmail.com

One possible way to understand upper mantle processes better is to study ultramafic xenoliths hosted in alkali basalt, kimberlite and lamprophyre. Upper mantle xenoliths from the five volcanic fields of Plio-Pleistocene alkali basalts in the Carpathian-Pannonian region (Styrian Basin, Little Hungarian Plain, Bakony-Balaton Highland, Nógrád-Gömör Volcanic Field, Perșani Mountains) have provided matter to petrologic and geochemical studies in the last few decades (e.g. EMBEY-ISZTIN *et al.*, 1989, 2001; DOWNES *et al.*, 1992; SZABÓ, 2004; DOBOSI *et al.*, 2010). In our work we concentrate on the Nógrád-Gömör Volcanic Field (NGVF), located at the northern part of the Carpathian-Pannonian region. We collected xenoliths from the central part of the area, including five quarries from Medves Plateau and Babi Hill that have barely been studied before.

From more than 200 collected xenoliths, over 80 samples were chosen for detailed petrographic examinations. The main aim was to determine texture types and modal compositions of the xenoliths regarding the rock-forming minerals, as well as describe fabrics of the mantle constituents (olivine, orthopyroxene, clinopyroxene and spinel). Based on the results, we present a detailed summary about the NGVF xenoliths, showing how they can be correlated with xenoliths originating from better researched marginal areas, introducing a

texture type (poikilitic) (Fig. 1), as well as a rock type (wehrlite) (Fig. 2) that have not been reported before in earlier studies on the central part of the NGVF. It will also be presented that wehrlites show features of different texture types in the form of clinopyroxene-spinel aggregations sitting in the original texture, which indicates a possible mantle metasomatism that took place before the ascension of the host basaltic melt. Presence of amphibole in some of the xenoliths also confirms that this process enriched the upper mantle in incompatible elements.

References

- DOBOSI, G., JENNER, G.A., EMBEY-ISZTIN, A. & DOWNES, H. (2010): Geological Society, London, Special Publications, 337: 177–194.
- DOWNES, H., EMBEY-ISZTIN, A. & THIRLWALL, M.F. (1992): Contributions to Mineralogy and Petrology, 109: 340–354.
- EMBEY-ISZTIN, A., SCHARBERT, H.G., DIETRICH, H. & POULTIDIS, H. (1989): Journal of Petrology, 30: 79–105.
- EMBEY-ISZTIN, A., DOBOSI, G., ALTHERR, R. & MEYER, H.P. (2001): Tectonophysics, 331: 285–306.
- SZABÓ, Cs., FALUS, Gy., ZAJACZ, Z., KOVÁCS, I. & BALI, E. (2004): Tectonophysics, 393: 119–137.




Fig. 1. Olivine (ol) inclusion in orthopyroxene (opx) as an indication of poikilitic textured lherzolite; transmitted light, N+




Fig. 2. Aggregation of clinopyroxene (cpx) and vermicular spinel (sp) in a wehrlite; transmitted light, 1N