



University of Novi Sad
**INSTITUTE
OF FOOD
TECHNOLOGY**



XV International
Feed Technology
Symposium



COST -
„Feed for Health“
joint Workshop



PROCEEDINGS

Novi Sad, 03-05 October, 2012.

ISBN 978-86-7994-032-2

**XV INTERNATIONAL FEED TECHNOLOGY SYMPOSIUM
“FEED-TO-FOOD” / COST FEED FOR HEALTH JOINT
WORKSHOP, NOVI SAD, 2012.**

Publisher

University of Novi Sad
Institute of Food Technology
Bulevar cara Lazara 1.
21000 Novi Sad

Main editor

Dr Jovanka Lević

Editors

Mr Slavica Sredanović
Dr Olivera Đuragić

Abstract/Paper Review

All abstracts and papers are reviewed by International Scientific Committee and competent researchers

Technical editor

Bojana Kokić

Cover

Boris Bartula, BIS, Novi Sad, Serbia

Printed by

“Futura” – Novi Sad, Serbia

Number of copies

200 copies

CONTENT

FEED TO FOOD	1
<i>Jovanka Lević, Slavica Sredanović</i>	
COST ACTION FEED FOR HEALTH 2008-2012 MAIN ACHIEVEMENTS	2
<i>Luciano Pinotti</i>	
VALIDATION OF IMMUNOASSAYS FOR THE DETECTION OF PROCESSED RUMINANT PROTEINS IN NON-RUMINANT PROTEINS	3
<i>Leo W.D. van Raamsdonk, Rob J.C.F. Margry, Robert G.C. Van Kaathoven, Monique G.E.G. Bremer</i>	
REINTRODUCTION OF PROCESSED ANIMAL PROTEINS IN FEED: FILLED GAPS AND GAPS TO BE FILLED	4
<i>Pascal Veys, Marie-Caroline Lecrenier, Vincent Baeten</i>	
IMPLEMENTATION OF THE REAL-TIME PCR AS OFFICIAL METHOD OF DETECTION OF PROCESSED ANIMAL PROTEINS IN THE EUROPEAN UNION REFERENCE LABORATORY NETWORK	5
<i>Olivier Fumière, Aline Marien, Pascal Veys, Vincent Baeten, Gilbert Berben</i>	
PRODUCTION OF FERMENTED SAUSAGES WITH OLIVE OIL, PROBIOTICS AND DIETARY FIBERS	7
<i>Taxiarhoulia Magra, Nikolaos Soutos, Elias Papapanagiotou, Ioannis Amvrosiadis</i>	
DIETARY SPECIES-SPECIFIC PROBIOTIC CAN CONTRAST MULTIRESTANT E. COLI ISOLATES IN THE GUT OF VEAL CALVES	8
<i>Alessandro Agazzi, Simone Stella, Erica Tirloni, Serena Marocco, Barbara Ripamonti, Carla Bersani, Giovanni Savoini</i>	
NEWBORN CALF FED SPECIES-SPECIFIC PROBIOTIC: EFFECTS ON GROWTH PERFORMANCE, HEALTH STATUS, MICROBIOLOGICAL AND HEMATOLOGICAL PARAMETERS AND CELL MEDIATE IMMUNE RESPONSE	9
<i>Alessandro Agazzi, Serena Marocco, Erica Tirloni, Simone Stella, Barbara Ripamonti, Carla Bersani, Vittorio dell'Orto, Giovanni Savoini</i>	
THERMAL ANALYSIS OF REHYDRATED PORK MEAT	10
<i>Sanja Ostojić, Darko Micić, Snežana Zlatanović, Lato Pezo, Danijela Šuput, Branislav Simonović, Ljubinko Lević</i>	
ROBUSTNESS OF NIRS CALIBRATIONS FOR TRANSFERENCE TO PORTABLE INSTRUMENTS	11
<i>Sagrario Modroño, Ana Soldado, Adela Martínez-Fernández, Begoña de la Roza-Delgado</i>	
INFLUENCE OF Se SUPPLEMENTATION OVER TRACE ELEMENTS IN COW MILK	12
<i>Amelia González-Arrojo, Ana Soldado, Fernando Vicente, M^a Luisa Fernández Sánchez, Alfredo Sanz-Medel, Begoña de la Roza-Delgado</i>	

AROMABIOTIC POULTRY FOR AN IMPROVED FEED EFFICIENCY AND BREAST MEAT YIELD IN BROILERS <i>Inne Gantois, Jan Vervloesem, Rejhana Semic, Andras Muzsek, László Pál, Károly Dublec</i>	13
SALBIOTIC, THE ULTIMATE HURDLE AGAINST SALMONELLA <i>Inne Gantois, Jan Vervloesem, Rejhana Semic</i>	14
THE USE OF LACUNAE AREA/FRAGMENT AREA RATIO AS A MARKER IN DISTINGUISH BETWEEN LAND ANIMAL VS. SEA MAMMALS IDENTIFICATION <i>L. Pinotti, V. Baeten, P. Veys, T. Fearn, M. Ottoboni, F. Cheli, V. Dell'Orto</i>	15
MANGANESE IN CHICKEN AND SWINE FEED: A REGIONAL MONITORING IN ITALY <i>Daniela Marchis, Paola Brizio, Matteo Pellegrino, Stefania Gavinelli, Renata Tarasco, Giuseppina Amato, Stefania Squadrone, Maria Cesarina Abete</i>	16
WHEAT MILLING AND MYCOTOXIN FRACTIONATION IN BY-PRODUCTS: A SYSTEMATIC REVIEW <i>Federica Cheli, Luciano Pinotti, Luciana Rossi, Vittorio Dell'Orto</i>	17
ANTIOXIDANT CAPACITY OF SELECTED GRAINS AND FEEDS <i>Vida Rezar, Alenka Levart, Darja Kocjan Ačko, Janez Salobir</i>	18
EFFECT OF LONG-TERM FEEDING OF CRUDE GLYCERINE ON PERFORMANCE, CARCASS TRAITS, MEAT QUALITY AND METABOLIC PARAMETERS OF FINISHING BULLS <i>Milan Marounek, Luděk Bartoň, Petr Homolka</i>	19
STUDY OF TRADITIONAL PONTIAKOS CAVOURMAS MADE FROM WILD BOAR MEAT <i>Timotheos Zougkouridis, Nikolaos Soultos, Taxiarchoula Magra, Ekaterini Papavergou, Ioannis Amvrosiadis</i>	20
PHYTANIC ACID IN ORGANIC MILK – A MATTER OF FEEDING <i>Mette Krogh Larsen, Brita Ngum Che, Troels Kristensen, Trine Kastrup Dalsgaard, Lars I. Hellgren, Jette Feveile Young</i>	21
CONSUMER PERCEPTION OF ANIMAL FEED IN RELATION TO FOOD SAFETY <i>Nataša Vukelić, Jasmina Živković, Djordje Okanović, Nikola Puvača</i>	22
TRANSFER OF VOLATILES FROM OREGANO OR CARAWAY ESSENTIAL OILS INTO COW'S MILK <i>Johan Lejonklev, Mette Marie Løkke, Mette Krogh Larsen, Grith Mortensen, Mikael Agerlin Petersen, Martin Riis Weisbjerg</i>	23
THE IMPACT OF MYCOTOXINS ON THE INTESTINE <i>Isabelle Oswald</i>	24
EFFECTS OF PROCESSING METHOD ON INTAKE, MILK YIELD AND MILK FATTY ACID PATTERN WHEN FEEDING COWS LINSEED-ENRICHED CONCENTRATE FEEDS <i>Anjo Elgersma, Jeroen Brok, Seerp Tamminga</i>	25

EFFECT OF FORAGE DRY MATTER CONTENT AND BALE SIZE ON FERMENTATION CHARACTERISTICS AND AEROBIC STABILITY OF ALFALFA ENSILED IN ROUND BALES	
<i>Sema Yaman, Engin Ünay, Vedat Karakaş, Murat Yıldırım, Şinasi Aşkar, Ali Bilgen, Mustafa Avcı, Tugay Ayaşan</i>	26
QUALITY CONTROL OF FEED IN REPUBLIC OF SRPSKA	
<i>Tanja Malčić, Tatjana Docić-Kojadinović, Duška Jokić</i>	27
GARLIC IMPAIRS ACTINOBACILLUS PLEUROPNEUMONIAE IN VITRO AND ALLEVIATES PLEUROPNEUMONIA IN A PIG MODEL	
<i>Petra Becker, Piet van Wikselaar, Monique Mul, Arjan Pol, Bas Engel, Jan Wijdenes, Carola van der Peet-Schwering, Henk Wisselink, Norbert Stockhofe-Zurwieden</i>	28
FOOD AND FEED-RELATED PATHOGEN AND TOXIN BINDERS FOR AN IMPROVED GUT HEALTH	
<i>Petra Becker, Piet van Wikselaar</i>	29
POTENTIAL OF MINOR OILSEED MEALS AS FEEDSTUFF FOR ORGANIC & LOW INPUT DAIRY SYSTEMS	
<i>Catalin Dragomir, Marketta Rinne, David Yanez Ruiz</i>	30
ASSESSMENT OF SOME HYGIENIC PARAMETERS OF ANIMAL FEEDS IN SERBIA	
<i>Ana Varga, Jovanka Lević, Ivana Čabarkapa, Bojana Kokić, Dragana Plavšić, Ljubiša Šarić</i>	31
TECHNOLOGICAL RESEARCH ON THE PRODUCTION OF FUNCTIONAL COOKED SAUSAGE FOR CHILDREN	
<i>Toni Doncheva, Djordje Okanović</i>	33
GROWTH PARAMETERS OF CARP FED MIXTURES CONTAINING DIFFERENT LEVELS AND ORIGIN OF PROTEINS	
<i>Marko Stanković, Vesna Poleksić, Nada Lakić, Zorka Dulić, Ivana Živić, Zoran Marković</i>	34
RED DEER (CERVUS ELAPHUS) – NEW PERSPECTIVE ON THE FARMS IN SLOVAKIA	
EFFECT OF SILAGE QUALITY ON BARK BROWSING AND NUTRIENTS DIGESTIBILITY	
<i>Matúš Rajs ký, Miroslav Vodnansky, Mária Chrenková, Rudolf Kropil, Dušan Rajs ký</i>	40
CHARACTERISTICS OF BEVERAGE OBTAINED FROM MILK PERMEATE	
<i>Dragana Ilić-Udovičić, Spasenija Milanović, Mirela Ilić</i>	45
EFFECTIVENESS OF FAGOPYRI HERBA FEED SUPPLEMENTATION IN NORMAL AND HIGH-FAT FED RATS	
<i>Aleksandra Mišan, Maja Đurendić-Brenesel, Ivan Milovanović, Miljan Milić, Vladimir Pilija, Đuro Vukmirović, Radmilo Čolović, Anamarija Mandić, Jovanka Lević</i>	52
LIMITATION OF FLAXSEED USAGE IN ANIMAL NUTRITION	
<i>Nikola Puvača, Vidica Stanačev, Dragan Milić, Bojana Kokić, Ivana Čabarkapa, Vladislav Stanačev</i>	58

USE OF PHYTOGENIC PRODUCTS FOR PIG AND BROILER DISEASES	
<i>Ljiljana Kostadinović, Jovanka Lević</i>	64
OPTIMIZATION OF PORK MEAT REHYDRATION PROCESS USING SEQUENTIAL QUADRATIC PROGRAMMING METHOD	
<i>Lato Pezo, Danijela Šuput, Biljana Dojčinović, Snežana Zlatanović, Olgica Kovačević, Ljubinko Lević</i>	75
FOOD INDUSTRY PERSPECTIVE ON INNOVATION IN BOSNIA AND HERZEGOVINA	
<i>Radoslav Grujić, Mitar Perušić, Jasna Mastilović</i>	81
VALIDATION METHODS FOR THE DETERMINATION OF β AGONISTS RESIDUES IN FEED	
<i>Risto Uzunov, Zehra Hajrulai-Musliu, Vlatko Ilieski, Biljana Stojanovska Dimzoska, Elizabeta Dimitrieska Stojkovic, Pavle Sekulovski, Velimir Stojkovski</i>	88
SIGNIFICANCE OF PROTEOLYTIC PROCESSES IN SILAGE FOR MODERN NUTRITION OF RUMINANTS	
<i>Bora Dinić, Nenad Đorđević, Zoran Lugić</i>	94
PREDICTION OF METABOLISABLE ENERGY OF FEED FOR POULTRY BY USE OF NON-LINEAR MODELS	
<i>Radmilo Čolović, Dragan Palić, Kedibone Y. Modika, Penny Barnes</i>	103
OCCURRENCE OF POTENTIAL MYCOTOXIN PRODUCING FUNGI ON MAIZE KERNEL IN HUNGARY	
<i>Beáta Tóth, Orsolya Török, Xénia Pálfi, Éva Toldi Tóth, Ákos Mesterházy, János Varga</i>	109
EXTRUSION-COOKING IN FEED PREMIXES STABILIZATION	
<i>Leszek Mościcki</i>	115
FATTY ACID COMPOSITION OF CARP MEAT: A REVIEW	
<i>Dušica Ivanov, Olivera Đuragić, Radmilo Čolović, Dragan Palić, Rade Jovanović, Jovanka Lević, Dragana Ljubojević</i>	119
AUTOCHTHONOUS BREED OF CHICKEN IN SERBIA: RESEARCH OR DEVELOPMENT	
<i>Zlatica Pavlovski, Zdenka Škrbić, Miloš Lukić</i>	127
SPECIFICS OF GOAT NUTRITION IN THE PRODUCTION CYCLE	
<i>Nurgin Memiši, Miroslav Žujović, Snežana Ivanović</i>	133
NUTRITIONAL VALUE OF RYE (<i>SECALE CEREALE</i> L.), TRITICALE (<i>TRITICO SECALE</i>) AND THEIR USE IN ANIMAL NUTRITION	
<i>Zuzana Formelová, Mária Chrenková, Ľubica Chrastinová, Mária Poláčiková</i>	141
TESTING OF CERTAIN CHEMICAL QUALITY CHARACTERISTICS OF VARIOUS SERBIAN MEAT PRODUCTS	
<i>Marija Vukašinović, Vladimir Kurćubić, Vesna Kaljević, Pavle Mašković, Milun Petrović, Slavica Vesković-Moračanin</i>	148
COMMON FOOD BORNE BACTERIAL DISEASES OF POULTRY	
<i>Ljiljana Suvajdzic, Jovanka Levic, Maja Velhner, Sava Pavkov, Aleksandar Potkonjak, Dubravka Milanov, Ivana Cabarkapa</i>	154

INFLUENCE OF <i>Kitaibelia vitifolia</i> EXTRACT ON COLOUR AND TEXTURE OF SREMSKA SAUSAGE	
<i>Djordje Okanović, Vladimir Kurćubić, Pavle Mašković, Marija Jokanović, Danijela Vranić, Slobodan Lilić, Natalija Džinić</i>	166
WHEAT ADMIXTURES AND THE POSSIBILITY OF THEIR VALORISATION	
<i>Slavko Filipović, Šandor Kormanjoš, Vera Radović, Rade Jovanović, Nikola Đukić, Jelena Filipović, Nikola Hristov</i>	172
CHANGES IN EAR COLOR OF SPELT WHEAT AS A RESULT OF DIFFERENT INTENSITY OF <i>ALTERNARIA SPP.</i> INFECTION	
<i>Jovana Vučković, Jovana Brkljača, Tea Brlek, Jelena Ćulafić, Aleksandra Bočarov-Stančić, Milica Aćimović, Marija Bodroža-Solarov</i>	179
INFLUENCE OF STORAGE CONDITIONS ON QUALITY OF RAPESEED OIL CAKE FOR FEED	
<i>Vanja Jurišić, Tajana Krička, Ana Matin, Neven Voća</i>	185
MONITORING OF RESIDUES OF PACKAGING MATERIALS IN BAKERY PRODUCTS USED AS FEED INGREDIENT	
<i>Leo W.D. van Raamsdonk, Victor G.Z. Pinckaers, Jef J.M. Vliege, Harry J. van Egmond</i>	191
ESTIMATION OF CHEMICAL AND FATTY ACID COMPOSITION OF SOME MARINE FISH SPECIES FROM RETAIL STORES	
<i>Dragana Ljubojević, Miroslav Ćirković, Nikolina Novakov, Vesna Đorđević, Dejana Trbović, Vesna Matekalo-Sverak, Danijela Vranić</i>	195
THE INFLUENCE OF DIFFERENT OSMOTIC SOLUTIONS ON NUTRITIVE PROFILE DURING OSMOTIC DEHYDRATION OF PORK	
<i>Milica Nićetin, Vladimir Filipović, Biljana Ćurčić, Lato Pezo, Violeta Knežević, Jasmina Gubić, Tatjana Kuljanin</i>	203
SALMONELLA FROM ANIMAL FEED: BIOFILM FORMING ABILITIES AND ANTIMICROBIAL SUSCEPTIBILITY	
<i>Dubravka Milanov, Dragica Stojanović, Bojana Prunić, Ivana Čabarkapa, Ljiljana Suvajdžić</i>	209
TESTING CONCENTRATE FEED MIXTURES WITH INCREASED SELENIUM CONCENTRATION IN ORDER TO PRODUCE EGGS AS FUNCTIONAL FOOD	
<i>Nataša Tolimir, Rade Jovanović, Marijana Maslovarić, Marina Vukić Vranješ, Milanka Drinić</i>	217
SHARE OF <i>PENICILLIUM</i> SPECIES IN MYCOPOPULATIONS ISOLATED FROM FEEDS	
<i>Marija Škrinjar, Mira Ač, Slavica Vesković-Moračanin, Vladislava Šošo, Nevena Blagojev</i>	223
GELATINIZATION OF STARCH AND METHODS FOR ITS DETERMINATION	
<i>Bojana Kokić, Dragan Palić, Olivera Đuragić, Nedeljka Spasevski, Tea Brlek, Dušica Ivanov, Radmilo Čolović</i>	229

IMPROVING THE NUTRITIVE VALUE OF CHICKEN MEAT BY THE ADDITION OF KOMPO OMEGA SUN AS FUNCTIONAL COMPONENT ENRICHED WITH LINSEED OIL <i>Đuro Vukmirović, Dušica Ivanov, Radmilo Čolović, Jovanka Lević, Rade Jovanović, Olivera Đuragić, Nedeljka Spasevski</i>	238
QUALITY OF WHEAT BY-PRODUCTS AIMED FOR ANIMAL FEEDING <i>Šandor Kormanjoš, Slavko Filipović, Ivana Čabarkapa, Vera Radović, Jelena Filipović, Nikola Hristov</i>	246
THE EFFECT OF CONCENTRATION OF MOLASSES ON TECHNOLOGICAL AND MICROBIOLOGICAL PARAMETERS OF OSMODEHYDRATED MEAT <i>Vladimir Filipović, Biljana Čurčić, Milica Nićetin, Lato Pezo, Violeta Knežević, Dragana Plavšić</i>	253
OPTIMIZATION OF PORK OSMOTIC DEHYDRATION PROCESS USING FUZZY SYNTHETIC EVALUATION <i>Biljana Čurčić, Vladimir Filipović, Milica Nićetin, Lato Pezo, Gordana Koprivica, Tatjana Kuljanin, Danijela Šuput</i>	260
OSMOTIC DEHYDRATION IMPACT ON MICROBIOLOGICAL PROFILE OF PACKED PORK MEAT <i>Danijela Šuput, Vera Lazić, Lato Pezo, Milica Nićetin, Vladimir Filipović, Biljana Čurčić, Nevena Krkić</i>	266
ASSESSMENT OF MINCED AND GRILL MEAT MICROBIOLOGICAL SAFETY IN YEAR 2012 <i>Ana Varga, Dragana Plavšić, Bojana Kokić, Tatjana Tasić, Ljubiša Šarić, Jasmina Gubić, Bojana Šarić</i>	273
COMPARISON OF THE NUTRITIONAL PROFILE OF GM MAIZE MON 89034 × NK603 AND CONVENTIONAL MAIZE <i>Mária Polačiková, Mária Chrenková, Zuzana Formelová, Ľubica Chrastinová, Silvia Pomikalová</i>	278
INFLUENCE OF SLAUGHTER AGE ON MEAT QUALITY OF GOAT <i>Snežana Ivanović, Zoran Stojanović, Boris Pisinov, Ksenija Nešić, Ivan Pihler, Nevena Maksimović, Nikola Stanišić</i>	284
HYDROLYTIC AND OXIDATIVE CHANGES OF UNPACKED AND PACKED BEEF <i>Danijela Vranic, Lazar Turubatovic, Slobodan Lilic, Zoran Petrovic, Milan Milijasevic, Radivoj Petronijevic, Dejana Trbovic</i>	290
ASSESSMENT OF ANTIMICROBIAL ACTIVITY OF EIGHT ESSENTIAL OILS AGAINST SALMONELLA ENTERITIDIS <i>Ivana Čabarkapa, Jovanka Lević, Slavica Sredanović, Bojana Kokić, Dubravka Milanov, Ljiljana Suvajdžić, Sava Pavkov</i>	296
IMPORTANCE OF CLINICAL AND PATHOLOGICAL DIAGNOSTICS OF MYCOTOXICOSIS IN FATTENING TURKEYS CAUSED BY T-2 TRICHOHECENE <i>Miloš Kapetanov, Dubravka Potkonjak, Igor Stojanov, Milica Živkov-Baloš, Sandra Jakšić</i>	304

TOTAL PHOSPHORUS, PHYTATE AND PHYTASE ACTIVITY OF SOME CEREALS GROWN IN ALBANIA AND USED IN NON RUMINANT FEED RATIONS	310
<i>Etleva Delia, Myqerem Tafaj, Klaus Männer</i>	
ECONOMIC FACTORS WITH INFLUENCE ON FEED MANUFACTURING	317
<i>Jasna Stevanovic, Aleksandar Drljacic, Biljana Miljkovic</i>	
APPLICATION OF NIR TECHNOLOGY FOR THE QUALITY CONTROL OF SUNFLOWER AND SOYBEAN MEAL AVAILABLE ON THE SERBIAN MARKET	323
<i>Marijana Maslovarić, Rade Jovanović, Snežana Janković, Nataša Tolimir, Marina Vukić – Vranješ</i>	
MOLASSES AS A FEED SUPPLEMENT	329
<i>Rada Jevtić-Mučibabić, Jasna Grbić, Bojana Filipčev, Vojislav Banjac, Tatjana Kuljanin, Jasmina Živković</i>	
THE INFLUENCE OF PHYTOGENIC ADDITIVES AND ORGANIC ACIDS SUPPLEMENTATION TO DIET ON BROILER CARCASS COMPOSITION AND MEAT YIELD	334
<i>Tatjana Tasić, Predrag Ikonić, Đorđe Okanović, Natalija Džinić, Jasmina Gubić, Dragan Milić, Marina Vukić - Vranješ</i>	
EFFECT OF HEMPSEED (CANNABIS SATIVA sp.) SUPPLEMENTATION TO DIET ON PERFORMANCE, CARCASS AND INTESTINAL ORGAN TRAITS IN JAPANESE QUAIL (Coturnix coturnix japonica)	340
<i>Yusuf Konca, Selma Büyükkiliç Beyzi</i>	
DETERMINATION OF THE EFFECTS AND TYPE OF LIQUID ADDITIVES ON MIXTURE HOMOGENEITY IN VARIOUS MIXERS	346
<i>Olivera Đuragić, Radmilo Čolović, Bojana Kokić, Jovanka Lević, Đuro Vukmirović, Slavica Sredanović, Dušica Ivanov</i>	
QUALITY MANAGEMENT	352
<i>Ružica Milovanović</i>	
QUALITY PARAMETERS OF MAIZE BIOMASS AS A FEED RAW MATERIAL	359
<i>Dušanka Terzić, Milica Radosavljević, Marija Milašinović-Šeremešić, Goran Todorović, Zorica Pajić, Milomir Filipović</i>	
CHANGES IN THE TECHNOLOGICAL QUALITY AND SAFETY OF WINTER WHEAT INFECTED WITH MYCOBIOTA	366
<i>Marija Bodroža-Solarov, Jovana Brkljača, Jovana Vučković, Tea Brlek, Ferenc Balaž, Maja Manojlović</i>	
COWS AND WETHERS NITROGEN EXCRETION DEPEND ON NUTRITION	371
<i>Mariana Petkova</i>	
MINERAL ADSORBENTS AND ENZYMES IN POULTRY PRODUCTION	377
<i>Vera Radović, Slavko Filipović, Šandor Kormanjoš</i>	

PERFORMANCE CHARACTERISTICS OF WEANED RABBITS FED GRADED LEVELS OF DRY CASSAVA PEEL FORTIFIED WITH SOYBEAN RESIDUE BASAL DIET	386
<i>Olayeni Victor Olutayo</i>	
INFLUENCE OF DIFFERENT PROTEIN SUPPLEMENTS ON SHEEP MILK QUANTITY AND QUALITY	392
<i>Marin Yossifov</i>	
DISCONTINUITY IN GRAIN DRYER OPERATION AS A CAUSE OF HIGHER ENERGY CONSUMPTION	404
<i>Mirko Babić, Ljiljana Babić, Ivan Pavkov, Milivoj Radojčin</i>	
BIODEGRADATION OF AFLATOXIN B1 BY FUNGAL SPECIES	410
<i>Aleksandra Bočarov-Stančić, Marija Bodroža-Solarov, Ferenc Bagi, Vera Stojšin, Jovana Vučković, Vladimir Pantić</i>	
DSM NUTRITIONAL PRODUCTS – YOUR PARTNER FOR EUBIOTIC SOLUTIONS	416
<i>Matthias Wiemann, Christophe Paulus, Zoran Grbic</i>	

OCCURRENCE OF POTENTIAL MYCOTOXIN PRODUCING FUNGI ON MAIZE KERNEL IN HUNGARY

Beáta Tóth¹, Orsolya Török¹, Xénia Pálfi¹, Éva Toldi Tóth¹,
Ákos Mesterházy¹, János Varga²

¹Cereal Research Nonprofit Ltd., 6726 Szeged, Alsó kikötő sor 9, Hungary

²University of Szeged, Faculty of Science and Informatics, Department of Microbiology, 6726 Szeged, Közép fasor 52, Hungary

ABSTRACT

Maize is one of the most important ingredients of animal feed formulations. *Fusarium* species are important pathogens of maize, causing various diseases, and contamination of maize kernel by mycotoxins including trichothecenes, zearalenone and fumonisins. *Aspergillus* and *Penicillium* species and their mycotoxins including aflatoxins, ochratoxins, fumonisins and patulin are also frequently encountered on cereal products. We investigated the occurrence of these species and their mycotoxins on maize in various maize growing areas in Hungary in 2010 and 2011 years after harvest. Surface-sterilized cereal seeds were placed on selective media, and the isolated fungal strains were identified using morphological methods. 81.94% and 14.33% of the samples were found to be contaminated with potentially toxigenic isolates in 2010 and 2011, respectively. Species identifications of selected isolates have been carried out using sequence-based methods. Regarding *Fusarium* species, in 2010, when the weather was rainy, *F. graminearum* and *F. subglutinans* dominated, while in 2011 with hot and dry summer *F. verticillioides* was the predominant species identified. *F. culmorum* could not be detected in any of the samples. Regarding *Aspergilli*, several *Aspergillus flavus* isolates were identified, which are potential aflatoxin producers. Besides, other mycotoxin producer species were also isolated, including black *Aspergilli* which potentially produce ochratoxins and fumonisins, and *A. clavatus*, which produces patulin. Other genera (*Alternaria*, *Nigrospora*, *Epicoccum*, *Cladosporium*) were found in smaller proportions. Besides, the protective maize endophyte *Acremonium zeae* was also identified in some of the samples. Further studies are in progress to examine the mycotoxin producing abilities of the isolates, mycotoxin content of the maize samples, and the applicability of *Acremonium zeae* isolates for lowering fungal burden and mycotoxin contamination of maize.

Keywords: maize, mycotoxins, aflatoxins, fumonisins, *Fusarium*, *Aspergillus*

INTRODUCTION

Maize is among the most important ingredients of feed formulations worldwide. *Fusarium* species are among the most important mycotoxin producing pathogens of maize. These species cause several diseases on maize including

ear rot and stalk rot (Figure 1), and contaminate maize kernel with various mycotoxins including trichothecenes, zearalenone and fumonisins. The most important maize pathogens are *F. graminearum*, *F. culmorum*, *F. verticillioides*, *F. proliferatum* and *F. subglutinans*. Besides, although not considered to be major causes of plant disease, *Aspergillus* and *Penicillium* species may also be responsible for several disorders in various plants including maize, and most importantly cause mycotoxin contamination [10]. The most notorious plant pathogens are black *Aspergilli* and *A. flavus* which may cause ear rot and are also important as postharvest pathogens. In contrast with specialized plant pathogens such as powdery mildews, rusts or most *Fusarium* species, these species are opportunistic pathogens without host specialization as proved in *A. flavus* [10]. The most important aspect of food and feed spoilage caused by these organisms is the formation of mycotoxins, which may have harmful effects on human and animal health. Several mycotoxins produced by *Aspergilli* or *Penicillia* have been identified in maize in previous studies, the economically most important of which are aflatoxins, ochratoxins and fumonisins [10]. In this study, we examined the mycobiota of maize samples collected in 2010 and 2011 from various maize growing regions of Hungary using morphological and sequence-based identification methods.



Figure 1. *Fusarium* ear rot of maize

MATERIAL AND METHODS

Sample collection

The samples were collected in 9 and 10 maize growing regions in 2010 and 2011, respectively. Maize grains were surface sterilized using ethanol, and plated onto dichloran rose bengal agar media [1]. Plates were incubated at 25°C in darkness and monitored periodically for characteristic mycelium growing from the kernels. Single colonies were purified and transferred to malt extract agar

(MEA) media. Isolates were subcultured as single conidia on MEA, Czapek-yeast extract agar (CYA) and potato dextrose agar (PDA) plates [8].

Identification of fungal isolates

Morphological identification of fungal isolates came from maize grains have been done according to standard textbooks and monographs [2, 6, 8]. For sequence based identification, the cultures used for the molecular studies were grown on malt peptone broth for 2 days, and DNA was extracted from the mycelia using the Masterpure™ yeast DNA purification kit (Epicentre Biotechnol.) according to the instructions of the manufacturer. Parts of the ITS, Tri101 and calmodulin genes were amplified and sequenced as described previously [3, 5]. Sequences were compared using nucleotide-nucleotide BLAST (blastn) with default settings (<http://blast.ncbi.nlm.nih.gov>) to the Genbank database, and to our own sequence database. Species identifications were determined from the lowest expect value of the BLAST output.

RESULTS AND DISCUSSION

The overall fungal contamination rate of the samples was 81.94% in 2010, while only 14.33% of the samples were found to be contaminated in 2011 (Table 1). The lower contamination rate observed in 2011 is possibly due to the weather conditions. While the summer was rainy in 2010, it was dry and hot in 2011. Possibly due to the humid weather conditions, *Penicillia* were frequently isolated from the samples collected in 2010, while this genus was virtually absent in the samples collected in 2011. On the other hand, *Aspergillus* infection rates were higher in 2011 than in 2010. The higher prevalence of *Aspergilli* might have been caused by the warmer weather conditions in 2011, as these species (e.g. *A. flavus* or black *Aspergilli*) prefer higher temperatures. However, *Fusaria* were present in the samples in both years in large proportions (Table 1).

The number of primary isolates of each sample was restricted upon the grounds of colony and microscopic features and only the diverging ones were maintained for further investigations. 340 and 90 isolates were recovered in 2010 and 2011, respectively.

Table 1. Occurrence of mycotoxigenic fungi on maize in Hungary in 2010-2011

Year	Average infection of grains (%)	% of isolated fungal strains (based on sequence-based identification)			
		<i>Aspergillus</i> sp.	<i>Penicillium</i> sp.	<i>Fusarium</i> sp.	other genera
2010	81.94	2.02	27.56	70.30	0.12
2011	14.33	8.27	0.00	60.25	31.48

Occurrence of *Fusaria* in maize kernel

In 2010, when the weather was rainy, *F. graminearum* and *F. subglutinans* dominated, while in 2011 with hot and dry summer *F. verticillioides* was the predominant species identified. *F. culmorum* could not be detected in any of the samples, in contrast with the results of the previous 1977 survey [4]. *F. graminearum* favors higher temperatures than *F. culmorum* and the observed shift might be an indication of climate change. However, *F. proliferatum* was detected in both years. Differences could not be observed in the species distribution of *Fusaria* in different locations (data not shown). Besides, *F. sporotrichioides* and *F. oxysporum* were also identified in some locations. Regarding the species distribution of *F. graminearum sensu lato* isolated from Hungarian maize, all isolates proved to belong to *F. graminearum sensu stricto* based on sequence analysis of the Tri101 gene of the isolates (data not shown). *Fusarium boothii* or *F. meridionale* could not be identified, in contrast with studies carried out in other countries including South Africa and Argentina [7]. Fumonisin and DON content of the maize samples was also analyzed using HPLC-MS. All samples were contaminated by these toxins under the EU limit (data not shown).

Occurrence of *Aspergilli* and *Penicillia* in maize kernel

Among the examined samples, several ones were found to be contaminated by members of section *Flavi* of the genus *Aspergillus* based on colony morphology and microscopic features. Although several *Aspergillus* species have been identified recently which are able to produce aflatoxins, *A. flavus*, *A. parasiticus* and *A. nomius* are the economically most important species regarding aflatoxin contamination of agricultural products [9]. These species can readily be distinguished using sequence analysis of part of their β -tubulin or calmodulin genes [9]. Species assignment of the isolates was carried out using partial sequence analysis of their calmodulin gene. All isolates assigned to section *Flavi* based on morphological features were found to belong to the *A. flavus* species based on sequence data.

Besides *A. flavus*, several other potential mycotoxin producers were identified in the samples. The patulin producer *A. clavatus* and black *Aspergilli* able to produce both ochratoxins and fumonisins were recovered from several samples. Among black *Aspergilli*, *A. niger* was identified most frequently (41 isolates), although *A. tubingensis* (10 isolates) was also isolated from some samples. Regarding *Penicillia*, several mycotoxin producers were identified (e.g. *P. crustosum*, *P. brevicompactum*, *P. chrysogenum* and *P. viridicatum*). Other potentially mycotoxigenic genera (*Alternaria*, *Nigrospora*, *Epicoccum*, *Cladosporium*) were found in smaller proportions (data not shown). Besides, the protective maize endophyte *Acremonium zeae* was also identified in some of the samples for the first time in Central Europe [11]. This species has been shown to inhibit infection and mycotoxin accumulation caused by *Fusaria* and *A. flavus* by producing the antibiotic compounds pyrrocidins [11].

Aflatoxin content of the samples was analyzed using HPLC-MS. None of the samples were found to be contaminated by any of the aflatoxins (B₁, B₂, G₁, G₂). Examination of ochratoxin content of the samples is in progress.

CONCLUSIONS

During a survey of mycotoxin producing molds in Hungarian maize samples in 2010 and 2011, 81.94% and 14.33% of the samples were found to be contaminated with potentially toxigenic isolates, respectively. Regarding *Fusarium* species, in 2010, when the weather was rainy, *F. graminearum* and *F. subglutinans* dominated, while in 2011 with hot and dry summer *F. verticillioides* was the predominant species identified. *F. culmorum* could not be detected in any of the samples. Among Aspergilli, several *Aspergillus flavus* isolates were identified, which are potential aflatoxin producers. Besides, other mycotoxin producing species, including black Aspergilli which potentially produce ochratoxins and fumonisins, and *Penicillium* species producing a range of mycotoxins have also been identified. The most contaminated samples came from 2010, possibly due to the rainy, humid weather conditions. Samples came from 2011 were found to be infected less severely, but the *Aspergillus* infection was higher than in the previous year. Further studies are in progress to examine the mycotoxin producing abilities of the isolates, mycotoxin content of the maize samples, and the applicability of *Acremonium zeae* isolates for lowering fungal infection and mycotoxin contamination of maize.

ACKNOWLEDGEMENTS

This work was supported by OTKA grant Nos. K84122 and K84077, and by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences (B. Tóth). The project is co-financed by the European Union through the Hungary-Serbia IPA Cross-border Co-operation Programme (ToxFreeFeed, HU-SRB/1002/122/062).

REFERENCES

1. King, A.D.Jr., Hocking, A.D., Pitt, J.I.: *Dichloran-rose bengal medium for enumeration and isolation of molds from foods*, Appl. Environ. Microbiol., 37 (1979), 959–964.
2. Leslie, J.F., Summerell, B.A.: *The Fusarium Laboratory Manual*. Blackwell Publishing Professional, Ames, USA, 2006, p. 388.
3. Malihipour, A.: *Genetic analysis of resistance to Fusarium head blight in wheat (Triticum spp.) using phenotypic characters and molecular markers*, Ph.D. Thesis, University of Manitoba, Winnipeg, Canada, 2010.
4. Mesterházy, Á., Vojtovics, M.: *A kukorica Fusarium spp. okozta fertőzőttségének vizsgálata 1972-1975-ben (Rate of Fusarium spp.*

- infection in maize in 1972-1975*), Növénytermelés 26 (1977), 367-378.[in Hungarian]
5. **Pildain, M.B., Frisvad, J.C., Vaamonde, G., Cabral, D., Varga, J., Samson, R.A.:** *Two novel aflatoxin-producing Aspergillus species from Argentinean peanuts*, Int. J. Syst. Evol. Microbiol., 58 (2008), 725-735.
 6. **Raper, K.B., Fennell D.I.:** *The genus Aspergillus*. Williams and Wilkins, Baltimore, USA, 1965, p. 686.
 7. **Sampietro, D.A., Díaz, C.G., Gonzalez, V., Vattuone, M.A., Ploper, L.D., Catalan, C.A., Ward, T.J.:** *Species diversity and toxigenic potential of Fusarium graminearum complex isolates from maize fields in northwest Argentina*, Int. J. Food Microbiol. 145 (2011), 359-264.
 8. **Samson, R.A., Hoekstra, E.S., Frisvad, J.C.:** *Introduction to Food- and Airborne Fungi*, 7th edition. CBS Fungal Biodiversity, Center, Utrecht, Netherlands, 2004, p. 389.
 9. **Varga, J., Frisvad, J.C., Samson, R.A.:** *A reappraisal of fungi producing aflatoxins*, World Mycotoxin J. 2 (2009), 263-277.
 10. **Varga, J., Tóth, B., Mesterházy, Á., Téren, J., Fazekas, B.:** *Mycotoxigenic fungi and mycotoxins in foods and feeds in Hungary*, in: An overview on toxigenic fungi and mycotoxins in Europe. Eds. **A. Logrieco, A. Visconti**. Kluwer Academic Publishers, Amsterdam, Netherlands, 2004, p. 123-139.
 11. **Wicklow, D.T., Roth, S., Deyrup, S.T., Gloer, J.B.:** *A protective endophyte of maize: Acremonium zeae antibiotics inhibitory to Aspergillus flavus and Fusarium verticillioides*, Mycol. Res. 109 (2005), 610–618.