

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Systematic study on free radical hydrothiolation of unsaturated monosaccharide derivatives with exo- and endocyclic double bonds[†]

László Lázár,^a Magdolna Csávás,^b Ádám Hadházi,^b Mihály Herczeg,^{a,c} Marietta Tóth,^c László Somsák,^c Terézia Barna,^d Pál Herczegh^b and Anikó Borbás*^{a,b}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

Exo- and endocyclic double bonds of glycals and terminal double bonds of enoses were reacted with various thiols by irradiation with UV light in the presence of a cleavable photoinitiator. The photoinduced radical-mediated hydrothiolation reactions showed highly varying overall conversions depending not only on the substitution pattern and electron-density of the double bond but also on the stability of the thiyl radical. Out of the applied thiols thiophenol, producing the most stabilized thiyl radical, exhibited the lowest reactivity toward each type of alkenes revealing the inverse proportionality between the stability of the thiyl radical and conversion of alkene in the thiol-ene coupling reaction. In most cases, the hydrothiolations took place with full regio- and stereoselectivities. Successful addition of 1,2:3,4-di-*O*-isopropylidene-6-thio- α -D-galactopyranose to a 2,3-unsaturated *N*-acetylneuraminic acid derivative, providing a (3→6)-*S*-linked pseudodisaccharide, demonstrated that the endocyclic double bond of Neu5Ac-2-ene, bearing an electron-withdrawing substituent, shows sufficient reactivity in the photoinduced thiol-ene coupling reaction.


Introduction

Over the last decade, the photoinduced thiol-ene coupling, free-radical addition of a thiol to a nonactivated carbon-carbon double bond, has been recognised as a robust ligation tool possessing many of the attributes of click chemistry.^{1,2} The great synthetic potential of the reaction has been amply demonstrated in areas of polymer chemistry and material sciences for network formation,³ dendrimer synthesis,⁴ and polymer functionalization.⁵

The photoinduced thiol-ene coupling was also applied as a metal-free click process for *S*-glycoconjugation producing glycodendrimers,⁶ calix[4]arene-based glycoclusters,⁷ *S*-linked protein glycoconjugates⁸ as well as thioglucoside-containing micellar structures appropriate for controlled drug delivery.⁹ In these approaches thioglycosylation occurred by addition of 1-thioglycoses across the terminal double bonds of alkenyl tags of dendritic or polymeric scaffolds. Carbohydrates equipped with alkenyl auxiliaries have been used also as the ene reactants in the thio-click strategy to afford alkyl-tethered glycosyl-cysteines¹⁰ or glycopeptides,¹¹ β -cyclodextrin-based saccharide clusters¹² and sucrose-containing polymers.¹³

The carbohydrate skeleton itself with an exo- or endocyclic double bond has scarcely been applied as the ene partner, with only three articles published until now.^{14–16} Dondoni and coworkers were the first who exploited the thiol-ene reaction for the synthesis of *S*-disaccharides by reacting sugar thiols with exo- and endoglycals.^{14,15} They found that hydrothiolation of the exocyclic double bond of hex-5-enopyranoside and pent-4-enofuranoside derivatives with 1-thioglycoses afforded *S*-disaccharides with high yields and diastereoselectivities of up to 99%.¹⁴ Addition of 1-thioglycoses across the endocyclic double bond of glycals showed regioselectivity but lack of

stereoselectivity, furnishing the stereoisomeric 1-deoxy *S*-disaccharides in about 1:1 ratios.¹⁵



Scheme 1. Sequential thiyl-addition and hydrogen abstraction steps during a thiol-ene coupling, and mechanism for the initiation with the cleavable photoinitiator 2,2-dimethoxy-2-phenylacetophenone (DPAP)

Incorporation of 2-acetoxy-glycals and 2,3-unsaturated glycosides within the thiol-ene coupling strategy was reported for

the first time by our group.¹⁶ We have shown that reactions between 2-acetoxy-D-glucal and a range of thiols including amino acid, peptide, and sugar thiols gave 1,2-*cis*- α -S-glucoconjugates and S-disaccharides with full regio- and stereoselectivities in good to excellent yields. Addition of 1-thioglycoses to a 2,3-unsaturated glycoside (Ferrier glycal) was found to proceed also with high selectivity offering an easy access to 3-deoxy-S-disaccharides.¹⁶

Regarding the great potential of hydrothiolation of the easily available glycals in the synthesis of thioglycosides, which are especially valuable glycoside mimetics, further exploration and exploitation of the reaction can be foreseen. Here we present a systematic study of the reactivity of the acetylated and benzylated derivatives of 2-hydroxy-D-glucal towards a range of thiols in the photoinduced thiol-ene coupling reaction, in comparison to the reactivity of saccharides bearing an exomethylene moiety. Free radical hydrothiolation of the endocyclic double bond of sialic acid glycal is also shown.















Results and discussion

To study the scope of the hydrothiolation reaction of unsaturated monosaccharides, 2-acetoxy-3,4,6-tri-O-acetyl-D-glucal **1**¹⁷ was reacted with thiols **2b-h** applying the optimized conditions established in our recent work for the synthesis of **3a**.¹⁶ Thus, the reactions were carried out in toluene at room temperature with a 2:1 thiol:ene ratio by irradiation at λ_{max} 365 nm for 3 x 15 min in the presence of 2,2-dimethoxy-2-phenylacetophenone (3 x 0.15 equiv) as the cleavable photoinitiator^{1a} (Table 1). In the case of ethanethiol and 2-methylpropane-2-thiol, higher thiol:ene ratio was applied because of the volatility of the reagents. When the reaction showed low conversion after 45 min, further portions of thiol and initiator were added and the irradiation was continued.¹⁸

Unexpectedly, thiophenol did not react with compound **1**. Our attempts to elicit a reaction between **1** and **2b** by applying high excess of thiol, increasing the amount of DPAP, and prolongation of the irradiation time were unsuccessful.

Aromatic thiols are better chain transfer reagents in free-radical additions than aliphatic thiols, since in the former case the energy required to break the S-H bond is lowered by the resonance stabilization of the thiyl radical formed.¹⁹ This fact seems to contradict with the observed reactivities of ethanethiol and thiophenol towards compound **1**. We assume, that the resolution of this contradiction resides in the reversibility of the reaction. The reactivity of the olefin partner in a thiol-ene free-radical chain reaction is dependent on the extent of substitution. Terminal alkenes are found to be the most reactive, while internal 1,2-disubstituted alkenes exhibit much lower overall reaction rates. The reduced rates and conversions observed for 1,2-disubstituted enes are a result of the reversibility of the addition of the thiyl radical across the internal double bond.^{1,19} In this case the reversible equilibrium in the addition step depicted in Scheme 2 is shifted backward by the high stability of the aromatic thiyl radical causing the complete lack of reaction between **1** and thiophenol.


Table 1. Free radical addition of thiols to 2-acetoxy glycal **1**

Thiol	thiol:ene ratio	Product	Yield ^a (%)
			
HSEt 2a	15:1		81
 2b	50:1	no reaction ^b	
 2c	4:1		35 ^c
 2d	4:1		10 ^c
 2e	15:1		25 ^c
 2f	2:1		67
 2g	2:1		31 ^c
 2h	2:1		64 ^d

^aYield of isolated compounds after 3 x 15 min irradiation under Ar, using 3 x 0.1equiv. DPAP; ^b6 x 15 min irradiation, 6 x 0.1 equiv. DPAP; ^c4 x 15 min irradiation, unreacted **1** was recovered with 55-60%; ^dThe reaction was carried out in abs. MeOH.

Hydrothiolation of 2-acetoxy glycal **1** with thiols **2c-2h** took place with full regio- and stereoselectivities affording the corresponding α -D-1-thioglycosides, however, with various conversions. The reactivities of **2c** and **2d** towards **1** was strongly influenced by the stability of the thiyl radicals formed by irradiation. The benzyl-thiyl radical generated from **2c** afforded the hydrothiolation product **3c** with 35% yield, while the 4-

methoxybenzyl-thiyl radical, being more stabilized due to the electron-donating methoxy group, furnished the product **3d** only with 10%.



Scheme 2. Reversible addition of the thiyl radical to the endocyclic double bond of **1**

Reaction of 2-methylpropane-2-thiol with glycal **1** gave the α -thioglucoside **3e** with 25% isolated yield. The low conversion can be explained by either the stability of the thiyl radical or a steric congestion between the 2-acetoxy moiety and the bulky *tert*-butyl group, both anchored to the α -side of the sugar ring in the intermediate C-2 radical.

The different conversion in the reaction of **1** with 2,3-di-O-acetyl-1-thioglycerol **2f**²⁰ compared to that with acetonide-protected 1-thioglycerol **2g**²¹ provided another example for the inverse proportionality between reactivity and stability of the thiyl radical in the reversible addition step. While the less stable radical formed from **2f** possessing electron withdrawing substituents afforded **3f** in 67% yield, the more stable thiyl radical generated from **2g** furnished product **3g** with a yield of only 31%.

Finally, the sodium sulfonatoethyl mercaptane **2h** (Mesna), used as a detoxifying adjuvant in cancer chemotherapy, was reacted with **1**. We were pleased to find that compound **2h** added readily across the endocyclic double bond of **1** in MeOH and gave **3h** in good yield. To the best of our knowledge, this is the first example of using a salt as the thiol partner in the photoinduced hydrothiolation reaction.

Next, we decided to study the photoinduced hydrothiolation reactions of the benzylated 2-hydroxyglycal **4**.²² It is known, that the overall reaction rate of the thiol-ene reaction is directly related to the electron density on the ene: for a given thiol, electron-rich enes react much more rapidly than electron-poor ones.^{1,19} Compound **4**, possessing a more electron-rich double bond than that of **1**, was expected to show higher conversions with the thiols which exhibited low reactivity towards compound **1**. Hydrothiolation of **4** with ethanethiol went to completion rapidly and with full selectivity to give the expected α -thioglucoside **5a** in 71% yield (Table 2). The slight decrease of the yield compared to that of **3a** was a result of the lability of the benzyl-type protecting groups under shortwave UV irradiation causing partial decomposition.²³⁻²⁵

To our delight, when glycal **4** was reacted with thiophenol, a fair conversion of **4** was observed after 45 min irradiation affording the α -glucoside **5b** in a yield of 21%. Applying higher excess of thiol or longer exposure to UV light did not give rise noticeable increase of the conversion. Despite the moderate yield, this reaction clearly demonstrates that the highly stable thiophenyl radical can be trapped by an internal double bond of sufficient electron-density and the equilibrium of the thiyl addition can be forced toward the irreversible hydrogen abstraction step to give the final addition product.

Reaction of 2-methylpropane-2-thiol with **4** gave α -thioglucoside **5e** with 40% isolated yield. The good conversion of

4 with **2e** revealed that rather the stability of the thiyl radical than steric hindrance had caused the low reactivity of **2e** towards the acetoxy-glycal **1**.

Table 2. Free radical addition of thiols to 2-benzoyloxy glycal **4**

Thiol	thiol:ene ratio	Product	Yield ^a (%)
HSEt 2a	15:1	5a	71
 2b	2:1	5b	21
 2e	15:1	5e	40
 2i	2:1	5i	46 ^b
 2j	2:1	5j	59


^aYield of isolated compounds after 3 x 15 min irradiation under Ar, using 3 x 0.1 equiv. DPAP; ^bThe reaction was carried out in abs. MeOH.

Previously, we have found that reaction between **1** and 1-thioglycerol **2i** showed very low conversion of the glycal either toluene-MeOH or MeOH was used as the solvent, furnishing the corresponding α -thioglucoside with only 15-17% yield.¹⁶ As we expected, thiol **2i** reacted more readily with the more electron-rich double bond of **4** providing product **5i** with 46% yield.

Finally, S-disaccharide **5j** was prepared by the reaction between **4** and the sterically hindered di-O-isopropylidenated sugar thiol **2j**.²⁶ Despite complete conversion of the glycal observed, the isolated yield of **5j** was only 59% because decomposition also occurred due to lability of the benzyl-

protecting groups under irradiation.²⁷

Thereafter, we studied the photoinduced hydrothiolation reactions of glycal **6** with ethanethiol and thiophenol under the optimized conditions. (Scheme 3). In this case, addition of thiophenol across the endocyclic double bond of **6** failed, revealing again the lack of reactivity of the resonance stabilized aromatic thiyl radical towards a relatively electron-poor internal double bond. Reaction between glycal **6** and ethanethiol showed almost complete conversion after 45 min affording a mixture of the axially and equatorially linked 2-ethylthio derivatives **7** and **8** and thioglycoside **9** in a ratio of ~5:3:1.^{28,29} Because of the poor selectivity, which was also reported by Dondoni and co-workers,^{15,30} and the difficulty of the separation of the diastereoisomers formed, hydrothiolation of glycal **6** was not examined further.



Scheme 3. Reagents and conditions: a) 15 equiv. EtSH, toluene, 3 x 15 min irradiation, 3 x 0.1 equiv. DPAP; b) 50 equiv. PhSH toluene, 6 x 15 min irradiation, 6 x 0.1 equiv. DPAP.

Reactivity of thiophenol towards terminal double bonds of monosaccharides was studied next, by applying compounds **10**,³¹ **12**³² and **14**³³ as the ene partners (Table 3). The reactions were carried out in toluene with a 2:1 thiol:ene ratio. Hydrothiolation of the galactose derived alkene **10** afforded the addition product **11** in 46% yield. Reaction of 3-exomethylene-glucufuranose derivative **12** with thiophenol showed slightly lower efficiency but full selectivity furnishing compound **13** as a single diastereoisomer in 32% yield. We were pleased to find that in the reaction of exoglycal **14** with thiophenol almost complete conversion of the glycal occurred after 45 min providing the C-glycosyl derivative **15** stereoselectively, in high yield. The exclusive formation of the β -glycoside **15** can be explained by the preferred axial attack on the D-glucopyranosyl radical.³⁴ Then, reactivity of **14** toward ethanethiol was also tested. Fast and complete reaction was observed to afford the β -configured **16** in 93% yield. It is important to note, that sulfides **15** and **16** had a great propensity for suffering oxidation upon standing in air, especially in the presence of ethyl acetate. When the crude **15** was chromatographed with an ethyl acetate-containing eluent, beside the hydrothiolation product **15**, a significant amount of the sulfoxide derivative **15b** (8%) was also isolated. Therefore, ethyl acetate was avoided during work-up and purification of **15** and **16**.


Table 3. Hydrothiolations of terminal double bonds in monosaccharide derivatives

Alkene	Thiol	Product	Yield ^a (%)
	PhSH		46
	PhSH		32
	PhSH		74 ^b 61 ^c
			8 ^c (R,S)
	EtSH		93 ^d

^aYield of isolated compounds after 3 x 15 min irradiation under Ar, using 3 x 0.1 equiv. DPAP and 2 equiv. of thiol; ^bpurified by column chromatography in 98:2 CH₂Cl₂:acetone; ^ccolumn chromatography of the reaction mixture of **15** in 75:25 *n*-hexane-EtOAc; ^d15 min irradiation, 0.1equiv. DPAP, 5 equiv. of EtSH.

Finally, we tested the applicability of the 2,3-unsaturated *N*-acetylneuraminic acid (Neu5Ac-2-ene) derivative **17**,³⁵ an intermediate in the synthesis of the influenza neuraminidase inhibitor Relenza, in the thiol-ene coupling reaction. We envisioned the synthesis of an *S*-linked mimetic of the natural α (2 $\rightarrow$ 6)-linked sialyl-galactoside structure by hydrothiolation of **17** with 1,2:3,4-di-*O*-isopropylidene-6-thio- α -D-galactopyranose **18**.³⁶ Previously, thiol **18** was applied successfully for hydrothiolation of the 2-acetoxy-glycal **1** to afford the corresponding *S*-disaccharide in 69% yield.¹⁶ In the present case, reaction of **17** and the galactose-6-thiol derivative **18** led to ~50-60% consumption of the glycal, estimated by tlc,³⁷ and resulted in a mixture of stereoisomers from which only the main product **19** could be isolated in 23% yield (Scheme 3). The low conversion could be a result of the lower electron-density of the double bond in Neu5Ac-2-ene possessing the electron-withdrawing carboxyl moiety. The equatorial orientation of the C-1 group and axial orientation of the *S*-linkage were evidenced by crosspeaks

between H-3 and both H-2 and H-4 hydrogens of the sialyl residue appeared in the ROESY spectrum of **19**. Formation of the pseudodisaccharide **19** as the main product can be explained by preferred attack on the less substituted carbon at the less hindered side in the thiyl radical addition step and the preferred axial attack on the C-2 radical in the hydrogen abstraction step. Despite the moderate yield, the approach allowing a simple one-step transformation of the easily available **17** into 2-S-linked sialomimetics, stable analogues of the biorelevant *N*-acetylneuraminic acid glycosides,³⁸ is worthy of further investigation.



Scheme 4. Reagents and conditions: 2 equiv of **18**, 5 x 15 min irradiation, 5 x 0.1 equiv. DPAP, 23% for **19**; unreacted **17** was recovered with 35% yield.

Conclusions

Photoinduced radical-mediated hydrothiolation of acetyl or benzyl-protected endoglycals and monosaccharides bearing an exocyclic double bond at C1-, C3- or C6-position with a range of thiols was studied. In most cases, the thiol-ene coupling reactions took place with full regio- and stereoselectivities, however the conversions highly varied. It was found that the overall conversion of a given ene substantially depended on the stability of the thiyl radical, most probably due to the reversible nature of the radical addition step. Thus, inverse proportionality between the stability of the thiyl radical and conversion of the reactants was observed. The highly resonance stabilized thiophenyl radical proved to be especially useful for testing the reactivity of the different enes in the thiol-ene coupling reactions. Attempted reaction between thiophenol and either 2-acetoxy-3,4,6-tri-*O*-acetyl-D-glucal **1** or 3,4,6-tri-*O*-acetyl-D-glucal **6** failed even by applying 50 equivalents of the thiol. This hydrothiolation agent displayed moderate reactivity towards both the electron-rich internal double bond of the benzylated 2-hydroxy-glycal **4**, and the electron-poor terminal double bonds of exoglycals **10** and **12** furnishing the final products in 21-46% yields. The reaction of thiophenol and exoglycal **14** afforded the corresponding β -C-glycoside in 74% yield. In this case the highly reactive electron-rich terminal double bond of **14** captured the thiyl radical and, therefore, forced the equilibrium of the thiyl radical addition step

towards the hydrogen abstraction step affording the product in high yield.

Successful hydrothiolation of a 2,3-unsaturated *N*-acetylneuraminic acid derivative revealed that the endocyclic double bond of Neu5Ac-2-ene, bearing an electron-withdrawing substituent, exhibits sufficient reactivity in thiol-ene coupling reaction.

Experimental section

General Information

Ethanethiol, 2-methylpropane-2-thiol, thiophenol, phenylmethanethiol, (4-methoxyphenyl)methanethiol, sodium 2-mercaptoethanesulfonate and 1-thioglycerol were purchased from Sigma Aldrich Chemical Co. Optical rotations were measured at room temperature with a Perkin-Elmer 241 automatic polarimeter. TLC was performed on Kieselgel 60 F254 (Merck) with detection by immersing into 5 % ethanolic sulfuric acid soln. followed by heating. Column chromatography was performed on Silica gel 60 (Merck 0.063-0.200 mm). Organic solutions were dried over MgSO_4 , and concentrated in vacuum. The ^1H (360, 400 and 500 MHz) and ^{13}C NMR (90.54, 100.28 and 125.76 MHz) spectra were recorded with Bruker DRX-360, Bruker DRX-400 and Bruker Avance II 500 spectrometers. Chemical shifts are referenced to Me_4Si or DSS (0.00 ppm for ^1H) and to the residual solvent signals (CDCl_3 : 77.00 ppm, CD_3OD : 49.15 ppm, $\text{DMSO}-d_6$: 39.52 ppm for ^{13}C). The coupling constant values (J) are given in Hz. Elemental analyses (C, H, S, N) were performed using an Elementar Vario MicroCube instrument. The photocatalytic reactions were carried out at room temperature by irradiation with a Hg-lamp with a borosilicate vessel giving maximum emission at 365 nm.

General method A for the photoinduced addition of ethanethiol and 2-methylpropane-2-thiol to glycals (**1**, **4**, **6** and **14**)

To a solution of the starting glycal (1.00 mmol) in dry toluene (7 mL) ethanethiol/2-methylpropane-2-thiol (5.00 mmol) and 2,2-dimethoxy-2-phenylacetophenone (DPAP, 25 mg, 0.10 mmol) were added. The solution was deoxygenated by argon bubbling and irradiated at room temperature for 15 min. The addition of DPAP and the thiol, deoxygenation and irradiation were repeated twice more (except for the reaction of **14** and ethanethiol which showed complete conversion of the glycal after 15 min). Then the solution was concentrated and the residue was purified by column chromatography.

General method B for the photoinduced addition of thiols to endo- and exocyclic double bonds of monosaccharide derivatives

To a solution of the starting unsaturated monosaccharide (1.00 mmol) in dry toluene^a (7 mL) thiol (2.0-4.0 equiv) and 2,2-dimethoxy-2-phenylacetophenone (DPAP, 25 mg, 0.10 mmol) were added. The solution was deoxygenated and irradiated at room temperature for 15 min. Addition of DPAP and irradiation were repeated twice more. Then the solution was concentrated^b and the residue was purified by column chromatography.

^aIt is indicated when other solvent was used.

^bWhen thiophenol was used as the reagent, the solution was diluted with CH_2Cl_2 , washed with 1M aq NaOH, dried over MgSO_4 , concentrated and purified.

**Phenylmethyl
glucopyranoside (3c)**

2,3,4,6-tetra-O-acetyl-1-thio- α -D-

Compound **1** (330 mg, 1.00 mmol) and phenylmethanethiol (**2c** 2x235 μ L, 4.00 mmol) were reacted according to general method **B** using 4 irradiation cycles. The crude product was purified by column chromatography (97:3 CH₂Cl₂-acetone) to give **3c** (160 mg, 35%) as a syrup. [α]_D²² +211.5 (c 0.68 in MeOH), (lit.³⁹ [α]_D +190), *R*_f 0.30 (97:3 CH₂Cl₂-acetone); Elemental Analysis: Found: C, 54.6; H, 5.7, S, 7.0. Calc. for C₂₄H₂₆O₉S: C, 54.5; H, 5.8, S, 7.1%; ¹H NMR (360 MHz, CDCl₃, Me₄Si): δ (ppm) 2.01, 2.02, 2.03, 2.10 (12 H, 4 x s, 4 x COCH₃), 3.72 (2 H, ABq, *J* 13.5, SCH₂), 3.87 (1 H, dd, *J*_{5,6A} 1.8, *J*_{6AB} 12.3, 6-H_A), 4.24 (1 H, dd, *J*_{5,6B} 4.6, *J*_{6AB} 12.3, 6-H_B), 4.36-4.38 (1 H, m, 5-H), 5.00-5.07 (2 H, m, 2-H, 4-H), 5.37 (1 H, t, *J* 9.7, 3-H), 5.53 (1H, d, *J*_{1,2} 5.8, 1-H), 7.19-7.35 (5 H, m, arom.); ¹³C NMR (90 MHz, CDCl₃): δ (ppm) 20.5, 20.5, 20.5, 20.6 (4 x COCH₃), 33.9 (SCH₂), 61.6 (C-6), 67.6, 68.4, 70.3, 70.5 (C-2, C-3, C-4, C-5), 81.0 (C-1), 127.2, 128.5, 128.7, 137.1 (6 C, arom.), 169.5, 169.5, 169.8, 170.4 (4 x CO).

Unreacted **1** was recovered from the reaction mixture (182 mg, 55%)

**(4-Methoxyphenyl)methyl
glucopyranoside (3d)**

2,3,4,6-tetra-O-acetyl-1-thio- α -D-

Compound **1** (330 mg, 1.00 mmol) and (4-methoxyphenyl)methanethiol (2x280 μ L, 4.0 mmol) were reacted according to general method **B** using 4 irradiation cycles. The crude product was purified by column chromatography (99:1 CH₂Cl₂-acetone) to give **3d** (49 mg, 10%) as a syrup. [α]_D²² +197.1 (c 0.51 in CHCl₃), *R*_f 0.33 (99:1 CH₂Cl₂-acetone); Elemental Analysis: Found: C, 53.9; H, 5.8, S, 6.65. Calc. for C₂₂H₂₈O₁₀S: C, 54.5; H, 5.8, S, 6.6%; ¹H NMR (360 MHz, CDCl₃, Me₄Si): δ (ppm) 2.00, 2.01, 2.03, 2.10 (12 H, 4 x s, 4 x COCH₃), 3.67 (2 H, ABq, *J* 13.4, SCH₂), 3.80 (3 H, s, OCH₃), 3.96 (1 H, dd, *J*_{5,6A} 2.0, *J*_{6AB} 12.3, 6-H_A), 4.24 (1 H, dd, *J*_{5,6B} 4.6, *J*_{6AB} 12.3, 6-H_B), 4.38-4.41 (1 H, m, 5-H), 4.99-5.06 (2 H, m, 2-H, 4-H), 5.37 (1 H, t, *J* 9.8, 3-H), 5.51 (1H, d, *J*_{1,2} 5.8, 1-H), 6.83 (2 H, d, *J* 8.6, arom.), 7.21(2 H, d, *J* 8.6, arom.). ¹³C NMR (90 MHz, CDCl₃): δ (ppm) 20.6, 20.6, 20.6, 20.7 (4 x COCH₃), 33.3 (SCH₂), 55.3 (OCH₃), 61.9 (C-6), 67.7, 68.6, 70.4, 70.7 (C-2, C-3, C-4, C-5), 81.0 (C-1), 114.0, 129.0, 130.0, 158.9 (6 C, arom.), 169.6, 169.6, 169.9, 170.6 (4 x CO).

Unreacted **1** was recovered from the reaction mixture (197 mg, 60%)

2-Methylpropane-2-yl

2,3,4,6-tetra-O-acetyl-1-thio- α -D-

glucopyranoside (3e)

Compound **1** (165 mg, 0.500 mmol) and thiol (**2e**, 3 x 281 μ L, 7.50 mmol) were reacted in toluene (5 mL) according to general method **A** using 4 irradiation cycles. The crude product was purified by silica gel chromatography in 7:3 *n*-hexane-EtOAc, to give **3e** (52 mg, 25%) as white needles. mp 73-74 °C (from *n*-hexane-EtOAc) (lit.⁴⁰ mp 63-65 °C); [α]_D²² +154.4 (c 0.30 in CHCl₃), (lit.⁴⁰ [α]_D +185); *R*_f 0.60 (6:4 *n*-hexane-EtOAc); Elemental Analysis: Found: C, 51.6; H, 6.6; S, 7.7. Calc. for C₁₈H₂₈O₉S: C, 51.4; H, 6.7; S, 7.6%; ¹H NMR (360 MHz, CDCl₃, Me₄Si): δ (ppm) 1.35 (9 H, s, 3 x SCCH₃), 2.02, 2.03, 2.06, 2.07 (12 H, 4 x s, 4 x COCH₃), 4.05 (1 H, dd, *J* 2.0, *J* 12.3), 4.30 (1 H, dd, *J* 4.7, *J* 12.3), 4.46-4.51 (1 H, m), 4.95 (1 H, dd, *J* 5.9, *J* 10.5), 5.03 (1 H, t, *J* 9.7), 5.28 (1 H, t, *J* 9.9), 5.86 (1 H, d, *J*_{1,2} 5.9, 1-H);

¹³C NMR (90 MHz, CDCl₃): δ (ppm) 20.6, 20.6, 20.6, 20.8 (4 x COCH₃), 31.3 (3 x SCCH₃), 44.4 (Cq), 61.9 (C-6), 67.6, 68.6, 70.5, 70.8 (C-2, C-3, C-4, C-5), 80.1 (C-1), 169.6, 169.8, 169.9, 170.6 (4 x CO).

Unreacted **1** was recovered from the reaction mixture (97 mg, 59%)

[(2*R,S*)-2,3-Di-O-acetoxy]propyl 2,3,4,6-tetra-O-acetyl-1-thio- α -D-glucopyranoside (3f)

Compound **1** (330 mg, 1.00 mmol) and thiol (**2f**, 384 mg, 2.00 mmol) were reacted in toluene (7 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 6:4 *n*-hexane-EtOAc, to give **3f** (351 mg, 67%) as a colourless syrup; [α]_D²² +132.5 (c 0.82 in CHCl₃); *R*_f 0.32 (6:4 *n*-hexane-EtOAc); Elemental Analysis: Found: C, 48.3; H, 5.85; S, 6.1. Calc. for C₂₁H₃₀O₁₃S: C, 48.3; H, 5.8; S, 6.1%; ¹H NMR (360 MHz, CDCl₃, Me₄Si): δ (ppm) 2.02, 2.04, 2.07, 2.09 (18 H, 4 x br s, 6 x CH₃), 2.71-2.77 (1 H, m), 2.85-2.92 (1 H, m), 4.07-4.14 (2 H, m), 4.28-4.40 (3 H, m), 4.99-5.08 (2 H, m), 5.13-5.22 (1 H, m), 5.28-5.35 (1 H, m), 5.70, 5.75 (1 H, 2 x d, *J* 5.6, *J* 5.6); ¹³C NMR (90 MHz, CDCl₃): δ (ppm) 20.1, 20.3, 20.3 (6 x CH₃), 29.6, 29.8 (SCH₂), 61.3, 61.4 (OCH₂), 63.1, 63.3 (C-6), 67.4, 67.6, 67.8, 67.9, 69.4, 69.7, 69.9, 70.1, 70.6 (C-2, C-3, C-4, C-5, C-2-propyl), 81.8, 82.4 (C-1), 169.0, 169.1, 169.3, 169.4, 169.6, 169.9 (6 x CO).

[(4*R,S*)-2,2-Dimethyl-1,3-dioxolane-4-yl]methyl 2,3,4,6-tetra-O-acetyl-1-thio- α -D-glucopyranoside (3g)

Compound **1** (165 mg, 0.500 mmol) and thiol (**2g**, 148 mg, 1.00 mmol) were reacted in toluene (5 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 97:3 CH₂Cl₂-acetone, to give **3g** (74 mg, 31%) as a colourless syrup; [α]_D²² +141.6 (c 0.43 in CHCl₃); *R*_f 0.42 (95:5 CH₂Cl₂-acetone); Elemental Analysis: Found: C, 50.1; H, 6.35; S, 6.5. Calc. for C₂₀H₃₀O₁₁S: C, 50.2; H, 6.3; S, 6.7%; ¹H NMR (360 MHz, CDCl₃, Me₄Si): δ (ppm) 1.35, 1.42, 1.43 (6 H, 3 x s, 2 x CqCH₃), 2.02, 2.04, 2.04, 2.07, 2.10 (12 H, 5 x br s, 4 x COCH₃), 2.57-2.85 (2 H, m), 3.65-3.75 (1 H, m), 4.06-4.10 (2 H, m), 4.22-4.33 (2 H, m), 4.38-4.46 (1 H, m), 5.01-5.08 (2 H, m), 5.34-5.40 (1 H, m), 5.70, 5.75 (1 H, 2 x d, *J* 5.7, *J* 5.8); ¹³C NMR (90 MHz, CDCl₃): δ (ppm) 20.5, 20.6, 20.6 (4 x COCH₃), 25.4, 25.5, 26.7, 26.8 (2 x CqCH₃), 32.8, 32.9 (SCH₂), 61.8, 61.9 (OCH₂), 67.7, 67.7, 68.4, 68.5, 70.2, 70.5, 74.8, 75.2 (C-2, C-3, C-4, C-5, C-4-dioxolane), 68.7 (C-6), 81.9, 82.7 (C-1), 109.6, 109.7 (Cq), 169.5, 169.7, 169.8, 170.5 (4 x CO).

Unreacted **1** was recovered from the reaction mixture (93 mg, 57%)

Sodium sulfonatoethyl 2,3,4,6-tetra-O-acetyl-1-thio- α -D-glucopyranoside (3h)

Compound **1** (165 mg, 0.500 mmol) and thiol (**2h**, 164 mg, 1.00 mmol) were reacted in MeOH (5 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 75:25 CH₂Cl₂-MeOH, to give **3h** (158 mg, 64%) as a colourless syrup; [α]_D²² +126.1 (c 0.28 in MeOH); *R*_f 0.31 (8:2 CH₂Cl₂-MeOH); Elemental Analysis: Found: C, 38.7; H, 5.0; S, 13.4. Calc. for C₁₆H₂₃NaO₁₂S₂: C, 38.9; H, 4.7; S, 13.0%; ¹H NMR (360 MHz, CDOD₃, Me₄Si): δ (ppm) 2.00, 2.03, 2.04, 2.09 (12 H, 4 x s, 4 x CH₃), 2.90-3.16 (4 H, m, SCH₂, CH₂S), 4.12 (1 H, dd, *J* 1.6, *J* 12.2), 4.27 (1 H, dd, *J* 5.9, *J* 12.2), 4.43-4.47 (1 H, m), 4.99-5.07 (2 H, m), 5.30 (1 H, t, *J* 9.8), 5.75

- (1 H, d, J 5.6, 1-H); ^{13}C NMR (90 MHz, CDOD_3): δ (ppm) 20.6, 20.6, 20.6, 20.8 (4 x CH_3), 26.4 (CH_2S), 52.9 (NaO_3SCH_2), 63.5 (C-6), 69.2, 70.0, 71.6, 71.9 (C-2, C-3, C-4, C-5), 83.7 (C-1), 171.3, 171.3, 171.5, 172.6 (4 x CO).
- 5 Ethyl 2,3,4,6-tetra-*O*-benzyl-1-thio- α -D-glucopyranoside (5a)**
Compound **4** (261 mg, 0.500 mmol) and thiol (**2a**, 3 x 185 μL , 7.50 mmol) were reacted in toluene (5 mL) according to general method **A**. The crude product was purified by silica gel chromatography in 85:15 *n*-hexane-EtOAc, to give **5a** (207 mg, 71%) as white crystals. mp 86-87 °C (from EtOH), (lit.⁴¹ mp 88-90 °C); $[\alpha]_{\text{D}}^{22} +96.1$ (c 0.56 in CHCl_3); (lit.⁴¹ $[\alpha]_{\text{D}} +108$); R_f 0.60 (8:2 *n*-hexane-EtOAc); Elemental analysis: (Found: C, 73.9; H, 6.8; S, 5.6. Calc. for $\text{C}_{36}\text{H}_{40}\text{O}_5\text{S}$: C, 73.9; H, 6.9; S, 5.5%); ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 1.27 (3 H, t, J 7.4, CH_3), 2.45-2.63 (2 H, m, CH_2), 3.61-3.67 (2 H, m), 3.76 (1 H, dd, J 3.5, J 10.7), 3.81-3.89 (2 H, m), 4.19 (1 H, br d, J 9.7), 4.44, 4.47, 4.60, 4.65, 4.73, 4.76, 4.83, 4.95 (8 x 1 H, 8 x d, J 12.1, J 10.6, J 12.1, J 11.8, J 12.2, J 11.0, J 10.8, J 10.8, 4 x CH_2Ph), 5.41 (1 H, d, $J_{1,2}$ 4.4, 1-H), 7.12-7.39 (20 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 14.7 (CH_3), 23.6 (CH_2), 68.5 (C-6), 70.4, 77.4, 79.4, 82.5, 83.0 (C-1, C-2, C-3, C-4, C-5), 72.3, 73.4, 74.9, 75.6 (4 x CH_2Ph), 127.6-138.7 (24 C, arom).
- Phenyl 2,3,4,6-tetra-*O*-benzyl-1-thio- α -D-glucopyranoside (5b)**
Compound **4** (261 mg, 0.500 mmol) and thiol (**2b**, 102 μL , 1.00 mmol) were reacted in toluene (5 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 99:1 CH_2Cl_2 -EtOAc, to give **5b** (67 mg, 21%) as a white needles. mp 77-78 °C (from EtOH) (lit.⁴² mp 81-82 °C); $[\alpha]_{\text{D}}^{22} +142.7$ (c 0.33 in CHCl_3), (lit.⁴² $[\alpha]_{\text{D}} +154$); R_f 0.58 (8:2 *n*-hexane-EtOAc); Elemental Analysis: Found: C, 76.1; H, 6.5; S, 5.0. Calc. for $\text{C}_{40}\text{H}_{40}\text{O}_5\text{S}$: C, 75.9; H, 6.4; S, 5.1%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 3.61 (1 H, br d, J 10.2), 3.66-3.72 (1 H, m), 3.77 (1 H, dd, J 3.7, J 10.7), 3.86-3.93 (2 H, m), 4.33 (1 H, br d, J 9.8), 4.41, 4.49, 4.58, 4.68, 4.76, 4.81, 4.85, 5.00 (8 x 1 H, 8 x d, J 12.0, J 10.8, J 12.0, J 11.7, J 11.8, J 10.9, J 10.8, J 10.8, 4 x CH_2Ph), 5.64 (1 H, d, $J_{1,2}$ 3.7, 1-H), 7.14-7.50 (25 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 68.5 (C-6), 71.1, 77.3, 79.7, 82.5 (C-2, C-3, C-4, C-5), 72.5, 73.3, 75.1, 75.7 (4 x CH_2Ph), 87.0 (C-1), 127.0-138.6 (30 C, arom).
- 2-Methylpropane-2-yl 2,3,4,6-tetra-*O*-benzyl-1-thio- α -D-glucopyranoside (5e)**
Compound **4** (261 mg, 0.500 mmol) and thiol (**2e**, 3 x 281 μL , 7.50 mmol) were reacted in toluene (5 mL) according to general method **A**. The crude product was purified by silica gel chromatography in 88:12 *n*-hexane-EtOAc, to give **5e** (122 mg, 40%) as white needles. mp 91-92 °C (from EtOH) (lit.⁴³ mp 97-98 °C); $[\alpha]_{\text{D}}^{22} +72.9$ (c 0.40 in CHCl_3), (lit.⁴³ $[\alpha]_{\text{D}} +120$); R_f 0.55 (85:15 *n*-hexane-EtOAc); Elemental Analysis: Found: C, 74.4; H, 7.1; S, 5.2. Calc. for $\text{C}_{38}\text{H}_{44}\text{O}_5\text{S}$: C, 74.5; H, 7.2; S, 5.2%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 1.38 (9 H, s, 3 x CH_3), 3.57 (1 H, dd, J 1.9, J 10.6), 3.64-3.69 (1 H, br t, J 8.6, J 9.7), 3.74-3.83 (3 H, m), 4.25-4.28 (1 H, m), 4.41, 4.45, 4.62 (3 x 1 H, 3 x d, J 12.2, J 11.0, J 12.1, CH_2Ph), 4.70 (2 H, s, CH_2Ph), 4.75, 4.83, 4.96 (3 x 1 H, 3 x d, 3 x J 10.8, CH_2Ph), 5.54 (1 H, d, $J_{1,2}$ 4.8, 1-H), 7.12-7.39 (20 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 31.5 (3 x CH_3), 43.7 (Cq), 68.4 (C-6), 70.4, 77.5, 79.4, 81.7, 82.9 (C-1, C-2, C-3, C-4, C-5), 72.4, 73.4, 74.9, 75.7 (4 x CH_2Ph), 127.5-138.7 (24 C, arom).
- [(2*R,S*)-2,3-Dihydroxy]propyl 2,3,4,6-tetra-*O*-benzyl-1-thio- α -D-glucopyranoside (5i)**
Compound **4** (261 mg, 0.500 mmol) and thiol (**2i**, 87 μL , 1.00 mmol) were reacted in toluene (5 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 1:1 CH_2Cl_2 -EtOAc, to give **5i** (144 mg, 46%) as white crystals. mp 60-61 °C (from EtOH); $[\alpha]_{\text{D}}^{22} +71.2$ (c 0.58 in CHCl_3); R_f 0.51 (1:1 CH_2Cl_2 -EtOAc); Elemental Analysis: Found: C, 70.2; H, 6.6; S, 4.9. Calc. for $\text{C}_{37}\text{H}_{42}\text{O}_7\text{S}$: C, 70.45; H, 6.7; S, 5.1%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 2.50-2.79 (3 H, m), 3.38-3.85 (9 H, m), 4.20-4.27 (1 H, m), 4.42-4.57 (3 H, m, CH_2Ph), 4.64-4.77 (3 H, m, CH_2Ph), 4.82, 4.94 (2 x 1 H, 2 x d, J 10.9, J 10.8, CH_2Ph), 5.33 (1 H, d, $J_{1,2}$ 4.2, 1-H), 7.13-7.37 (20 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 34.3 (SCH_2), 65.2 (OCH_2), 68.6, 68.7 (C-6), 70.7, 70.8, 71.2, 77.4, 79.3, 79.4, 82.2, 84.3, 84.9 (C-1, C-2, C-3, C-4, C-5, C-2'), 72.5, 72.5, 73.3, 73.3, 74.9, 75.6 (4 x CH_2Ph), 127.5-138.4 (24 C, arom).
- 1,2:5,6-Di-*O*-isopropylidene-3-*S*-(2,3,4,6-tetra-*O*-benzyl- α -D-glucopyranosyl)-3-thio- α -D-glucofuranose (5j)**
Compound **4** (261 mg, 0.500 mmol) and thiol (**2j**, 276 mg, 1.00 mmol) were reacted in toluene (5 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 85:15 *n*-hexane-EtOAc, to give **5j** (235 mg, 59%) as white crystals. mp 89-90 °C (from EtOH). $[\alpha]_{\text{D}}^{22} +89.0$ (c 0.73 in CHCl_3); R_f 0.32 (85:15 *n*-hexane-EtOAc); Elemental Analysis: Found: C, 69.0; H, 6.8; S, 3.9. Calc. for $\text{C}_{46}\text{H}_{54}\text{O}_{10}\text{S}$: C, 69.15; H, 6.8; S, 4.0%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 1.24, 1.31, 1.43, 1.50 (12 H, 4 x s, 4 x CH_3), 3.58-3.90 (6 H, m), 4.03-4.23 (4 H, m), 4.44-4.49 (2 H, m), 4.54-4.64 (3 H, m), 4.72-4.85 (4 H, m), 4.94 (1 H, d, J 10.8), 5.77 (1 H, d, $J_{1,2}$ 3.2, 1-H), 5.84 (1 H, d, $J_{1,2'}$ 4.7, 1'-H), 7.11-7.38 (20 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 25.3, 26.0, 26.5, 26.9 (4 x CH_3), 50.0 (C-3), 67.7 (C-6), 68.3 (C-6'), 71.0, 74.4, 77.1, 79.1, 80.4, 82.0, 84.4 (C-2, C-4, C-5, C-2', C-3', C-4', C-5'), 71.5, 73.3, 74.9, 75.5 (4 x CH_2Ph), 86.4 (C-1'), 104.7 (C-1), 109.2, 111.7 (2 x Cq), 127.4-138.4 (24 C, arom).
- 3,4,6-Tri-*O*-acetyl-1,5-anhydro-2-*S*-ethyl-D-mannitol (7), 3,4,6-tri-*O*-acetyl-1,5-anhydro-2-*S*-ethyl-D-glucitol (8) and ethyl 3,4,6-tri-*O*-acetyl-2-deoxy-1-thio- α -D-arabinohexopyranoside (9)**
Compound **6** (544 mg, 2.00 mmol) and ethanethiol (**2a**, 3 x 740 μL , 30.00 mmol) were reacted according to general method **A**, to give a mixture of **7**, **8** and **9** which were separated by silica gel chromatography in 1:1 *n*-hexane-Et₂O. (After column chromatography, compound **9** was contaminated by the unreacted starting material, from which could only be separated after catalytic hydrogenation and a subsequent purification by silica gel chromatography in 1:1 *n*-hexane-EtOAc).
- 7:** (230 mg, 34%) colourless syrup; $[\alpha]_{\text{D}}^{22} -34.4$ (c 0.61, CHCl_3); R_f 0.18 (1:1 *n*-hexane-Et₂O); Elemental Analysis: Found: C, 50.15; H, 6.5; S, 9.45. Cald. for $\text{C}_{14}\text{H}_{22}\text{O}_7\text{S}$: C, 50.3; H, 6.6; S, 9.6.; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 1.24 (3 H, t, J 7.4, CH_3CH_2), 2.06, 2.09, 2.10 (9 H, 3s, 3 x CH_3), 2.58-2.64 (2 H, q, CH_2), 3.37-3.39 (1 H, m, 2-H), 3.58-3.62 (1 H, m, 5-H), 3.83 (1 H, dd, $J_{1\beta,2}$ 2.1, $J_{1\alpha,1\beta}$ 12.2, 1 β -H), 4.09 (1 H, dd, $J_{1\alpha,2}$ 2.6,

$J_{1\alpha,1\beta}$ 12.2, 1- α H), 4.13 (1 H, dd, $J_{5,6A}$ 2.7, $J_{6A,B}$ 12.2, 6- H_A), 4.19 (1 H, dd, $J_{5,6B}$ 5.5, $J_{6A,B}$ 12.2, 6- H_B), 5.10 (1 H, dd, $J_{2,3}$ 4.4, $J_{3,4}$ 9.3, 3-H), 5.28 (1 H, t, $J_{4,5}$ 9.2, 4-H); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 14.5 (CH_3CH_2), 20.5, 20.5, 20.6 (3 x CH_3), 26.4 (CH_2), 45.3 (C-2), 62.3 (C-6), 66.5 (C-4), 69.1 (C-1), 73.7 (C-3), 76.6 (C-5), 169.2, 170.0, 170.4 (3 x CO).

8: (163 mg, 24%) colourless syrup; $[\alpha]_{\text{D}}^{22} +71.1$ (c 0.35, CHCl_3); R_f 0.30 (1:1 *n*-hexane-Et₂O); Elemental Analysis: Found: C, 50.05; H, 6.7; S, 9.4. Calcd. for $\text{C}_{14}\text{H}_{22}\text{O}_7\text{S}$: C, 50.3; H, 6.6; S, 9.6; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 1.24 (3 H, t, J 7.4, CH_3CH_2), 2.03, 2.08, 2.09 (9 H, 3s, 3 x CH_3), 2.52-2.62 (2 H, m, CH_2), 2.84-2.91 (1 H, m, 2-H), 3.40 (1 H, t, $J_{1\beta,2}$ 11.9, 1 β -H), 3.58-3.63 (1 H, m, 5-H), 4.09 (1 H, dd, $J_{5,6A}$ 2.1, $J_{6A,B}$ 12.3, 6- H_A), 4.14 (1 H, dd, $J_{1\alpha,2}$ 5.1, $J_{1\alpha,1\beta}$ 11.9, 1 α -H), 4.25 (1 H, dd, $J_{5,6B}$ 4.8, $J_{6A,B}$ 12.3, 6- H_B), 4.98-5.00 (2 H, m, 3-H, 4-H); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 15.1 (CH_3CH_2), 20.5, 20.6, 20.6 (3 x CH_3), 25.3 (CH_2), 45.1 (C-2), 62.3 (C-6), 69.5 (C-4), 70.2 (C-1), 73.9 (C-3), 76.4 (C-5), 169.6, 170.1, 170.5 (3 x CO).

9: (46 mg, 7%) white crystals, mp 50-52 °C (EtOH), (lit.⁴⁴ mp 51-52 °C); $[\alpha]_{\text{D}}^{22} +207.7$ (c 0.25, CHCl_3) (lit.⁴⁴ $[\alpha]_{\text{D}} +183$); R_f 0.36 (1:1 *n*-hexane-Et₂O). Analysis: Found: C, 50.4; H, 6.75; S, 9.6. Calcd. for $\text{C}_{14}\text{H}_{22}\text{O}_7\text{S}$: C, 50.3; H, 6.6; S, 9.6; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 1.29 (3 H, t, J 7.4, CH_3CH_2), 2.01, 2.05, 2.09 (9 H, 3s, 3 x CH_3), 2.13-2.30 (2 H, m), 2.50-2.68 (2 H, m), 2.04 (1 H, dd, J 1.8, J 11.9), 4.33-4.42 (2 H, m), 4.98 (1 H, t, J 9.5), 5.21-5.28 (1 H, m), 5.45 (1 H, d, J 5.5); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 14.7 (CH_3CH_2), 20.7, 20.7, 20.9 (3 x CH_3), 24.9 (CH_2), 35.2 (C-2), 62.3 (C-6), 68.0, 69.4, 69.6 (C-3, C-4, C-5), 79.6 (C-1), 169.9, 170.0, 170.6 (3 x CO).

30 **6-Deoxy-1,2:3,4-di-O-isopropylidene-7-S-phenyl- α -D-galactopyranose (11)**

Compound **10** (256 mg, 1.00 mmol) and thiol (**2b**, 205 μL , 2.00 mmol) were reacted in toluene (7 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 98:2 CH_2Cl_2 -EtOAc, to give **11** (168 mg, 46%) as a colourless syrup. $[\alpha]_{\text{D}}^{22} -15.3$ (c 0.30 in CHCl_3); R_f 0.49 (99:1 CH_2Cl_2 -acetone); Elemental Analysis: Found: C, 62.4; H, 7.2; S, 8.65. Calcd. for $\text{C}_{19}\text{H}_{26}\text{O}_5\text{S}$: C, 62.3; H, 7.15; S, 8.75%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 1.32, 1.33, 1.44, 1.59 (12 H, 4 x s, 4 x CH_3), 1.73-1.82 (1 H, m, 6- H_A), 2.01-2.11 (1 H, m, 6- H_B), 2.95-3.03 (1 H, m, 7- H_A), 3.11-3.18 (1 H, m, 7- H_B), 3.99-4.04 (1 H, m, 5-H), 4.09 (1 H, dd, $J_{4,5}$ 1.6, $J_{3,4}$ 7.9, 4-H), 4.30 (1 H, dd, $J_{2,3}$ 2.3, $J_{1,2}$ 5.0, 2-H), 4.59 (1 H, dd, $J_{2,3}$ 2.3, $J_{3,4}$ 7.9, 3-H), 5.52 (1 H, d, $J_{1,2}$ 5.0, 1-H), 7.12-7.33 (5 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 24.3, 24.9, 25.9, 26.0 (4 x CH_3), 29.3, 29.6 (C-6, C-7), 65.6, 70.4, 70.8, 72.8 (C-2, C-3, C-4, C-5), 96.4 (C-1), 108.5, 109.0 (2 x Cq), 125.7, 128.7, 129.0 (5 C, arom), 136.2 (Cq, arom).

3-Deoxy-3-C-(phenylthiomethyl)-1,2:5,6-Di-O-isopropylidene- α -D-allofuranose (13)

Compound **12** (256 mg, 1.00 mmol) and thiol (**2b**, 205 μL , 2.00 mmol) were reacted in toluene (5 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 98:1:1 CH_2Cl_2 -EtOAc-TEA, to give **13** (117 mg, 32%) as white crystals. mp 41-42 °C; $[\alpha]_{\text{D}}^{22} +99.0$ (c 0.42 in CHCl_3); R_f 0.42 (98:2 CH_2Cl_2 -EtOAc); Elemental Analysis: Found: C, 62.05; H, 7.3; S, 8.9. Calcd. for $\text{C}_{19}\text{H}_{26}\text{O}_5\text{S}$: C, 62.30; H, 7.15; S, 8.75%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm)

1.32, 1.37, 1.42, 1.53 (12 H, 4 x s, 4 x CH_3), 2.13-2.21 (1 H, m, 3-H), 3.09 (1 H, dd, $J_{3,\text{SCH}_2\text{A}}$ 11.5, $J_{\text{SCH}_2\text{AB}}$ 13.5, SCH_2A), 3.52 (1 H, dd, $J_{3,\text{SCH}_2\text{B}}$ 3.2, $J_{\text{SCH}_2\text{AB}}$ 13.5, SCH_2B), 3.77 (1 H, dd, $J_{3,4}$ 9.7, $J_{4,5}$ 7.5, 4-H), 3.89 (1 H, dd, $J_{5,6A}$ 5.6, J_{6AB} 8.2, 6- H_A), 3.94-3.99 (1 H, m, 5-H), 4.09 (1 H, dd, $J_{5,6B}$ 5.9, $J_{6A,B}$ 8.2, 6- H_B), 4.76 (1 H, t, J 3.6, 2-H), 5.71 (1 H, d, $J_{1,2}$ 3.6, 1-H), 7.14-7.40 (5 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 25.3, 26.3, 26.6, 26.8 (4 x CH_3), 27.9 (SCH_2), 48.8 (C-3), 67.9 (C-6), 76.7 (C-5), 80.7 (C-4), 81.0 (C-2), 104.8 (C-1), 109.6, 112.0 (2 x Cq), 125.5, 128.2, 128.8 (5 C, arom), 136.3 (Cq, arom).

1-S-Phenyl-2,6-anhydro-1-deoxy-3,4,5,7-tetra-O-benzoyl-D-glycero-D-gulo-heptitol (15) and 1-(phenylsulfinyl)-2,6-anhydro-1-deoxy-3,4,5,7-tetra-O-benzoyl-D-glycero-D-gulo-heptitol (15b)

Compound **14** (225 mg, 0.380 mmol) and thiol (**2b**, 78 μL , 0.760 mmol) were reacted in toluene (4 mL) according to general method **B**. The crude product was purified by silica gel chromatography in 98:2 CH_2Cl_2 -acetone, to give **15** (197 mg, 74%) as a colourless syrup. $[\alpha]_{\text{D}}^{22} +4.0$ (c 0.50 in CHCl_3); R_f 0.45 (75:25 *n*-hexane-EtOAc); Elemental Analysis: Found: C, 69.9; H, 4.7; S, 4.4. Calcd. for $\text{C}_{41}\text{H}_{34}\text{O}_9\text{S}$: C, 70.1; H, 4.9; S, 4.6%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 3.16 (1 H, dd, $J_{1A,2}$ 8.0, J_{1AB} 14.3, 1- H_A), 3.27 (1 H, dd, $J_{1B,2}$ 2.7, J_{1AB} 14.3, 1- H_B), 3.97-4.03 (1 H, m, 2-H), 4.09-4.14 (1 H, m, 6-H), 4.43 (1 H, dd, $J_{6,7A}$ 5.3, J_{7AB} 12.2, 7- H_A), 4.59 (1 H, dd, $J_{6,7B}$ 2.8, J_{7AB} 12.2, 7- H_B), 5.59 (1 H, t, J 9.6), 5.70 (1 H, t, J 9.7), 5.91 (1 H, t, J 9.5), 7.08-7.54 (17 H, m, arom), 7.80-8.06 (8 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 35.8 (C-1), 63.1 (C-7), 69.6, 72.0, 74.1, 76.0, 77.8 (C-2, C-3, C-4, C-5, C-6), 126.2-135.9 (30 C, arom), 165.1, 165.3, 165.8, 166.0 (4 x CO).

When the crude **15** was purified by silica gel chromatography using 75:25 *n*-hexane-EtOAc as the eluent, the sulfoxide derivative **15b** was isolated in 8% yield and pure **15** was isolated in 61% yield.

Compound **15b**: Colourless syrup. $[\alpha]_{\text{D}}^{22} +23.5$ (c 0.47 in CHCl_3); R_f 0.56 (1:1 *n*-hexane-EtOAc); Elemental Analysis: Found: 68.2; H, 4.6; S, 4.65%. Calcd. for $\text{C}_{41}\text{H}_{34}\text{O}_{10}\text{S}$: C, 68.5; H, 4.8; S, 4.5%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 2.91-3.04 (2 H, m), 3.13 (1 H, dd, J 4.5, J 13.6), 3.34 (1 H, dd, J 7.5, J 13.6), 3.90-3.95 (1 H, m), 4.00-4.06 (1 H, m), 4.25-4.39 (3 H, m), 4.43-4.56 (2 H, m), 4.68 (1 H, dd, J 2.8, J 12.3), 5.43-5.54 (2 H, m), 5.62 (1 H, t, J 9.8), 5.73 (1 H, t, J 9.8), 5.83 (1 H, t, J 9.6), 5.96 (1 H, t, J 9.6), 7.23-8.11 (50 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 58.4, 60.8 (C-1), 63.1 (C-7), 69.3, 69.4, 71.7, 72.0, 72.1, 72.5, 73.8, 74.1, 76.1, 76.4 (C-2, C-3, C-4, C-5, C-6), 123.8-143.8 (60 C, arom), 165.1, 165.2, 165.5, 165.7, 165.7, 166.2 (8 x CO).

1-S-Ethyl-2,6-anhydro-1-deoxy-3,4,5,7-tetra-O-benzoyl-D-glycero-D-gulo-heptitol (16)

Compound **14** (140 mg, 0.236 mmol) and thiol **2a** (87 μL , 1.18 mmol) were reacted in toluene (3 mL) according to general method **A**. The crude product was purified by silica gel chromatography in 98:2 CH_2Cl_2 -acetone, to give **16** (143 mg, 93%) as a colourless syrup; $[\alpha]_{\text{D}}^{22} +23.5$ (c 0.47 in CHCl_3); R_f 0.45 (75:25 *n*-hexane-EtOAc); Elemental Analysis: Found: C, 67.65; H, 5.05; S, 5.0. Calcd. for $\text{C}_{37}\text{H}_{34}\text{O}_9\text{S}$: C, 67.9; H, 5.2; S, 4.9%; ^1H NMR (360 MHz, CDCl_3 , Me_4Si): δ (ppm) 1.14 (3 H, t, J 7.4, CH_3) 2.57-2.71 (2 H, m, CH_2S), 2.74-2.83 (2 H, m, 1- H_{AB}),

4.00–4.06 (1 H, m, 2-H), 4.14–4.19 (1 H, m, 6-H), 4.46 (1 H, dd, $J_{6,7A}$ 5.4, J_{7AB} 12.2, 7-H_A), 4.66 (1 H, dd, $J_{6,7B}$ 2.7, J_{7AB} 12.2, 7-H_B), 5.59 (1 H, t, J 9.6), 5.68 (1 H, t, J 9.7), 5.93 (1 H, t, J 9.6), 7.22–7.56 (12 H, m, arom), 7.80–8.07 (8 H, m, arom); ^{13}C NMR (90 MHz, CDCl_3): δ (ppm) 14.4 (CH_3), 27.3 (CH_2S), 32.8 (C-1), 63.2 (C-7), 69.6, 71.8, 74.2, 76.0, 80.0 (C-2, C-3, C-4, C-5, C-6), 128.1–133.3 (24 C, arom), 165.1, 165.2, 165.8, 166.0 (4 x CO). **1,2:3,4-Di-O-isopropylidene-6-S-(5-acetamido-4,7,8,9-tetra-O-acetyl-2,6-anhydro-5-deoxy-3-yl-D-glycero-D-galacto-uronic acid methyl ester)-6-thio- α -D-galactopyranose (19)** Compound **17** (473 mg, 1.00 mmol) and thiol **18** (553 mg, 2.0 mmol) were reacted according to general method **B**; the addition of DPAP (5 x 25 mg, 5 x 0.100 mmol), deoxygenation and 15 min irradiation were repeated 5 times. The crude product was separated by column chromatography in 7:3 *n*-hexane-EtOAc to two fractions. First fraction with higher mobility (230 mg, R_f 0.78) proved to be disulfide derivative of **18**.⁴⁵ The second fraction (650 mg, R_f 0.05) was a mixture of compounds with very similar mobility, including two main components. This mixture was purified again by column chromatography in 1:9 *n*-hexane-EtOAc, to give **19** (173 mg, 23%) as a colourless syrup, and unreacted glycal **17** (163 mg, 35%; R_f 0.38). Compound **19**: $[\alpha]_D^{24}$ -20.3 (c 0.25, CHCl_3); R_f 0.37 (1:9 *n*-hexane-EtOAc). Elemental Analysis: Found: C, 51.4; H, 6.3; N, 1.9; S, 4.3. Calc. for $\text{C}_{32}\text{H}_{47}\text{NO}_{17}\text{S}$: C, 51.3; H, 6.3; N, 1.9; S, 4.3%; ^1H NMR (CDCl_3 , 400 MHz): δ (ppm) 1.32, 1.40, 1.56 (12 H, 3 x s, 4 x CH_3), 1.89, 2.02, 2.05, 2.10, 2.13, 2.16 (15 H, 5 x s, 5 x CH_3), 2.69–2.74 (1 H, m, 6-H_A), 2.84–2.90 (1 H, m, 6-H_B) 3.63–3.65 (1 H, m, 3-H), 3.79 (3 H, s, OCH_3), 3.82–3.84 (2 H, m, 5'-H, 6-H), 4.16–4.22 (2 H, m, 5-H, 9-H_A), 4.27–4.29 (2 H, m, 2'-H, 4'-H), 4.36 (1 H, d, $J_{2,3}$ 1.1, 2-H), 4.57–4.59 (1 H, m, 3'-H), 4.66–4.70 (1 H, m, 9-H_B), 5.26–5.30 (2 H, m, 7-H, 8-H), 5.46 (1 H, d, $J_{1,2}$ 4.7, 1'-H), 5.66 (1 H, d, J 9.3, NH); ^{13}C NMR (CDCl_3 , 100 MHz): δ (ppm) 20.7, 20.8, 20.9 (4 x CH_3), 23.2 (CH_3), 24.3, 24.8, 25.9 (4 x CH_3), 32.4 (C-6'), 47.1 (C-5), 48.7 (C-3), 52.4 (C-1), 62.4 (C-9), 66.9 (C-5'), 68.3 (C-7), 70.4, 70.6, 70.7 (C-2', C-3', C-4'), 71.6 (C-8), 73.1 (C-4), 78.0 (C-2), 96.5 (C-1'), 109.1, 108.6 (2 x Cq), 167.4, 170.2, 170.3, 170.5, 170.6 (6 x CO).

Notes and references

^aResearch Group for Carbohydrates, University of Debrecen, PO Box 94, H-4010 Debrecen, Hungary. Fax:3652 512914;

^bDepartment of Pharmaceutical Chemistry, Medical and Health Science Center, University of Debrecen, PO Box 70, H-4010 Debrecen, Hungary.

⁴⁵ Fax:3652 512914; E-mail: borbas.aniko@pharm.unideb.hu

^cDepartment of Organic Chemistry, University of Debrecen, PO Box 20, H-4010 Debrecen, Hungary.

^dDepartment of Genetics and Applied Microbiology, University of Debrecen, 4032 Debrecen, Hungary

† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

‡ Acknowledgements

The work was supported by the TÁMOP 4.2.1/B-09/1/KONV-2010-0007 project. The project is co-financed by the European Union and the European Social Fund. Financial support of the Hungarian Research Fund (OTKA K 79126 to HP) and the Bolyai János Research Fellowship (to MT) of the Hungarian Academy of Sciences are also acknowledged.

- 1 (a) C. E. Hoyle, T. Y. Lee and T. Roper, *J. Polym. Sci. Part A: Polym. Chem.* 2004, **42**, 5301; (b) C. E. Hoyle and C. N. Bowman, *Angew. Chem. Int. Ed.* 2010, **49**, 1540.
- 2 (a) A. Dondoni, *Angew. Chem. Int. Ed.* 2008, **47**, 8995; (b) A. Dondoni and A. Marra, *Chem. Soc. Rev.* 2012, **41**, 573.
- 3 C. N. Bowman, *Macromolecules* 2003, **36**, 7964.
- 4 K. L. Killops, L. M. Campos and C. J. Hawker, *J. Am. Chem. Soc.* 2008, **130**, 5062.
- 5 P. Jonkheijm, D. Weinrich, M. Köhn, H. Engelkamp, P. C. M. Christianen, J. Kuhlmann, J. C. Maan, D. Nüsse, H. Schroeder, R. Wacker, R. Breinbauer, C. M. Niemeyer and H. Waldmann, *Angew. Chem. Int. Ed.* 2008, **47**, 4421.
- 6 M. Köhn, J. M. Benito, C. Ortiz Mellet, T. K. Lindhorst and J. M. García Fernández, *ChemBioChem* 2004, **5**, 771.
- 7 M. Fiore, A. Chambery, A. Marra and A. Dondoni, *Org. Biomol. Chem.* 2009, **7**, 3910.
- 8 (a) N. Floyd, B. Vijayakrishnan, J. R. Koepp and B. G. Davis, *Angew. Chem. Int. Ed.* 2009, **48**, 7798; (b) M. Fiore, M. Lo Conte, S. Pacifico, A. Marra and A. Dondoni, *Tetrahedron Lett.* 2011, **52**, 444.
- 9 G. Chen, S. Amajjahe and M. H. Stenzel, *Chem. Commun.* 2009, 1198.
- 10 F. Cardona and B. La Ferla, *J. Carbohydr. Chem.* 2008, **27**, 203.
- 11 A. Dondoni, A. Massi, P. Nanni and A. Roda, *Chem.–Eur. J.* 2009, **15**, 11441.
- 12 D. A. Fulton and J. F. Stoddart, *J. Org. Chem.* 2001, **66**, 8309.
- 13 R. A. Ortiz, A. E. G. Valdez, M. G. M. Aguilar and M. L. B. Duarte, *Carbohydr. Polym.* 2009, **78**, 282.
- 14 M. Fiore, A. Marra and A. Dondoni, *J. Org. Chem.* 2009, **74**, 4422.
- 15 S. Staderini, A. Chambery, A. Marra and A. Dondoni, *Tetrahedron Lett.* 2012, **53**, 702.
- 16 L. Lázár, M. Csávas, M. Herczeg, P. Herczegh and A. Borbás, *Org. Lett.* 2012, **14**, 4650.
- 17 R. Davuluri and L. M. Lerner, *Carbohydr. Res.* 1972, **22**, 345.
- 18 TLC monitoring of test reactions of **1** with **2a** showed very similar conversions after a 15 min and a 1 h of irradiation, therefore, 15 min of irradiation cycles were applied in all cases. It was found that after 15 min irradiation the 0.1 equiv of initiator was completely consumed and the next cycle of irradiation only slightly promoted the reaction without adding another portion of DPAP. Adding 3 x 0.1 equiv of DPAP proved to be more effective than adding 0.3–0.5 equiv of DPAP in one portion before starting the reaction. Our attempts to increase the conversion of glycal **1** with thiols **2b** or **2c** by applying 4 equiv of the thiols and irradiating for 6x15 min of the reaction mixtures were unsuccessful.
- 19 K. Griesbaum, *Angew. Chem. Int. Ed.* 1970, **9**, 273.
- 20 (a) D. Horton and M. L. Wolfrom, *J. Org. Chem.* 1962, **27**, 1794; (b) G. Mugunthan, Dharmarajan Sriram, Perumal Yogeewari and K. P. Ravindranathan Kartha, *Carbohydr. Res.* 2011, **346**, 2401.
- 21 M. Goubert, I. Canet and M-E. Sinibaldi, *Eur. J. Org. Chem.* 2006, 4805.
- 22 P. A. Gent and R. Gigg, *J. Chem. Soc. Perkin Trans. 1*, 1974, 1446.
- 23 Although Lindhorst and co-workers reported that benzyl-type protecting groups were labile under shortwave UV irradiation,²⁴ other reports appeared describing successful thiol-ene reaction in the presence of benzyl and *p*-methoxybenzyl protecting groups using ultraviolet irradiation at 365 nm.²⁵ We have found that after 30 min irradiation the decomposition was neglectable, longer exposure to UV light progressively decreased the yield mainly by making the chromatographic purification difficult.
- 24 C. Kieburg, M. Dubber and T. K. Lindhorst, *Synlett* 1997, 1447.
- 25 (a) M. Nitz, B. W. Purse and D. R. Bundle, *Org. Lett.* 2000, **2**, 2939; (b) A. Lubineau, H. Lortat-Jacob, O. Gavard, S. Sarrazin and D. Bonaffé, *Chem.–Eur. J.* 2004, **10**, 4265.
- 26 P. A. Risbood, T. S. Phillips and L. Goodman, *Carbohydr. Res.* 1981, **94**, 101.
- 27 Acetylated counterpart of disaccharide **5j** has been achieved with exactly the same yield by hydrothiolation of **1** with **2j** (see ref 16.) In that case, the moderate yield was the result of the moderate conversion of the less reactive glycal **1**.
- 28 Photochemical addition of ethanethiol to tri-O-acetyl-D-glucal (**6**) has previously been described by Kushida et al.,²⁹ who reported that 25 h irradiation of **6** dissolved in ethanethiol (500 mg / 18 mL) afforded a

- 1:1 mixture of **7** and **8** with 87% overall yield. The products were not characterized.
29. Y. Araki, K. Matsuura, Y. Ishido and K. Kushida, *Chem. Lett.* **1973**, 383.
 30. Dondoni et al. published a detailed study on the hydrothiolation of glycal **6** and its C3 and C4-epimers (see ref 15). They also observed the lack of stereoselectivity of the addition and the formation of 1-deoxy-S-disaccharides in nearly 1:1 mixtures of the diastereomers.
 31. J. S. Brimacombe and A. K. M. S. Kabir, *Carbohydr. Res.* 1986, **150**, 35.
 32. E. M. Acton, R. N. Goerner, H. S. Uh, K. J. Ryan, D. W. Henry, C. E. Cass and G. A. LePage, *J. Med. Chem.* 1979, **22**, 518.
 33. (a) M. Tóth and L. Somsák, *J. Chem. Soc., Perkin. Trans.1*, **2001**, 942; (b) M. Tóth, K. E. Kövér, A. Bényei and L. Somsák, *Org. Biomol. Chem.* 2003, **1**, 4039; (c) M. Tóth, S. Kun, L. Somsák, D. Goyard, in P. Kovac (Ed.) *Carbohydrate Chemistry: Proven Synthetic Methods*, CRC Press, Boca Raton, 2012, Vol. 1. pp. 367-375.
 34. (a) J. Dupuis, B. Giese, D. Ruege, H. Fischer, H. G. Korth and R. Sustman, *Angew. Chem. Int. Ed. Engl.* 1984, **23**, 896; (b) J. Gervay, T. M. Flaherty and D. Holmes, *Tetrahedron*, 1997, **53**, 16355; (c) J.-P. Praly, *Adv. Carbohydr. Chem. Biochem.* 2000, **56**, 65-151.
 35. (a) P. Meindl and H. Tuppy, *Monatsh. Chem.* 1969, **100**, 1295. (b) K. Ikeda, K. Konishi, K. Sano and K. Tanaka, *Chem. Pharm. Bull.* 2000, **48**, 163. (c) E. J. Horn and J. Gervay-Hague, *J. Org. Chem.* 2009, **74**, 4357.
 36. Q. Tian, S. Zhang, Q. Yu, M.-B. He and Y.-S. Yang, *Tetrahedron* 2007, **63**, 2142.
 37. The conversion of glycal **17** remained unchanged when higher excess of thiol was used and exposure to UV light was prolonged for 6 h. Disulfide derivative arised from homocoupling of the thiol **18** was also observed in the reaction mixture.
 38. (a) E. E. Simanek, G. J. McGarvey, J. A. Jablonowski and C.-H. Wong, *Chem. Rev.* 1998, **98**, 833; (b) M. J. Kiefel and M. von Itzstein, *Chem. Rev.* 2002, **102**, 471; (c) T. Angata and A. Varki, *Chem. Rev.* 2002, **102**, 439.
 39. T. Ogawa and M. Matsui, *Carbohydr. Res.* 1977, **54**, C17.
 40. H. B. Wood, Jr., B. Coxon, H. W. Diehl and H. G. Fletcher Jr., *J. Org. Chem.* 1964, **29**, 461.
 41. F. Weygand and H. Ziemann, *Justus Liebigs Ann. Chem.* 1962, **657**, 179.
 42. R. J. Ferrier and R. H. Furneaux, *Carbohydr. Res.* 1976, **52**, 63.
 43. M. Yanase and M. Funabashi, *J. Carbohydr. Chem.* 2000, **52**, 53.
 44. S. Sabesan and S. Neira, *J. Org. Chem.* 1991, **56**, 5468.
 45. M. Adinolfi, D. Capasso, S. Di Gaetano, A. Iadonisi, L. Leone, A. Pastore, *Org. Biomol. Chem.* 2011, **9**, 6278.