

FOR SLEEP? A PROSPECTIVE CONTROLLED TRIAL IN PATIENTS BANDAGES AFTER OTOPLASTY

Menezes
Brazilian Society

MH, Martins DM,
Neto JI, Matzenbacher CA

prospective, randomized, controlled study
the postoperative complications
use or not of a ear bandage following
population was made up of 15
who had undergone bilateral
and Furna's procedure exclusively
to resection of cartilage) for the
ears under local anesthesia from
Patients ranged in age from 7
16. Data on complications were
under normal postoperative care
All patients received after surgery
bandage only for first 24 hours, than
carefully and changed it for a ear
The ear bandage was kept for one
was randomized to use it for 24 hours
perceived benefits of bandaging have
previous study, and due to complains
hearing and itch, we proposed this
significant difference in complications
at postoperative visits. The finding
and effective the use only at night of

1335E SEVERE GRANULOMATOSUS REACTION AFTER POLYALKYLIMIDE (BIO-ALCAMID) OF LIP AUGMENTATION

Presenter: Janos Varga, PhD

Affiliation: University of Szeged

Country: Hungary

Authors: Varga J, Kis E, Mohos G, Varga A, Bende B,
Kocsis A, Kemeny L

Injectable filler materials can be valuable to aesthetic surgeons. To date, hardly any short-term and long-term complication of polyalkylimide injections (Bio-Alcamid) have been reported. We present and discuss the history and the treatment of a 29 years old patient who had such complication.

