

Stability of gamma-valerolactone under neutral, acidic, and basic conditions

Claire Yuet Yan Wong¹ · Alex Wing-Tat Choi¹ · Matthew Y. Lui¹ · Bálint Fridrich² · Attila K. Horváth³ · László T. Mika² · István T. Horváth¹

Received: 1 November 2016 / Revised: 13 November 2016 / Accepted: 15 November 2016 / Published online: 1 December 2016
© Springer Science+Business Media New York 2016

Abstract Dry gamma-valerolactone (GVL) is stable for several weeks at 150 °C and its thermal decomposition only proceeds in the presence of appropriate catalysts. Since GVL does not react with water up to 60 °C for several weeks, it could be used as a green solvent at mild conditions. At higher temperatures, GVL reacts with water to form 4-hydroxyvaleric acid (4-HVA) and reaches the equilibrium in a few days at 100 °C. Aqueous solutions of acids (HCl and H₂SO₄) catalyze the ring opening of GVL even at room temperature, which leads to the establishment of an equilibrium between GVL, water, and 4-HVA. Although the 4-HVA concentration would be below 4 mol% in the presence of acids, it could be higher than the concentration of a reagent or a catalyst precursor, not to mention a catalytically active species. The latter could be especially worrisome as 4-HVA could be an excellent bi- or even a tri-

dentate ligand for transition metals. Aqueous solution of bases (NaOH and NH₄OH) also catalyzes the reversible ring opening of GVL. While in the case of NaOH, the product is the sodium salt of 4-hydroxyvalerate, the reversible reaction of GVL, with NH₄OH results in the formation of 4-hydroxyvaleric amide. The reversible ring opening of (*S*)-GVL in the presence of HCl or NaOH has no effect on the stability of the chiral center.

Keywords Gamma-valerolactone · Water · 4-hydroxy-valeric acid · Sulfuric acid · Hydrochloric acid · Ammonium hydroxide · Solvents · Equilibrium · Stability

Introduction

Solvents are important to provide one or more liquid phases for chemical reactions and processes [1]. While some solvents are available from nature [2], even in large quantities such as water, most of the solvents are man-made from fossil resources [3]. Traditionally, solvents were developed and/or selected primarily to achieve chemical and/or physical objectives and the impacts of their widespread use on the environment and/or health were not seriously considered until the emergence of green chemistry [4]. One of the key principles of green chemistry is the elimination of solvents in chemical processes or the replacement of hazardous solvents with environmentally benign solvents. Furthermore, the development of solvent-free processes has been suggested as the best prevention, especially when either one of the substrates or the products is a liquid and can be used as the solvent. However, if a solvent is required, the selection of a green solvent with no or limited impact on health and the environment should be preferred [5].

Gamma-valerolactone (GVL) is a cyclic ester containing a methyl group at the fifth position of a dihydrofuran-2(3H)-one

Dedicated to Professor George A. Olah on the occasion of his 90th birthday.

Electronic supplementary material The online version of this article (doi:10.1007/s11224-016-0887-6) contains supplementary material, which is available to authorized users.

✉ Attila K. Horváth
horvatha@gamma.ttk.pte.hu

✉ László T. Mika
laszlo.t.mika@mail.bme.hu


✉ István T. Horváth
istvan.t.horvath@cityu.edu.hk

¹ Department of Biology and Chemistry, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong

² Department of Chemical and Environmental Process Engineering, Budapest University of Technology and Economics, Műegyetem rkp. 3, Budapest H-1111, Hungary

³ Department of Inorganic Chemistry, University of Pécs, Ifjúság útja 6, Pécs H-7624, Hungary

ring. Although the (*S*) and (*R*) enantiomers of GVL have been known [6], most investigations have used its racemic form. We have




realized 10 years ago that GVL exhibits the most important characteristics of an ideal sustainable liquid [7–10] and could be used as a green solvent [2] or a feedstock for the production of both carbon-based consumer products [8, 11] and energy [7–10, 12]. GVL is renewable, occurs naturally, and used by the food industry frequently as a food additive. It has low melting (−31 °C), high boiling (207 °C), and flash (96 °C) points, a pleasant smell for easy recognition of leaks and spills, low toxicity, and high solubility in water to assist biodegradation. In addition, its vapor pressure is low, 0.65 kPa at 25 °C, and increases five times to 3.5 kPa at 80 °C [10].

The thermal stability of GVL is remarkably high and only 1% decomposed when a mixture of GVL and N₂ (1:10 mol/mol) was heated at 600 °C in a continuous reactor with a residence time of approximately 400 ms at 0.17 MPa [13]. By increasing the temperature to 800 °C, 98% of GVL decomposed, initially to 4-pentenoic acid, 1-butene, and CO₂ (Scheme 1), followed by the formation of typical products of high temperature radical decomposition. Isomerization of 4-pentenoic acid and 1-butene as well as other fragmentation reactions was also observed [13].

Isotope labeling experiments with H₂¹⁸O have established that GVL does not react with water under neutral conditions at room temperature for 3 months or at 60 °C for 28 days [10]. Since GVL does not form an azeotrope with water [10, 14], the latter can be readily removed by distillation resulting in a less energy demanding process for the production of GVL than that of absolute ethanol [15]. Furthermore, GVL does not form a measurable amount of peroxides in a glass flask under air in weeks, making it a safe material for large scale use [16].

It was also shown by using H₂¹⁸O, that GVL undergoes a reversible reaction with water in the presence of hydrochloric acid to form 4-hydroxyvaleric acid (4-HVA) [10] (Scheme 2) and the equilibrium lies far to the left at room temperature. The presence of small amount of 4-HVA was also observed during the conversion of ¹³C₆-labeled fructose to ¹³C₅-GVL in the presence of sulfuric acid in GVL as solvent [17].



Scheme 2 Reversible reaction of GVL with water


It should be noted that either at very high temperatures [13] or in the presence of acids at lower temperatures, the isomerization of 4-pentenoic acid and 1-butene as well as other fragmentation reactions proceed readily [18–20].

Olah and Ku reported that GVL can be completely protonated with HSO₃F-SbF₅ in liquid SO₂ to form two stable isomeric species in 75:25 ratio at −80 °C (Scheme 3), indicating a hindered rotation along the protonated C=O bond [21].


The potassium salt of 4-hydroxyvalerate was prepared by the ring-opening reaction of GVL with aqueous KOH (Scheme 4), which was then used to prepare the silver salt of 4-hydroxyvalerate [22]. The formation of the sodium salt of 4-hydroxyvalerate by the reaction of GVL with NaOH was confirmed by NMR [10]. While the reaction of GVL with aqueous NH₄OH resulted in 4-hydroxyvaleric amide, other aliphatic primary and secondary amines were used to prepare 4-hydroxy-N-alkyl or -dialkyl valeric amides [23]. Similarly, GVL has reacted with tetraalkylammonium hydroxydes to form tetraalkylammonium 4-hydroxyvalerates [24].

It was shown that GVL is one of the best solvents for acid catalyzed conversion of carbohydrates to levulinic and formic acids [25]. The selective hydrogenation of levulinic acid to 4-HVA was performed in GVL–water mixture by the use of homogeneous Ru catalysts [26]. GVL was shown to be an excellent solvent for heterogeneous Pd-catalyzed Hiyama [27], Heck [28], Sonogashira [29], and Catellani [30] reactions. Synthesis of GVL in GVL from paper wastes [12] as well as from corn stover and rice husk [31] was demonstrated by a one pot system containing sulfuric acid and the Shvo catalyst. The heterogeneous catalytic C–H arylation of 1,2,3-triazoles in GVL was used in the synthesis of pharmaceuticals [32]. The application of GVL as the reaction medium for Pd-catalyzed asymmetric hydroformylation and aminocarbonylation has also been reported [33, 34].

Since GVL has become a popular green solvent [35, 36], we report our studies on the stability of GVL in neutral, basic, and acidic conditions, which should help chemist and chemical engineers to decide how to use GVL as a green solvent.



Scheme 1 Decomposition of GVL at higher temperatures



Scheme 3 Protonation of GVL by $\text{HSO}_3\text{-SbF}_5$ in liquid SO_2 at $-80\text{ }^\circ\text{C}$

Materials and methods

Gamma-valerolactone, 37.0% HCl, 98% H_2SO_4 , 25% NH_4OH , biphenyl, dimethyl sulfone (DMSO_2), and 3-(trimethylsilyl)-1-propanesulfonic acid sodium salt (DSS-Na) were obtained from Sigma-Aldrich Ltd., Hong Kong. Thirty-seven percent DCl was purchased from Armar Chemicals. d_6 -Dimethyl sulfoxide (DMSO) and deuterium oxide (D_2O) were obtained from Cambridge Isotope Laboratories, Inc. Levulinic acid, $(S)\text{-}[(\text{RuCl}(\text{SEGP}(\text{HOS})))_2(\mu\text{-Cl})_3][\text{NH}_2\text{Me}_2]$, and $(S)\text{-GVL}$ were purchased from Sigma-Aldrich Ltd., Budapest, Hungary.


Dry GVL was prepared by stirring 100 mL GVL with about 5 g P_2O_5 for 3 days at room temperature followed by vacuum distillation at $65\text{ }^\circ\text{C}$ and 1.5 Hgmm.

The NMR experiments were carried out using Bruker AV III 300 or 400 or 600 MHz instruments. For quantitative ^1H experiments, yields were calculated using an internal standard such as biphenyl, DSS-Na and DMSO_2 with pulse delay larger than 10 s. Between NMR experiments or repeated heating cycles in an oil bath, samples were stored at $4\text{ }^\circ\text{C}$ unless otherwise specified.

The enantiomeric excess (*ee*) was determined by using a Finnigan trace gas chromatograph ultra (Thermo Electron Corporation) equipped with a HP-CHIRAL-20B capillary column ($30\text{ m} \times 0.25\text{ mm} \times 0.25\text{ }\mu\text{m}$) with H_2 carrier gas. For the analysis, 10 μL of the reaction mixture was added to 1 mL methanol.

Stability of GVL at $150\text{ }^\circ\text{C}$

Dry GVL (8.60 mL) was placed in a 15-mL pressure glass tube and heated at $150\text{ }^\circ\text{C}$ under air in an oil bath. Liquid samples (250 μL) were taken at different times and transferred to NMR tubes. After the addition of d_6 -DMSO (250 μL), the mixtures were analyzed by NMR.



Scheme 4 The reaction of GVL with aqueous bases

Stability of GVL in the presence of water at $150\text{ }^\circ\text{C}$

A solution of GVL (10.56 mL) and H_2O (1.5 mL) was placed in a 15-mL pressure glass tube and heated at $150\text{ }^\circ\text{C}$ under air in an oil bath. Liquid samples (250 μL) were taken at different times and transferred to NMR tubes. After the addition of a solution of 5 mg (0.023 mmol) 3-(trimethylsilyl)-1-propanesulfonic acid sodium salt (DSS-Na) in 250 μL d_6 -DMSO, the mixtures were analyzed by NMR.

Stability of GVL under aqueous acidic conditions at different acid concentrations

In 5 mm NMR tubes, GVL (1 mL, 10.5 mmol) was mixed with different concentrations of HCl (0.1 to 12 mol/L) or H_2SO_4 (0.1 to 14 mol/L) in amounts that the water contents of the mixtures were constant (0.15 g, 8.3 mmol). The final acid concentrations of the HCl and H_2SO_4 in solutions ranged from 0.02 to 2.0 mol/L and 0.01 to 4.05 mol/L, respectively. The NMR tubes were closed and taken for ^1H -NMR measurement immediately. Then, all tubes were stored at $23\text{ }^\circ\text{C}$ and taken for ^1H -NMR measurement in every week.

Stability of GVL in the presence of 25% NH_4OH

A solution of 2.16 mL GVL (22 mmol) and 0.79 mL 25% NH_4OH (12 mmol NH_3) was stirred at room temperature for 15 h. Liquid samples (200 μL) were then taken and transferred to a NMR tube. After the addition of D_2O (300 μL), the mixture was analyzed by NMR. ^1H NMR of the reaction mixture showed peaks for H_2O , GVL (70%), 4-hydroxyvaleric amide (28%), and 4-hydroxyvaleric acid (2%) (Fig. S4).

A solution of 10.9 mL of GVL (109 mmol) and 54 mL 25% NH_4OH (820 mmol) was stirred at room temperature for 15 h. Liquid samples (200 μL) were then taken and transferred to a NMR tube. After the addition of D_2O (300 μL), the mixture was analyzed by NMR. ^1H NMR of the reaction mixture revealed the presence of 99.9% 4-hydroxyvaleric amide and 0.1% of GVL. The mixture was then distilled under vacuum (3.75 mmHg) at $69\text{ }^\circ\text{C}$ resulting in 8.36 g distillate containing 99.7% of GVL and 0.3% 4-hydroxyvaleric amide.

Synthesis of $(S)\text{-GVL}$

The complete hydrogenation of 9.8 mmol levulinic acid was performed in the presence of 0.004 mmol of $(S)\text{-}[(\text{RuCl}(\text{SEGP}(\text{HOS})))_2(\mu\text{-Cl})_3][\text{NH}_2\text{Me}_2]$ in 1.46 mL methanol under 60 bar H_2 at $140\text{ }^\circ\text{C}$ for 20 h [37]. $(S)\text{-GVL}$ was separated by vacuum distillation ($78\text{--}80\text{ }^\circ\text{C}$, 2 mmHg) and isolated as a colorless liquid (560 mg). Yield: 59.6%, *ee*: 78%.

Stability of (S)-GVL in the presence of water, aqueous HCl, and aqueous NaOH


A mixture of 0.3 μL (S)-GVL (ee = 78%) and 30 μL of an aqueous solution with carefully adjusted pH (between 1 and 13 using aqueous HCl or NaOH) was stirred at different temperatures (25, 50, 75, and 100 $^{\circ}\text{C}$). After 24 h, 1 mL of EtOAc was added to the reaction mixture to form two phases. The upper organic phase was analyzed by chiral GC to establish the enantiomeric excess of the extracted (S)-GVL (Table S3).

Results and discussions

Dry GVL is stable at room temperature under air for a long time and remained stable at 150 $^{\circ}\text{C}$ for several days (Fig. S1). At much higher temperatures (>600 $^{\circ}\text{C}$), GVL can undergo a ring-opening reaction to form the isomers of pentenoic acid followed by the formation of butenes and CO_2 and other decomposition products [13].

Although GVL does not react with water under neutral conditions at room temperature for 3 months and at 60 $^{\circ}\text{C}$ for 28 days [10], GVL and H_2O reaches an equilibrium with 4-HVA (Scheme 2) after several days at 150 $^{\circ}\text{C}$ (Fig. S2). The rather slow approach to the equilibrium is probably due to the combination of the high activation energy of the ring-opening reaction and the very low acidity of water [38]. Thus, when GVL is used as a solvent in chemical reactions in the presence of water (which is either in the reactants or formed during the reactions) at or above 100 $^{\circ}\text{C}$, and the presence of 4-HVA has to be considered as a co-solvent or co-reagent.

The acid catalyzed hydrolysis of cyclic esters is well-known [39]. We have therefore investigated the ring-opening reaction with water in the presence of hydrochloric and sulfuric acids at room temperature. The acid concentrations of HCl and H_2SO_4 were varied between 0.02–2.0 mol/L and 0.01–4.05 mol/L, respectively; meanwhile, GVL and water were used at commensurable quantities. In situ, NMR has revealed that at very high acid concentrations, only relatively small amount of 4-HVA is present in the equilibrium (Table S1). The presence of 4-HVA was confirmed by in situ COSY measurements (Fig. S3). This experimental observation can be initially rationalized by the following three equilibria competing for water:



Since HCl and H_2SO_4 are both strong acids in presence of water, K_{a1} were set as 10^{10} and 10^8 for both. It should be noted that $K_{a2} = 0.0575$ was used in the case of sulfuric acid indicating that the dissociation of proton from HSO_4^- is not




Fig. 1 Initial sulfuric acid concentration dependence of [4-HVA]. ■: experimental —: calculated

complete at large sulfuric acid concentrations. Having the equilibrium concentration of 4-HVA, K_{ring} was obtained by nonlinear parameter estimation performed by ZiTa [40], a comprehensive program package used for determining the kinetic parameters of a given model. During the course of the calculations, the relative average deviation is minimized between the measured and calculated 4-HVA concentrations by optimizing the value of K_{ring} . In the case of using the data obtained in the presence of HCl or H_2SO_4 , the calculated K_{ring} was $(5.85 \pm 0.60) \times 10^{-3}$ and $(4.50 \pm 0.35) \times 10^{-3}$, respectively. The average deviation between the measured and calculated 4-HVA concentrations was 0.04 mol/L for HCl and 0.0374 mol/L for H_2SO_4 . Since the highest measured 4-HVA concentration exceeded 0.4 mol/L, it estimates a more than 10% error of the concentration determination. This difference and the fact that the equilibrium constant of the ring-






Fig. 2 Initial hydrochloric acid concentration dependence of [4-HVA]. ●: experimental —: calculated

Scheme 5. Reversible ring opening of (*S*)-GVL at different pH

opening reaction differs significantly, prompted us to improve the kinetic model. We have added the following two more equilibria into the previous system:



During the course of calculation $K_3 = K_w/[\text{H}_2\text{O}]^2 = 10^{-14}/55.55^2 = 3.24 \times 10^{-18}$ was fixed (at room temperature the ionization constant of water is approximately 10^{-14}) and the value of K_4 was fitted along with that of K_{ring} . As it was done before, the fitting procedure was performed first separately for different acids. Surprisingly, $K_{\text{ring}} = (5.62 \pm 0.20) \times 10^{-3} \text{ M}^{-1}$ and $K_4 = (4.00 \pm 1.10) \times 10^{-4}$ were obtained if hydrochloric acid was used as a catalyst, while $K_{\text{ring}} = (5.38 \pm 0.15) \times 10^{-3}$ and $K_4 = (3.96 \pm 0.85) \times 10^{-4}$ were found in the case of using sulfuric acid as a catalyst. Thus, in the final calculation procedure, we used all the experimental data independently of the nature of the acid applied to obtain K_{ring} and K_4 . These results suggest that indeed the ring-opening reaction is subject to general acid catalysis, so our final calculation procedure was executed using all experiments simultaneously. As a result of this calculation, we concluded that the value of $(5.57 \pm 0.18) \times 10^{-3} \text{ M}^{-1}$ for the equilibrium constant of the ring-opening reaction of GVL at room temperature, meanwhile $K_4 = (3.5 \pm 0.1) \times 10^{-4}$. Figures 1 and 2 show an acceptable agreement between the measured and calculated data and provides a strong support of the model developed.

Since the reaction of GVL with water leads to an equilibrium with 4-HVA, we have also studied its reaction with ammonium hydroxide [23] to see if the formation of 4-hydroxyvaleric amide is reversible. ^1H NMR of the mixture of 21.6 mmol GVL and 12.0 mmol ammonia (using a 25% aqueous NH_4SO_4 solution) has shown the presence of 70% GVL, 28% 4-hydroxyvaleric amide, and 2% 4-HVA after stirring at room temperature for 15 h (Fig. S4). By increasing the ammonia/GVL ratio to 1.2, the concentration of GVL decreased to about 60% indicating that excess amount of ammonia was necessary to drive the reaction to “completion”. Indeed, less than 1% GVL was present in the reaction mixture when the ammonia/GVL ratio was increased to 7.7 (Fig. S5). We have also noted that by heating the reaction mixture under vacuum, most of the GVL could be regenerated, as expected [41].

Enantiomerically pure GVL could be an excellent chiral building block in the synthesis of biologically active compounds and stability of its stereogenic center is crucial for this

purpose. It was already demonstrated by the use of ^{18}O -isotope labeling technique that the reversible ring-opening reaction of GVL in the presence of 0.3 mol/L HCl has no effect on the stability of the chiral center [37]. The ring opening of the chiral GVL resulted in the formation of chiral 4-HVA, which then can subsequently be used to introduce the chiral center under either acidic or alkaline conditions. Consequently, we have investigated the stability of the chiral center between pH =1–13 and 25–100 °C. The chiral center of (*S*)-GVL (ee = 78%) remained unchanged when the aqueous solution of (*S*)-GVL was stirred at 25, 50, 75, and 100 °C for 24 h, as expected. Similar results were observed in the presence of different amount of HCl (pH =1–6) and NaOH (pH =8–13) at 25–100 °C. After the extraction of the reaction mixture with EtOAc, the chiral GC analysis of the organic phase established that the enantiomeric excess of the extracted (*S*)-GVL was $77 \pm 1.75\%$ (Table S1). While the formation of (*S*)-4-HVA was observed at acidic conditions, the (*S*) sodium salt of 4-HVA was formed at basic conditions (Scheme 5).

Conclusions

We have now shown that dry GVL is stable for several weeks at 150 °C and its thermal decomposition only proceed in the presence of appropriate catalysts. Since GVL does not react with water up to 60 °C for several weeks, it could be used as a green solvent at mild conditions. At higher temperatures, GVL reacts with water to form 4-HVA and the equilibrium can be reached in a few days at 100 °C. The establishment of the equilibrium should be even faster at higher temperatures, though the equilibrium should be shifted back towards HVA and water. Aqueous solutions of acids or bases catalyze the ring opening of GVL even at room temperature, which lead to the establishment of an equilibrium between GVL, water and 4-HVA. Although the 4-HVA concentration would be below 4 mol% in the presence of acids, it could be higher than the concentration of a reagent or a catalyst precursor, not to mention a catalytically active species. The latter could be especially worrisome as 4-HVA could be an excellent bi- or even a tridentate ligand for transition metals.

Acknowledgements Some of this work was funded by the Innovation and Technology Support Program of the Innovation and Technology

Fund of the Government of the Hong Kong SAR (ITS/079/13). Any opinions, findings, conclusions or recommendations expressed in this material (or by members of the project team) do not reflect the views of the Government of the Hong Kong SAR, the Innovation and Technology Commission or the Panel of Assessors for the Innovation and Technology Support Program of the Innovation and Technology Fund. We also thank the Environment and Conservation Fund (ECF/31/2014) for partial financial support. L. T. Mika is grateful for the support of the National Research, Development and Innovation Office—NKFIH, Budapest, Hungary (PD116559), and the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, Budapest, Hungary. A. K. Horváth is grateful for the support of the National Research, Development and Innovation Office, NKFIH, Hungary (K116591).

References

- Reichardt C (2007) Solvents and solvent effects: an introduction. *Org Process Res Dev* 11:105–115
- Horváth IT (2008) Solvents from Nature. *Green Chem* 10:1024–1028
- Wypych G (2001) Handbook of solvents. ChemTec Publishing, Toronto
- Anastas PT, Warner JC (1998) Green chemistry: theory and practice. Oxford University Press, Oxford
- Kerton FM (2009) Alternative solvents for green chemistry. Royal Society of Chemistry, Cambridge
- Vigneron JP, Bloy V (1980) Preparation d'alkyl-4 gamma-lactones optiquement actives. *Tetrahedron Lett* 21:1735–1738
- Mehdi H, Horváth IT, Bodor A (2004) Dehydration and hydrogenation of carbohydrates with aqueous biphasic catalysts (CELL 95). 227th ACS National Meeting, Anaheim, CA
- Horváth IT (2006) Gamma-valerolactone: a sustainable liquid for energy and carbon based chemicals. The 10th Annual Green Chemistry and Engineering Conference, Washington DC
- Mehdi H, Tuba R, Mika LT, Bodor A, Torkos K, Horváth IT (2006) Catalytic conversion of carbohydrates to oxygenates. Renewable resources and renewable energy: a global challenge, Graziani M, Fornasiero P. (Eds), CRC Press: Boca Raton, FL, Chapter 4, 55–60
- Horváth IT, Mehdi H, Fábos V, Boda L, Mika LT (2008) γ -valerolactone – a sustainable liquid for energy and carbon-based chemicals. *Green Chem* 10:238–242
- Mehdi H, Fábos V, Tuba R, Bodor A, Mika LT, Horváth IT (2008) Integration of homogeneous and heterogeneous catalytic processes for a multi-step conversion of biomass: from sucrose to levulinic acid, gamma-valerolactone, 1,4-pentanediol, 2-methyl-tetrahydrofuran, and alkanes. *Top Catal* 48:49–54
- Fábos V, Lui MY, Mui YF, Wong YY, Mika LT, Qi L, Cséfalvay E, Kovács V, Szűcs T, Horváth IT (2015) The use of gamma-valerolactone as an illuminating liquid and lighter fluid. *ACS Sust Chem Eng* 3:1899–1904
- De Bruycker R, Carstensen H-H, Simmie JM, van Geem KM, Marin GB (2015) Experimental and computational study of the initial decomposition of gamma-valerolactone. *Proc Combust Inst* 35:515–523
- Havasi D, Mizsey P, Mika LT (2016a) Vapor–liquid equilibrium study of the gamma–valerolactone–water binary system. *J Chem Eng Data* 61:1502–1508
- Havasi D, Pátzay G, Kolarovszki Z, Mika LT (2016b) Isobaric vapor–liquid equilibria for binary mixtures of γ -valerolactone + methanol, ethanol, and 2-propanol. *J Chem Eng Data* 61:3326–3333
- Fábos V, Koczó G, Mehdi H, Boda L, Horváth IT (2009) Bio-oxygenates and the peroxide number: a safety issue. *Energy and Env Sci* 2:767–769
- Qi L, Horváth IT (2012) Catalytic conversion of fructose to γ -valerolactone in γ -valerolactone. *ACS Catal* 2:2247–2249
- Bond JQ, Alonso DM, Wang D, West RM, Dumesic JA (2010a) Integrated catalytic conversion of γ -valerolactone to liquid alkenes for transportation fuels. *Science* 327:1110–1114
- Serrano-Ruiz JC, Wang D, Dumesic JA (2010) Catalytic upgrading of levulinic acid to 5-nonanone. *Green Chem* 12:574–577
- Bond JQ, Alonso DM, West RM, Dumesic JA (2010b) Valerolactone ring-opening and decarboxylation over $\text{SiO}_2/\text{Al}_2\text{O}_3$ in the presence of water. *Langmuir* 26:16291–16298
- Olah GA, Ku AT (1970) Stable carbonium ions. CVIII. Protonated lactones and their cleavage reactions in fluorosulfonic acid-antimony pentafluoride solutions. *J Org Chem* 35:3916–3922
- Umland J, Witkowski S (1957) Reaction of silver 4-hydroxyvalerate with bromine. *J Org Chem* 22:345
- Chalid M, Heeres HJ, Broekhuis AA (2012) Ring-opening of γ -valerolactone with amino compounds. *J Appl Polym Sci* 123:3556–3564
- Fegyverneki D, Orha L, Láng G, Horváth IT (2010) Gamma-valerolactone based solvents. *Tetrahedron* 66:1078–1081
- Qi L, Mui YF, Lo SW, Lui MY, Akien GR, Horváth IT (2014) Catalytic conversion of fructose, glucose, and sucrose to 5-(hydroxymethyl) furfural and levulinic and formic acids in γ -valerolactone as a green solvent. *ACS Sustain Chem Eng* 4:1470–1477
- Tukacs JM, Király D, Strádi A, Novodárszki G, Eke Z, Dibó G, Kégl T, Mika LT (2012) Efficient catalytic hydrogenation of levulinic acid: a key step in biomass conversion. *Green Chem* 14:2057–2065
- Ismalaj E, Strappaveccia G, Ballerini E, Elisei F, Piermatti O, Gelman D, Vaccaro L (2014) γ -Valerolactone as a renewable dipolar aprotic solvent deriving from biomass degradation for the Hiyama reaction. *ACS Sustain Chem Eng* 2:2461–2464
- Strappaveccia G, Ismalaj E, Petrucci C, Lanari D, Marrocchi A, Drees M, Fachetti A, Vaccaro L (2015a) A biomass-derived safe medium to replace toxic dipolar solvents and access cleaner Heck coupling reactions. *Green Chem* 17:365–372
- Strappaveccia G, Luciani L, Bartollini E, Marrocchi A, Pizzo F, Vaccaro AL (2015b) γ -Valerolactone as an alternative biomass-derived medium for the Sonogashira reaction. *Green Chem* 17:1071–1076
- Rasina D, Kahler-Quesada A, Ziarelli S, Warratz S, Cao H, Santoro S, Ackermann L, Vaccaro L (2016) Heterogeneous palladium-catalysed Catellani reaction in biomass-derived γ -valerolactone. *Green Chem* 18:5025–5030
- Horváth IT (2014) Green or sustainable chemistry or both? *Chimica Oggi - Chem Today* 32:76–79
- Tian X, Yang F, Rasina D, Bauer M, Warratz S, Ferlin F, Vaccaro L, Ackermann L (2016) C–H Arylations of 1,2,3-triazoles by reusable heterogeneous palladium catalysts in biomass-derived γ -valerolactone. *Chem Commun* 52:9777–9780
- Pongrácz P, Kollár L, Mika LT (2016) A step towards hydroformylation under sustainable conditions: platinum-catalysed enantioselective hydroformylation of styrene in gamma-valerolactone. *Green Chem* 18:842–847
- Marosvölgyi-Haskó D, Lengyle B, Tukacs JM, Kollár L, Mika LT (2016) Application of γ -valerolactone as an alternative biomass-based medium for aminocarbonylation reactions. *Chem Plus Chem*. doi:10.1002/cplu.201600389
- Gu Y, Jerome F (2013) Bio-based solvents: an emerging generation of fluids for the design of eco-efficient processes in catalysis and organic chemistry. *Chem Soc Rev* 42:9550–9570

36. Soh L, Eckelman MJ (2016) Green solvents in biomass processing. *ACS Sustain Chem Eng* 4:5821–5837
37. Tukacs JM, Fridric B, Dibó G, Székely E, Mika LT (2015) Direct asymmetric reduction of levulinic acid to gamma-valerolactone: synthesis of a chiral platform molecule. *Green Chem* 17:5189–5195
38. http://www1.lsbu.ac.uk/water/water_dissociation.html (accessed on 26/10/2016)
39. Long FA, Paul MA (1957) Application of the H_0 acidity function to kinetics and mechanisms of acid catalysis. *Chem Rev* 57:935–1010
40. Peintler G, (1989–1998) ZiTa, version 5.0; a comprehensive program package for fitting parameters of chemical reaction mechanism; Attila József University: Szeged, Hungary
41. Bruice TC, Marquardt FH (1962) Hydroxyl group catalysis. IV. The mechanism of intramolecular participation of the aliphatic hydroxyl group in amide hydrolysis. *J Amer Chem Soc* 84:365–370