



CECE 2017
14th International Interdisciplinary Meeting on Bioanalysis
Veszprém, Hungary, 8-11 October, 2017

List of poster presentations

- P1. IDENTIFICATION OF O-GLYCOSYLATED (N-TERMINAL) PRO-B-TYPE NATRIURETIC PEPTIDE FORMS IN BLOOD PLASMA OF PATIENTS WITH SEVERE HEART FAILURE
Benno Amplatz, Bernhard Halfinger, Angelika Hammerer-Lercher, Bettina Sarg, Leopold Kremser, and Herbert H. Lindner
- P2. CAPILLARY ISOTACHOPHORESIS DETERMINATION OF TRACE OXIDIZED GLUTATHIONE IN BLOOD
Róbert Bodor, Andrea Nečasová, Alena Pechová, and Marián Masár
- P3. GLYCOSIMILARITY INDEX FOR BIOTHERAPEUTICS
Andras Guttman, Beata Borza, Marton Szigeti, Akos Szekrenyes, Laszlo Hajba
- P4. SEMIAUTOMATION OF LIQUID HANDLING FOR MICROELECTROMEMBRANE EXTRACTION OF COMPLEX SAMPLES
Miloš Dvořák, Andrea Šlampová, And Pavel Kubáň
- P5. DEVELOPMENT OF GRADIENT RP-HPLC METHOD FOR CHARACTERIZATION OF AMINO ACID COMPOSITION OF DEGRADATION PRODUCTS OF HUMIC ACIDS ISOLATED FROM PEAT AND SOIL AFTER PRE-COLUMN DERIVATIZATION
Róbert Góra, Milan Hutta, Erik Beňo, and Natália Bielčíková
- P6. STUDY OF VARIOUS PARAMETERS INFLUENCING CONTENT OF EXHALED BREATH CONDENSATE WITH IMPORTANCE FOR DIAGNOSIS OF GASTROESOPHAGEAL REFLUX DISEASE
Michal Greguš, Pavol Ďurč, Júlia Lačná, František Foret And Petr Kubáň
- P7. DETERMINATION OF THE OLIGOSACCHARIDE COMPOSITION IN WORT SAMPLES BY CAPILLARY ELECTROPHORESIS
Tamás Gábor Szilágyi, Beáta Hegyesné Vecseri, Zsuzsanna Kiss, László Hajba, András Guttman
- P8. MULTISCALE MODELING OF RECTIFYING BIPOLAR NANOPORE
Z. Ható, M. Valiskó, T. Kristóf, D. Gillespie, D. Boda
- P9. CFD MODELLING BASED INVESTIGATION OF THE ELECTROOSMOTIC AND FORCED CONVECTION FLOW IN CAPILLARY ELECTROPHORESIS APPLICATIONS
Gábor Járvas, Márton Szigeti, and András Guttman
- P10. SEARCH FOR MULTIPLE MYELOMA GLYCOBIOMARKERS BY CE-LIF
Zsuzsanna Kovács, Anna Farkas, and András Guttman

- P11. BIOBANKING OF PATIENT SAMPLES FOR GENO – GLYCOMIC LUNG DISEASE BIOMARKER STUDIES
Renáta Kun, Eszter Csánky, Miklós Szabó, Zsolt Rónai, Zsuzsanna Kovács, Márton Szigeti, Gábor Járvas, László Hajba, Boglárka Döncző, András Guttman
- P12. SINGLE-CELL ANALYSIS OF ACTIVE CASPASE 3/7 IN APOPTOTIC CELLS
Vojtěch Ledvina and Karel Klepárník
- P13. APPLICATION OF AMINO ACIDS AND PEPTIDES FOR OLIGOSACCHARIDE LABELING
Jana Krenkova, Jan Partyka, and Frantisek Foret
- P14. CAPILLARY ELECTROPHORETIC ANALYSIS OF EXHALED BREATH CONDENSATE IN THE DIAGNOSIS OF GASTROESOPHAGEAL REFLUX DISEASE
Petr Kubán, Pavol Ďurč, Júlia Lačná, Michal Greguš, František Foret, Jiří Dolina, Štefan Konečný, Martina Doubková, Dagmar Kindlová, Eva Pokojová, and Jana Skříčková
- P15. USING NANOPORES AS SENSORS: A COMPUTER SIMULATION STUDY
Eszter Máдай, Mónika Valiskó, and Dezső Boda
- P16. QUANTITATIVE ASSESSMENT AND AUTOMATIC ID OF GLYCANS OF CQA IMPORTANCE IN BIOLOGICS DEVELOPMENT
Andras Guttman, Marton Szigeti, Brigitta Meszaros, Jeff Chapman
- P17. DEVELOPMENT OF AN UHPLC-MS/MS METHOD FOR ANALYSIS OF A NOVEL CARDIOPROTECTIVE AGENT JAS 2 AND ITS METABOLITE
Hana Piskáčková, Petra Reimerová, Jaroslav Roh, Jana Sedláková, and Petra Štěrbová Kovaříková
- P18. *IN VITRO* BIOACTIVATION STUDY OF A NEW ANTIPROLIFERATIVE DRUG SOBUZOXANE IN BIOLOGICAL MATRICES
Petra Reimerová, Anna Jirkovská, Hana Piskáčková, Galina Karabanovich, Jaroslav Roh, and Petra Štěrbová-Kovaříková
- P19. COMBINATION OF PREPARATIVE ISOELECTRIC FOCUSING AND MALDI-TOF MS FOR IDENTIFICATION OF COLORED BACTERIA
Jiří Šalplachta, Marie Horká, Filip Růžička, and Karel Šlais
- P20. DETECTING MONO-PHOSPHORYLATED ISOBARIC PEPTIDES USING CESI-MS
Bettina Sarg, Klaus Faserl, and Herbert H. Lindner
- P21. COLD PLASMA: THE WAY TO IMPROVE THE SALD ICP MS ANALYSIS REPEATIBILITY
Marek Stiborek, Jan Preisler, Masoud Shekargoftar, Viktor Kanický, and Jakub Kelar
- P22. THE POWER OF GU AND THE IMPORTANCE OF TEMPERATURE CONTROL
András Guttman, Márton Szigeti, Jeff Chapman

- P23. DETERMINATION OF HOMOCYSTEINE IN URINE AND SALIVA BY MICROCHIP ELECTROPHORESIS
Peter Troška, Anna Mandžáková, Jasna Hradski, Juraj Ševčík, and Marián Masár
- P24. ANALYSIS AND IDENTIFICATION OF POLYPHENOLIC COMPOUNDS IN GREEN FOODS USING A COMBINATION OF HPLC-ESI-IT-TOF-MS/MS
Pavel Škvára, Júlia Kezmanová, Csilla Mišlanová, Andrea Vojs Staňová
- P25. SILICA MONOLITHIC CAPILLARY COLUMNS FOR HILIC SEPARATIONS
Dana Moravcová and Josef Planeta
- P26. MOLECULAR GLYCOHISTOPATHOLOGY BY CAPILLARY ELECTROPHORESIS
Boglárka Dönczö and András Guttman
- P27. ENHANCED SEPARATION AND CHARACTERIZATION OF DEAMIDATED PEPTIDES WITH CE-MS
Klaus Faser, Bettina Sarg, and Herbert H. Lindner

P15 MULTISCALE MODELING OF RECTIFYING BIPOLAR NANOPORE

ESZTER MÁDAI¹, MÓNICA VALISKÓ¹ AND DEZSŐ BODA^{1,2}

¹ Department of Physical Chemistry, University of Pannonia, Veszprém, Hungary

² Institute of Advanced Studies Kőszeg (iASK), Kőszeg, Hungary

Abstract

For nanopores, there can be made a difference between natural and artificial nanopores, as ion channels and carbon nanotubes or nanopores cauterized into foils. Artificial nanopores have microscopic-length and a nanoscopic radius-size. Ion transport through the nanopore is connected with an important property: the selectivity. Due to this property the nanopore can be used, for instance, as a nanoscopic porous sensor.

The basic concept of our model nanopore sensor is that the presence of an analyte molecule modulates the ionic current flowing through the pore. The analyte molecule (usually a biomolecule) is detected selectively if it is bound to the nanopore. If the binding is a reversible process involving physical forces that are weaker than chemical bonds, the probability that the analyte molecule binds to the nanopore is proportional to its concentration. From calibration curves (current vs. concentration) the analyte concentration can be determined.

In this modeling work, we create simple models the membrane, the pore and the electrolyte. Negative charges are located on the cylindrical pore's wall to make the pore cation selective. For the electrolyte, the Primitive Model of electrolytes is used, which treats the ions as hard spheres with their charges located in the centers, and the solvent as a constant dielectric continuum. The binding sites are modeled with a square-well potential. We applied the Local Equilibrium Monte Carlo (LEMC) method coupled with Nernst-Planck (NP) equation. This method (NP+LEMC) is suitable for study even micromolar concentrations.

We present a modeling study to analyze the device sensitivity (defined as I/I_0 , where I and I_0 are the currents in the presence and absence of the analyte, respectively) to the various model parameters: number, strength, and width of the binding sites; properties of the electrolyte and the analyte. Through the calculated concentration profiles, the molecular mechanisms of the system can be understood.