



ACTA

MINERALOGICA-PETROGRAPHICA

FIELD GUIDE SERIES

Volume 32

Szeged, 2017



15th Meeting of the
Central European Tectonic
Studies Group

László FODOR, Gábor HÉJA, Szilvia KÖVÉR, Gábor CSILLAG, Ádám Lajos CSICSEK:
Cretaceous deformation of the south-eastern Transdanubian Range Unit, and the effect
of inherited Triassic–Jurassic normal faults

Tivadar M. TÓTH, Ágnes FISER-NAGY, Félix, SCHUBERT:
Petrology and structural evolution of the metamorphic basement of the Pannonian
Basin

CETEG 2017 ABSTRACT VOLUME, PRE- AND POST-CONFERENCE FIELD TRIP
GUIDE



Cestmir Tomek: East Moesia continental subduction, slab extension, and the origin of intermediate earthquakes beneath Vrancea Mountains of Romania (**keynote talk**) (10h50-11h20)

István Kovács, László Lenkey, David H. Green, Tamás Fancsik, György Falus, János Kiss, László Orosz, Zsuzsanna Viktor, J. Angyal: The role of amphibole in the dynamics and rheology of the lithosphere–asthenosphere system (11:20-11:40)

László Aradi, Károly Hidas, István János Kovács, Csaba Szabó: Deformation and fluid-enhanced annealing in subcontinental lithospheric mantle beneath the Pannonian Basin (Styrian Basin, Eastern Austria) (11:40-12:00)

Session +

Ondrej Lexa: Presentation of softwares for structural geological purposes (free of use) (12:00-12:30)

12:30-13:50: Lunch break

Session 3: Connection between deformation and sedimentation (5) (13:50-15:30)

Chairman: Rastislav Vojtko

Krisztina Sebe: From post-rift phase to basin inversion – late-stage tectonic evolution of the Mecsek–Villány area, SW Hungary (13:50-14:10)

Tamás Csibri, Samuel Rybár, Michal Kováč: Depositional environment of the Lower and Middle Miocene conglomerates in key wells of the northern Blatné depression (Danube Basin) (14:10-14:30)

Michal Krobicki, T. Gagulski & G. Gorczyca: The early Late Cretaceous synsedimentary tectonic events in the PeriTethys region (Miechów Synclinorium and Silesian-Kraków Homocline, southern Poland) (14:30-14:50)

Nevenka Djerić, Špela Goričan, Nataša Gerzina Spajić, Renata Jach, Daniela Rehakova: The use of biostratigraphic tools in dating the remnants of the Alpine Tethys (Danubian Nappes, eastern Serbia) (14:50-15:10)

Éva Oravecz, László Fodor, Szilvia Kövér: Significance of Jurassic early deformation structures in the SW-Bükk Mts. (15:10-15:30)

15:30-15:50 Coffee break and posters

Session 4: Paleomagnetism, fractures (6) (15:50-17:50)

Chairman: Prokop Zavada

František Hrouda, Josef Ježek, Stuart Gilder, Michael Wack: Different sensitivities of paramagnetic and ferromagnetic minerals to ductile deformation in Intra-Carpathian Palaeogene of the Skorušinské vrchy Hills (15:50-16:10)

Emő Márton, Jozef Madzin, Jana Bučová, Jacek Grabowski, Dušan Plašienka, Roman Aubrecht: New paleomagnetic results from Permian and Triassic rocks from the Hronic Nappe (Nízke Tatry) and Paleogene sediments from the Central Carpathian Paleogene Basin, Western Carpathians (16:10-16:30)

Kristina Kolářová, Jan Černý, Rostislav Melichar: Magnetic and structural studies of Křivoklát-Rokycany volcanic complex, Barrandian, Czechia (16:30-16:50)

Nováková Lucie, Schnabl Petr, Štěpančíková Petra, Křížková Martina: Microtectonic studies of the faults (16:50-17:10)

Jan Barmuta, Piotr Hadro, Krzysztof Dzwiniel: Determination of fracture distribution based on seismic attributes and geomechanical modelling (17:10-17:30)

Jan Juráček, Rostislav Melichar: Post-Variscan tectonics of central part of the Bohemian Cretaceous Basin (Czech Republic) (17:30-17h50)

17:50-19:00 Poster session

19:00 Wine tasting and dinner

mineralization products of (1) and (2). In (3) the orogen parallel hot line is located in the South-Veporic zone, producing besides the uplifting of this zone, also the penetration of ore-bearing fluids into appropriate extensional fractures (C) in the contact zone between Veporicum and Gemericum, as well as in minor amount into Gemericum. The existence of two generations of pre-Neogene mineralization within the Western Carpathians (e.g. in Gemericum it was a dominant Variscan and minor Alpine mineralization) was a reason of misunderstanding and many decades lasting scientific disputes among geologists in Slovakia.

Neogene mineralization phase (4) is areally limited to solidified Miocene volcanic bodies (C), being either effusive (solidified lava flows) or extrusive (disseminations in volcanoclastic and pyroclastic sediments), as well as to their close vicinity (usually in the form of disseminated and stockwork mineralization). The origin of magma (A) is generally explained by the heat flow (B) related to asthenosphere upwelling due to subduction slab dipping from the north under Inner Western Carpathians.

The repeated metallogenetic phases in the Western Carpathians owe to existence of at least three orogenic (Wilson) cycles operating in the mobile zone between the Gondwana and Laurasia (Laurussia) in the territory of recent Slovakia. These metallogenetic phases have produced the mineral wealth of the country, including numerous deposits of public importance.

Acknowledgement. The contribution originated of the basis of numerous projects funded by the Ministry of Environment of the Slovak Republic, especially the projects Critical Raw Materials and Investigation of geological setting and compilation of geological maps in problematic areas in the Slovak Republic. It is also a contribution to EC Horizon 2020 project MINATURA 2020, as well as research grants VEGA-1/0079/15 and APVV-15-0050.

Microtectonic studies of the faults

Lucie Nováková¹, Petr Schnabl², Petra Štěpančíková¹, Martina Křížková¹

¹ Institute of Rock Structure and Mechanics of the CAS, v.v.i., V Holešovičkách 41, 182 09, Prague 8, Czech Republic, lucie.novakova@irms.cas.cz

² Institute of Geology of the CAS, v.v.i., Rozvojová 269, 165 00 Prague 6 - Lysolaje, Czech Republic

Fault zones were studied in microscopic scale to compare microfractures and grains within a fault plane with the striations visible on the fault surface in the field. Different lithologies were chosen in different geological conditions. Within the Central Europe, we have tested samples of the sediments, crystalline, and metamorphic rocks. We have established the methodology for studying and processing the samples as follows:

1. Sampling of oriented samples in the field
2. Preparation of the polished samples
3. Photography of a complete sample using an optical microscope (Fig. 1a)
4. Detail photography net of sample and picture composition (Fig. 1b)
5. Highlighting microfractures and grains in the composite picture manually (Fig. 1c) and semi- automatically (Fig. 1d)
6. Vectorization and topology imaging (Fig. 1e)
7. Automatic and manual reading of bearing

8. Visualization of the microfractures and grains (Fig. 1f)

The preferential orientations of microfractures and grains were compared with the striations on the fault surface. Anisotropy of magnetic susceptibility (AMS) was measured on five samples in order to correlate the microstructure and grain orientation of large mostly diamagnetic minerals with the orientation of paramagnetic and ferromagnetic minerals. The results showed two different behaviours. The more common one is the longest AMS axis k_1 aligned with striation and the orientation of the grains (gneiss and phyllite). The second behaviour, which was found only within the red quartzite, shows the shortest AMS axis k_3 parallel to the lineation.




Fig. 1 Sample processing step by step.

This work was carried out thanks to the support of the long-term conceptual development research plans RVO: 67985891 and RVO: 67985831.

Significance of Jurassic early deformation structures in the SW-Bükk Mts.

Éva Oravecz¹, László Fodor², Szilvia Kövér²

¹ Eötvös Loránd University, 1117 Budapest, Pázmány Péter sétány 1/c, Hungary, orav.eva@gmail.com

² MTA-ELTE Geological, Geophysical and Space Science Research Group, Hungarian Academy of Sciences at the Eötvös Loránd University, 1117 Budapest, Pázmány Péter sétány 1/c, Hungary

Field work in the quarry of Bükkzsérc and at the Patkó Cliffs (SW of Bükk Mts.) revealed several types of early, soft-sediment deformation structures in the Jurassic Bükkzsérc Limestone which was deposited in a subduction-related basin of the Neotethys Ocean and now appears as olistoliths in the accretionary Mónosbél Complex. These structures include small syn-sedimentary normal faults, ductile shear surfaces and slump folds, both of which were not recognized by previous structural geological and sedimentological studies (e.g. Németh 2007), although these structures have direct implications for the poorly known early deformation history of the Mónosbél Complex and possibly of the Bükk para-autochthonous unit.

The aim of our study was to estimate the Jurassic paleo-stress field for the Mónosbél Complex of the Bükk Mts. by measuring syn-sedimentary faults, and to estimate the paleoslope direction from the slump fold geometries.

In case of the normal faults, the key observations in our interpretation were that the faults are characterized by rounded, curved shapes, and that they don't have discrete fault planes. This can be explained by syn-sedimentary deformation in case of which the deformation in soft sediments is limited, while the diagenetic processes may have further effects on fault geometries as well.

Data were rotated back to its paleoposition by the average paleomagnetic data attributed for the Bükk Mts. (e.g. Márton & Márton 1996). Based on the back-rotated small-scale deformation structures we estimated a NE-SW extension as the Jurassic paleostress field, while the results of the slump fold analysis showed a SW paleoslope direction. Although these extensional directions match the previous broad estimations, we need to take into account that the olistolith itself could have had further rotation, so in order to finalize our estimations, additional measurements in other olistoliths and direct paleomagnetic data from the Bükkzsérc olistolith are still needed.

The research was supported by the research found NKFIH OTKA 113013 and the ÚNKP-16-1 New National Excellence Program of the Ministry of Human Capacities.

References

- Márton, E., Márton, P., 1996: Large scale rotations in North Hungary during the Neogene as indicated by paleomagnetic data — The Geological Society, London, Special Publications 105, p. 153-173.
Németh, N., 2007: Structural features of the quarry at the Patkó Cliffs, Bükkzsérc and its surroundings — *Földtani Közlemények* 137/4, p. 473-486.

Characteristic microstructures and deformation of the Dorozsma marble, Tisza Mega-unit, South Hungary

Nikoletta Papp, Andrea Varga, Béla Raucsik, Előd Mészáros

University of Szeged, Department of Mineralogy, Geochemistry and Petrology, Vulcano Research Group, H-6722, Szeged, Egyetem u. 2-6, Hungary, pappnicki@gmail.com

The basement of the Tisza Mega-unit, Hungary, comprises three Alpine zones: the Mecsek, Villány and Békés-Codru Zones. Within the Békés-Codru Zone, the metamorphic basement block near the village of Dorozsma can be subdivided into three tectonic units: the upper unit consists mainly of mica schist and gneiss with sporadic occurrence of metapegmatite, while the lower unit consists dominantly of amphibole-biotite gneiss and amphibolite. Between these tectonic blocks a few-meter-thick marble unit was identified (M. Tóth 2008).

Petrographic examination of marble core samples and archive thin sections was carried out from the boreholes of Dorozsma-4, Dorozsma-7 and Dorozsma-54. The studied rocks are greyish white, fine grained marble samples showing brittle-deformation features. Under a microscope the samples are fine grained marbles with heteroblastic texture. The carbonate crystal boundaries are dominantly sutured, embayed and rarely curved. All the morphological types of deformation twins appear in the samples, but dominantly twin types I, II and IV are present. The characteristic features of the samples are small dissolution cavities filled with fine crystalline carbonate, blocky quartz and, in some cases, with saddle dolomite. In the samples from the borehole Dorozsma-4 small inclusion free carbonate grains substitute the deformation twins of the large carbonate crystals. In addition, small carbonate grains and

Mg-chlorite flakes can be found among the large carbonate crystals. Based on the observed microtextural features the fine grained carbonatic edge could be the result of a low temperature dynamic recrystallization (Liu et al. 2002). Characteristic microstructures of the samples from well Dorozsma-7 are large sigmoid carbonate clasts in very fine grained, micritic matrix. The deformation twins of the carbonate clasts can be easily identified but show the signs of recrystallization. These samples beside the carbonate clasts also contain irregular shaped polycrystalline quartz grains with dissolved edges, undulose extinction and signs of dynamic recrystallization. The microstructures presented above provide an evidence for ductile deformation with a shearing component. The undirected fabric of the micritic matrix and the sigmoid carbonate clasts suggest that the main mechanism of the deformation was probably the grain boundary sliding during hydrolytic weakening (Passchier & Trouw 2005).

Consequently, the aforementioned microstructures of the studied marble samples from the boreholes Dorozsma-4 and Dorozsma-7 could be the results of hydrolytic weakening effect of hydrothermal fluids during deformation (Liu et al. 2002).

This research has been supported by the National Research, Development and Innovation Office (NKFIH K 108375).

References

- Liu, J., Walter, J., M. & Weber, K. 2002: Fluid-enhanced low-temperature plasticity of calcite marble: Microstructures and mechanisms. — Geological Society of America.
M. Tóth, T. 2008: Repedezett, metamorf fluidumtárolók az Alföld aljzatában. — DSc dissertation.
Passchier, C.S. & Trouw, R.A.J. 2005: *Microtectonics*. — Springer Verlag.

Pre-Variscan rift-related structure and metamorphism in the Domažlice Crystalline Complex, (the Teplá-Barrandian Domain, Bohemian Massif)

Vít Peřestý¹, Ondřej Lexa¹, Pavla Štípská², Petr Jeřábek¹, Martin Racek¹

¹ Institute of Petrology and Structural Geology, Charles University in Prague, Czech Republic (perev7an@natur.cuni.cz)

² Centre for Lithospheric Research, Czech Geological Survey, Prague, Czech Republic

The best preserved record of the pre-Variscan regional metamorphism in the Teplá-Barrandian Domain is traditionally ascribed to the Domažlice Crystalline Complex. In order to better constrain the nature and timing of this event, we have investigated metasedimentary rocks by structural analysis, in-situ U-Pb LASS ICP-MS monazite and xenotime dating and P-T modelling. The rocks were sampled across the biotite, garnet, staurolite-andalusite and sillimanite zones.

The structural analysis revealed dominantly gently ESE inclined fabric S1 which is modified by open large-scale folds with moderately to steeply NW-dipping axial planes S2. Textural relationships suggest that garnet1-, andalusite-, staurolite- and sillimanite-bearing mineral assemblages are post-dating the S1 fabric and pre-dating D2. The mineral growth related to the D2 event is almost missing. It represents rare younger garnet2 generation or