

during Middle and Late Devonian and were emplaced in the upper crust during Early Carboniferous. Magmas show strong involvement of the Cambrian lower crustal material concerning also the eclogites, and, based on

variable relative enrichment of mobile elements (LREEs and LILEs) among the samples, a possible heterogeneous contribution of the subduction component in the system.

REFERENCES

- Collett S., Štípská P., Schulmann K., Peřestý V., Soldner J., Anczkiewicz R., Lexa O. & Kylander-Clark A., 2018. Combined Lu-Hf and Sm-Nd geochronology of the Mariánské Lázně Complex: New constraints on the timing of eclogite- and granulite-facies metamorphism. *Lithos*, 304–307, 74–94.
- Timmermann H., Štědrá V., Gerdes A., Noble S.R., Parrish R.R. & Dörr W., 2004. The problem of dating high-pressure metamorphism: a U–Pb isotope and geochemical study on eclogites and related rocks of the Mariánské Lázně Complex, Czech Republic. *Journal of Petrology*, 45, 1311–1338.
- Žák J., Kratinová Z., Trubač J., Janoušek V., Sláma J. & Mrlina J., 2011. Structure, emplacement, and tectonic setting of Late Devonian granitoid plutons in the Teplá–Barrandian unit, Bohemian Massif. *International Journal of Earth Sciences*, 100, 7, 1477–1495.

Previous concepts and new data on the structural and magmatic evolution of the Bükk Mts., NE Hungary:

first step toward the reconsideration of geodynamic models

László Fodor^{1,2}, Szilvia Kövér¹, Gabriella B. Kiss³, Nevenka Đerić⁴, Melinda Fialowski¹, Anette Götz⁵, Zoltán Gulácsi⁶, János Haas¹, Zoltán Kovács^{1,9}, Éva Oravecz¹, Martin Reiser^{7,8}, Benjámín Scherman¹, Ralf Schuster⁷

¹Eötvös Loránd University, Hungarian Academy of Sciences, MTA-ELTE Geological, Geophysical and Space Science Research Group; Budapest, Hungary

²Hungarian Academy of Sciences, MTA-ELTE Volcanological Research Group; Budapest, Hungary

³Eötvös Loránd University, Department of Mineralogy; Budapest, Hungary

⁴University of Belgrade, Department of Paleontology; Belgrade, Serbia

⁵University Portsmouth, School Earth and Environmental Sciences; Portsmouth, UK

⁶Mining and Geological Survey of Hungary; Budapest, Hungary

⁷Geologische Bundesanstalt; Wien, Austria

⁸University of Innsbruck, Department of Geology; Innsbruck, Austria

⁹Eötvös Loránd University, Department of Petrology and Geochemistry, Lithosphere Fluid Research Lab; Budapest, Hungary

The Bükk Mts. are displaced fragments of the Inner Dinaric nappe system (Schmid et al. 2008). Two major concepts described their structure; one suggests four stratigraphic units as continuous, slightly deformed units (Less et al. 2005), while others considered these units as distinct nappes (Balla 1983, Csontos 1999, 2000). In our contribution we list some new, preliminary observations which may challenge both concepts and will hopefully provide a new one.

The Bükk Complex records the disintegration of the Triassic platform and deposition of Middle(?) Jurassic pelagic radiolarite. It is covered by the Lökvolgy distal turbidites, a slate with siltstone to sandstone intercalations. The overlying Mónosbél sequence contains shale, limestone intercalations, olistostromes and radiolarites. The Szarvaskő Complex is a Jurassic basaltic to gabbroic magmatic suite intruded in or flowed onto fine siliciclastics (Balla 1983). Finally, the Darnó Complex represents an *amélange* unit, composed of shales and embedded clasts of Triassic–Jurassic sediments and magmatic rocks (Kovács et al. 2011). All units were slightly metamorphosed and folded, the degree of alteration decreases upward. In the nappe-stack concept, the Mónosbél, Szarvaskő and Darnó Complexes are nappes, partly derived from the Neotethyan ophiolites (Csontos 1999, 2000):

- Newly described syn-diagenetic faults, slump folds can be connected to the passive margin setting of the Mónosbél Complex.
- Most, if not all olistolithes can be originated from the passive Adriatic margin, very few could have been scrapped off the Triassic part of the Neotethyan downgoing slab. Limestones are rich in clasts originated from coeval Bajocian–Bathonian carbonate platform.
- The Darnó Complex contains clasts from the overriding upper plate: blocks of Triassic and Jurassic radiolarites as well as Triassic basalts of advanced rifting origin (with peperitic red limestone) and Jurassic basalts similar to those occurring at Szarvaskő (Kiss et al. 2010, 2016, Kovács et al. 2011). We suggest that this is the only ophiolite-related tectono-sedimentary *mélange* in the Bükk area.
- There is a clear structural boundary at the top of the Lökvolgy Fm. (base Mónosbél Complex); folds are overturned, closed and have regular axial-plane foliation, while folds are open and have layer-parallel foliation, below and above the boundary. In addition, foliation, faults, or boudins form sigmoidal geometry, resembling to S–C foliation/sigma-clasts/faults in duplex; all indicating simple shear. Part of the shearing

with SSE to ESE vergency occurred in sub-horizontal bed position and probably characterise original tectonic emplacement. New fission-track zircon ages (140–134 Ma) would constraint this deformation as earliest Cretaceous.

- The geometry and magma-flow direction of the Szarvaskő Complex may point to a ring dyke-pillow basalt complex. New Sm-Nd dating on garnets from the contact of gabbro and plagiogranite yielded an age of 164.7 ± 1.6 Ma, in agreement with previous K-Ar ages (Árváné Sós et al. 1987; 166 ± 8 Ma on magmatic amphibole and 165 ± 5 Ma on muscovites of the contact hornfels), confirming a Callovian age of the magmatic rocks.
- Whole-rock geochemical data of the magmatites display MORB-like characteristics with subduction-related imprints, therefore previous authors (Harangi et al. 1996, Aigner-Torres & Koller 1999) suggested that the Jurassic magmatism of the Szarvaskő Complex is related to the opening of a back-arc/marginal basin. The differences from a real mid-oceanic ridge setting are also supported by the observed limited submarine

hydrothermal processes and the lava-unconsolidated sediment interactions, i.e. the presence of the siliciclastic peperitic facies (Kiss et al. 2016).

- However, there is a strong connection between the Szarvaskő and Bükk Complexes, as the clastics hosting the magmatites have strong affinities to the Bükkian Lökölgy Fm. and to the Vaskapu Sandstone Fm. It seems to be unrealistic to consider Szarvaskő as part of the upper plate ophiolite nappe system.

Our suggestion is that all complexes of the Bükk Mts. were derived from the same basin, situated on the down-going slab. Their original position was within the distal continental margin, possibly approaching the trench of the intra-oceanic subduction. Some exotic clasts could be detached during incipient thrusting and redeposited immediately. The Darnó complex, as a mélange, could be linked to the once overlying ophiolite.

The research was supported by the found NKFIH OTKA 113013, the ÚNKP-17-2 New National Excellence Program, Ministry of Human Capacities, and by the Austro-Hungarian Action Foundation (92öu11).

REFERENCES

- Aigner-Torres M. & Koller F., 1999. Nature of the magma source of the Szarvaskő complex (NE-Hungary). *Ofioliti*, 24, 1–12.
- Árváné Sós E., Balogh K., Ravaszné Baranyai L. & Ravasz Cs., 1987. Mezozoos magnás kőzetek K/Ar kora Magyarországi egyes területein. *Magyar Állami Földtani Intézet Évi Jelentése*, 295–308.
- Balla Z., 1983. Stratigraphy and tectonics of the Szarvaskő synform. *Annual Report Eötvös Geophysical Institute Hungary*, 1982, 42–65.
- Csontos L., 1999. A Bükk hegység szerkezetének főbb vonásai. Structural outline of the Bükk Mts (N. Hungary). *Földtani Közlöny*, 129, 611–651.
- Csontos L., 2000. A Bükk hegység mezozoos rétegtani újraértékelése. Stratigraphie réévaluation of the Bükk Mts (N. Hungary). *Földtani Közlöny*, 130, 95–131.
- Haas J., Pelikán P., Görög Á., Józsa S. & Ozsvárt P., 2013. Stratigraphy, facies and geodynamic settings of Jurassic formations in the Bükk Mountains, North Hungary: its relations with the other areas of the Neotethyan realm. *Geological Magazine*, 150, 1, 18–49.
- Harangi Sz., Szabó Cs., Józsa S., Szoldán Zs., Árváné Sós E., Balla M. & Kubovics I., 1996. Mesozoic Igneous Suites in Hungary: Implications for Genesis and Tectonic Setting in the Northwestern Part of Tethys. *International Geology Review*, 38, 336–360.
- Kiss G.B., Molnár F. & Palinkaš L.A., 2016. Hydrothermal processes related to some triassic and jurassic submarine basaltic complexes in northeastern Hungary, the dinarides and hellenides. *Geologia Croatica*, 69, 1, 39–64.
- Kiss G., Molnár F., Kovács S. & Palinkaš L., 2010. Field characteristics and petrography of the advanced rifting-related Triassic submarine basaltic blocks in the Jurassic mélange of the Darnó Unit. *Central European Geology*, 53, 2–3, 181–204.
- Kovács S., Haas J., Ozsvárt P., Palinkaš L.A., Kiss G., Molnár F., Józsa S. & Kövér Sz., 2011. Re-evaluation of the Mesozoic complexes of Darnó Hill (NE Hungary) and comparisons with Neotethyan accretionary complexes of the Dinarides and Hellenides – preliminary data. *Central European Geology*, 53/2–3, 205–231.
- Less Gy., Kovács S., Pelikán P., Pentelényi L. & Sásdi L., 2005. *Explanatory Book, Geological Map of the Bükk Mountains 1:50000*. Geological Institute Hungary.

3D structural model of Miocene depositional environment and faults for a future LBr-1 pilot CCS in the Vienna basin

Juraj Franců¹, Miroslav Pereszlényi¹, Lukáš Jurenka¹, Slavomír Nehyba², Přemysl Kyselák³, Oldřich Krejčí¹, Vít Hladík¹

¹ Czech Geological Survey; Leitnerova 22, 65869 Brno, Czech Republic; e-mail: juraj.francu@geology.cz

² Masaryk University, Faculty of Science, Department of Geological Sciences; Kotlarska 2, 61137 Brno, Czech Republic; e-mail: slavek@sci.muni.cz

³ MND a.s., Uprkova 807/6, 69501 Hodonín, Czech Republic; e-mail: kyselak@mnd.cz

The LBr-1 is an abandoned oil and gas field Lanžhot – Brodske located at the Czech and Slovak border in the

Vienna basin. The Lanžhot block forms a southern part of the Hodonín – Gbely horst in the Czech part of the