

ARCHIVES OF THE HUNGARIAN MEDICAL ASSOCIATION OF AMERICA



The official publication of the Hungarian Medical Association of America, Inc.

HMAA

Várgesztes Meeting

2024

ABSTRACTS

TARGETING HARD-TO-ATTACK PROTEIN-PROTEIN INTERACTION SURFACES USING UV-REACTIVE SURFACE FRAGMENTS

Bence Nagymihály, Edit Wéber, Péter Ábrányi-Balogh, Dóra Karancsiné Menyhárd, Nikolett Péczka, Márton Gadanecz, Gitta Schlosser, Zoltán Orgován, Ferenc Bogár, Dávid Bajusz, Gábor Kecskeméti, Zoltán Szabó, Éva Bartus, Attila Tököli, Gábor K. Tóth, Tibor V. Szalai, Tamás Takács, 6 Lászl (Department of Medical Chemistry, University of Szeged; HUN-REN-SZTE Biomimetic Systems Research Group; Medicinal Chemistry Research Group, HUN-REN Research Centre for Natural Sciences; Laboratory of Structural Chemistry and Biology, Institute of Chemistry, Eötvös Loránd University; MTA-ELTE Lendület Ion Mobility Mass Spectrometry Research Group, Institute of Chemistry; Eötvös Loránd University; Institute of Molecular Life Sciences, HUN-REN Research Centre for Natural Sciences)

Keywords: KRAS, peptidomimetics, HPLC-MS, NMR, fragment-based drug design

Aims: 80% of the human proteome consists of hard-to-attack proteins without targetable binding pocket. Efficient strategies like fragment-based drug development are not available for these proteins. Our goal was to map the surface of difficult-to-attack targets with peptidomimetic fragments and then increase the affinity with linking the foldamer fragments, thus extending fragment-based drug design to targets without a binding pocket.

Methods: Experiments were performed on one of the most difficult target proteins, KRas G12D, which has a high contribution to many tumors. We tested a 100-membered foldamer library, containing a UV-reactive group, so that the bound foldamers covalently labelled the target that could be identified by HPLC-MS. The binding site of the most effective fragments were determined by proteomic MS-MS analysis and ¹⁵N HSQC NMR titration. In dynamic covalent library experiments, foldamer dimers with higher affinity were created.

Results: The surface fragments were able to recognize the protein surface and bind to the target. We found numerous foldamers with a side chain-dependent binding pattern with KRas G12D. The UV-reactive marker made the screening efficient, and we were able to localize the binding site of the fragments. The foldamers found the

known protein-protein interaction surfaces of the target, but also discovered new, potentially attackable regions. In protein-templated dynamic covalent library experiments, dimers with higher affinity were created.

Conclusions: Using the protein surface mimetic foldamer library and connecting the fragments can be a viable general hit generation strategy for challenging protein targets.