

Mathematical Biosciences

A mathematical model for cancer dynamics with treatment and saboteur bacteria --Manuscript Draft--

Manuscript Number:	MBS-D-25-00222
Article Type:	Original research article
Keywords:	Cancer treatment; Therapeutic resistance; Differential equation model; Global dynamics; Bifurcations
Corresponding Author:	Anna Geretovszky HCEMM HUNGARY
First Author:	Anna Geretovszky
Order of Authors:	Anna Geretovszky Gergely Röst
Abstract:	We construct a mathematical model of cancer dynamics with chemotherapeutic treatment, in the presence of bacteria that are capable of metabolizing the chemotherapeutic drug, hence sabotaging the treatment. We investigate the possibility of complementing the cancer treatment with antibiotic drugs, thus eradicating the bacteria or at least mitigating their negative impact on the prospects of therapy. Our model is a nonlinear system of four differential equations, for which we perform a complete analysis, explicitly characterizing the four possible outcomes, depending on whether the cancer cells or the bacteria extinct or persist. Global stability results are proven by the iterative application of a comparison principle, and a bifurcation diagram is created to show the transitions between scenarios with respect to the controllable parameters. We apply our model to an experiment on mice with colon cancer and the drug Gemcitabine.
Opposed Reviewers:	

Highlights

1. Complementary antibiotic therapy to mitigate the sabotaging effect of bacteria in anti-cancer treatments.
2. New mathematical model describing the interactions between the cancer cells, bacteria and the drugs.
3. Identification of compound threshold parameters that determine the outcome of the treatment.
4. Full analysis of globally asymptotically stable equilibria and bifurcations.
5. The model accurately replicates real-world data from experiments on mice.

A mathematical model for cancer dynamics with treatment and saboteur bacteria

Anna Geretovszky^{a,b} and Gergely Röst^{a,b,c}

^aBolyai Institute, University of Szeged, Szeged H-6720, Hungary

^bHungarian Centre of Excellence for Molecular Medicine, Szeged H-6728, Hungary

^cNational Laboratory for Health Security, University of Szeged, Szeged H-6720, Hungary

Correspondence to be sent to: Anna Geretovszky, anna.geretovszky@hcemm.eu

ABSTRACT

We construct a mathematical model of cancer dynamics with chemotherapeutic treatment, in the presence of bacteria that are capable of metabolizing the chemotherapeutic drug, hence sabotaging the treatment. We investigate the possibility of complementing the cancer treatment with antibiotic drugs, thus eradicating the bacteria or at least mitigating their negative impact on the prospects of therapy. Our model is a nonlinear system of four differential equations, for which we perform a complete analysis, explicitly characterizing the four possible outcomes, depending on whether the cancer cells or the bacteria extinct or persist. Global stability results are proven by the iterative application of a comparison principle, and a bifurcation diagram is created to show the transitions between scenarios with respect to the controllable parameters. We apply our model to an experiment on mice with colon cancer and the drug Gemcitabine.

KEY WORDS

Cancer treatment; Therapeutic resistance; Differential equation model; Global dynamics; Bifurcations

1 Introduction

Cancer remains one of the most pressing challenges in modern medicine. It encompasses a diverse group of diseases characterized by uncontrolled cell division, leading to tumour formation and, in many cases, metastasis to distant organs. Without effective treatment, cancer is fatal, and its global burden continues to rise, partly due to population growth and aging. While advances in medical research, early detection, and risk factor reduction — such as decreased smoking rates — have contributed to a decline in age-standardized cancer mortality rates, current therapies remain ineffective for many patients. In 2025, the global incidence of cancer is projected to reach 21.33 million new cases, with an estimated 10.41 million deaths. Cancer is the second leading cause of mortality worldwide, accounting for approximately 14–19% of all deaths, following cardiovascular diseases ([1, 2, 3]).

Mathematical tools are increasingly being applied in biology and medicine, including oncology ([4, 5, 6]), to aid in the fight against cancer. Over the years, numerous mathematical models have been developed to study cancer dynamics, providing valuable insights into tumour growth, treatment responses, and disease progression. Comprehensive overviews of these models can be found in [7, 8].

There are several strategies for eliminating harmful cells in a patient's body, one of which is chemotherapy. This treatment employs chemical agents to target the tumour, making it particularly useful for inoperable cases. Chemotherapeutic drugs primarily attack rapidly dividing cells, a hallmark of cancer cells ([9, 10]). However, because these drugs are not selective for cancer cells alone, they also damage healthy tissues, leading to significant side effects. Therefore, optimizing treatment strategies to minimize the overall chemotherapy dosage

while maintaining efficacy is crucial. A major challenge in cancer therapy is the development of drug resistance, which can render chemotherapy ineffective. Depending on the type of cancer, there can be different reasons and mechanisms for the emergence of resistance ([9, 11]). One key process is natural selection: cells in the tumour mutate and the ones that acquired hereditary properties that are advantageous against chemotherapy become the majority ([12]). Another mechanism is cellular plasticity, where cells adapt their phenotype in response to environmental cues without genetic mutations. For instance, some cells develop the ability to pump drugs out of themselves using proteins, which is associated with the MDR (multidrug resistant) phenotype ([13]). Additionally, microvesicles, which play a role in intercellular communication between cancer cells, may also contribute to the development of resistance. These microvesicles can transfer resistance-conferring molecules from resistant to non-resistant cells, effectively spreading resistance throughout the tumour, much akin to the spreading of an infection ([14, 15]).

Several mathematical models have been developed to account for drug resistance in cancer treatment ([16]). The motivation for this paper is the findings of the article [17], which describe the relatively recent scientific discovery that microorganisms can influence the effectiveness of chemotherapy treatments targeting cancer tumour cells. Certain (not necessarily malignant) bacteria in the patient's body can metabolize chemotherapy drugs. Simply speaking, the bacteria "consume" the anti-cancer drug, hence sabotaging the treatment by preventing the drug from reaching the targeted cells, effectively making the tumour resistant to the therapy. In such cases, chemotherapy may need to be supplemented with antibiotics to reduce – or even eliminate – the sabotaging bacterial population, thus opening the way to a more effective chemotherapy treatment. However, excessive antibiotic use must be avoided, as it can further compromise the patient's already weakened immune system and contribute to antibiotic resistance. Therefore, identifying the optimal combination and dosage of both chemotherapeutic and antibiotic agents is crucial to maximizing treatment efficacy while minimizing adverse effects.

In this paper, we construct a dynamic model to represent the underlying biological mechanisms of tumour-bacteria-drug interactions. The model consists of a nonlinear system of four differential equations that describe the temporal variations of tumour size, the population of drug-metabolizing intra-tumour bacteria, and the concentration of both the chemotherapeutic drug and the antibiotics. To characterize the possible treatment outcomes, we introduce three combined threshold parameters that distinguish four scenarios based on the persistence or eradication of the tumour and the bacterial population. We identify the therapies that achieve the desired outcome, when both the tumour and the bacteria are eradicated. The main result of this paper is that for all four scenarios, we prove the global asymptotic stability of the equilibrium to which the solutions converge. The main tool of the proof is the iterative application of a comparison principle. We also conduct a bifurcation analysis in the parameter space of drug doses, and develop the full bifurcation diagram, demonstrating that single and simultaneous transcritical bifurcations separate these scenarios.

We calibrate a specific version of our model using real-world data, demonstrating its accuracy in capturing key biological dynamics. Finally, we explore potential extensions of our model, including more realistic tumour growth laws and the incorporation of pharmacokinetics and pharmacodynamics. These lead to more complicated systems, but we show that several of our qualitative results remain valid in these extended models.

2 Model

2.1 Construction

We aim to set up a minimal model that already includes all the relevant biological and pharmacological components. Possible further extensions and more general models will be discussed in Section 4. In constructing our model, we assume a uniform spatial distribution and a well-mixed system, so we use ordinary differential equations. We construct our dynamical model step-by-step, deriving from biological mechanisms, where every variable and parameter has a clear biological interpretation.

Let $T(t)$ denote the size of the tumour as a function of time. The choice of the growth function is heavily influenced by the specific type of cancer being modeled, as well as the environment in which tumour cells proliferate (e.g., the human body or an in vitro laboratory setting). In [18], various cancer types have been modeled using different functional forms. Among these, the power-law model $T'(t) = rT^a(t)$ was identified as the most accurate fit in the majority of cases. However, the authors highlight its extreme sensitivity to parameterization, making it unreliable for predictive purposes. Moreover, this model assumes unrestricted tumour growth, contradicting the biological reality that tumour expansion is constrained by factors such as nutrient availability and immune response. Additionally, the parameter a lacks direct biological interpretation, meaning the model is not mechanistic. For these reasons, we do not consider this model suitable for our study. In addition to the power-law model, the logistic model performed well, especially for cancer types that are usually




Fig. 1: (a) Based on biological intuition, a possible scenario for the change of tumour size ($T(t)$) in response to different externalities. Here, at time C the chemotherapy treatment is started, decreasing the tumour size. At time B the patient is infected by the bacteria, and at time A the patient receives the antibiotics treatment. (b) Recreation of the same scenario by a numerical simulation using our mathematical model. Additional to variation of the tumour size, we plot the drug concentrations and bacterial density as well.

94 treated with the chemotherapeutic drug under investigation in [17]. Hence we use this to describe the variation
95 of the tumour volume:

$$T'(t) = r_T T(t) (K_T - T(t)),$$

96 where r_T is the intrinsic tumour growth rate and K_T is the carrying capacity of the environment wrt. the tumour
97 size.

98 Let $C(t)$ denote the concentration of the chemotherapy drug as a function of time. In this section, we use
99 a constant drug dose (γ) and assume that the drug is absorbed and excreted from the body in proportion to
100 its concentration (with rate μ). The appearance of the drug also affects the size of the tumour: assuming mass
101 action law, the cytotoxic effect is proportional to the drug concentration and the number of tumour cells (with
102 rate δ). So our system of equations for the tumour and the chemotherapeutic drug is:

$$\begin{cases} T'(t) = r_T T(t) (K_T - T(t)) - \delta T(t) C(t), \\ C'(t) = \gamma - \mu C(t). \end{cases}$$

103 Let $B(t)$ denote the size of the bacterial population as a function of time. Its growth can also be modelled
104 by a logistic equation, with intrinsic growth rate r_B and carrying capacity K_B . The core phenomenon modelled
105 in this paper is that the bacteria can metabolize the chemotherapeutic drug, thus reducing its amount, where
106 again we use mass action law with rate λ . The resulting system of equations is

$$\begin{cases} T'(t) = r_T T(t) (K_T - T(t)) - \delta T(t) C(t), \\ C'(t) = \gamma - \mu C(t) - \lambda C(t) B(t), \\ B'(t) = r_B B(t) (K_B - B(t)). \end{cases}$$

107 Finally, we introduce the antibiotics that is used to kill the bacteria. Let $A(t)$ denote the concentration
108 of this as a function of time. Here again, we are working with a constant dose (α) and the antibiotic drug is
109 removed with rate σ . According to the mass action law, the bacterial population is killed by the antibiotics
110 in proportion to the concentration of the antibiotics and the amount of bacteria (with rate θ). This gives our
111 system of equations consisting all essential components:

$$\begin{cases} T'(t) = r_T T(t) (K_T - T(t)) - \delta T(t) C(t), \\ C'(t) = \gamma - \mu C(t) - \lambda C(t) B(t), \\ B'(t) = r_B B(t) (K_B - B(t)) - \theta B(t) A(t), \\ A'(t) = \alpha - \sigma A(t). \end{cases} \quad (1)$$

112 The model variables and the nonnegative parameters are summarized in Table 1.

variable/param.	meaning of the variable/parameter
T	size of the tumour
C	concentration of the chemotherapeutic drug
B	size of the bacterial population
A	concentration of the antibiotics
r_T	growth rate of the tumour
K_T	carrying capacity wrt. the tumour
δ	rate of death by the chemotherapeutic drug
γ	dose of the chemotherapeutic drug
μ	absorption or elimination of the chemotherapeutic drug
λ	rate of chemotherapeutic drug metabolized by the bacteria
r_B	growth rate of the bacterial population
K_B	carrying capacity wrt. the bacteria
θ	rate of death of bacteria by the antibiotics
α	dose of the antibiotics
σ	absorption or elimination of the antibiotics

Table 1: Variables and parameters in the equation system (1).

2.2 Analysis

In our model, we have defined variables that describe the size of a population or the concentration of a drug as a function of time. These can obviously only give realistic values if they are non-negative. For this reason, first we show that starting from non-negative initial values, the solutions remain non-negative.

Proposition 2.1. *The equation system (1) is well-defined. Starting from nonnegative initial values at t_0 , unique solutions exist on the whole $[t_0, \infty)$ interval, moreover, they remain nonnegative and bounded.* $\square$

Proof. By the Picard-Lindelöf theorem, the equation system (1) has a unique solution. For the nonnegativity part, we use the following well-known statement ([19] Prop. A.1). Let $x'(t) = F(t, x)$ be a differential equation system, where $x(t) = (x_1(t), \dots, x_n(t))$, $F(t, x) = (F_1(t, x), \dots, F_n(t, x))$, and $F(t, x)$ is defined for all $t \geq 0, x \in \mathbb{R}^n$. If the solutions of the initial value problems $x(t_0) = x^0$ are unique for $x^0 \in [0, \infty)^n, t_0 \geq 0$ and for all $i = 1, \dots, n$

$$F_i(t, x) \geq 0 \text{ whenever } x \in [0, \infty)^n, x_i = 0, t \geq 0,$$

then $x(t) \in [0, \infty)^n$ for all $t \geq t_0 \geq 0$, whenever $x(t_0) \in [0, \infty)^n$. This clearly holds for system (1), hence the solutions remain nonnegative.

The boundedness is concluded from the following observation: when $T(t) \geq K_T$, then $T'(t) < 0$, which means that $\limsup_{t \rightarrow \infty} T(t) \leq K_T$. The same stands in the case of $C(t), B(t)$ and $A(t)$, with bounds $\frac{\gamma}{\mu}, K_B$ and $\frac{\alpha}{\sigma}$, respectively. E.g., $\limsup_{t \rightarrow \infty} C(t) \leq \frac{\gamma}{\mu}$, $\limsup_{t \rightarrow \infty} B(t) \leq K_B$ and $\limsup_{t \rightarrow \infty} A(t) \leq \frac{\alpha}{\sigma}$.

Since the solutions are bounded, they exist on the whole $[t_0, \infty)$ interval. $\blacksquare$

Definition 2.2. Let us define the following compound threshold parameters:

$$\begin{aligned} \mathcal{F}_1 &:= \frac{K_B \sigma r_B}{\theta} - \alpha, \\ \mathcal{F}_2 &:= \frac{K_T \mu r_T}{\delta} - \gamma, \\ \mathcal{F}_3 &:= \frac{K_T \left(\mu + \lambda \left(K_B - \frac{\theta \alpha}{\sigma r_B} \right) \right) r_T}{\delta} - \gamma. \end{aligned}$$

We categorize each equilibrium $E_* = (T_*, C_*, B_*, A_*)$ into one of the following types:

- E_0 : trivial equilibrium, when both the tumour and the bacteria extinct, i.e. $T_* = 0$ and $B_* = 0$;
- E_T : tumour-free equilibrium, when the tumour extincts, but the bacteria persist, i.e. $T_* = 0$ and $B_* > 0$;
- E_B : bacteria-free equilibrium, when the tumour survives, but the bacteria extincts, i.e. $T_* > 0$ and $B_* = 0$;

- E_{coex} : coexisting equilibrium, when both the tumour cells and the bacteria persist, $T_* > 0$ and $B_* > 0$.

Theorem 2.3 (Equilibrium solutions). *The threshold parameters characterize which types of equilibria of system (1) exist, as follows:*

Case 1: if $\mathcal{F}_1 > 0$ and $\mathcal{F}_3 > 0$

a) if $\mathcal{F}_2 > 0$, then all four types of equilibria exist;

b) if $\mathcal{F}_2 \leq 0$, then E_0 , E_T and E_{coex} exist;

Case 2: if $\mathcal{F}_1 > 0$ and $\mathcal{F}_3 \leq 0$, then E_0 and E_T exist;

Case 3: if $\mathcal{F}_1 \leq 0$ and $\mathcal{F}_2 > 0$, then E_0 and E_B exist;

Case 4: if $\mathcal{F}_1 \leq 0$ and $\mathcal{F}_2 \leq 0$, then only E_0 exists.

The cases are depicted in Figure 2. □

Proof. We find an equilibrium solution (T^*, C^*, B^*, A^*) of the system (1) from the steady state equations

$$\begin{cases} 0 = r_T T^* (K_T - T^*) - \delta T^* C^*, \\ 0 = \gamma - \mu C^* - \lambda C^* B^*, \\ 0 = r_B B^* (K_B - B^*) - \theta B^* A^*, \\ 0 = \alpha - \sigma A^*. \end{cases}$$

From the last equation we get that $A^* = \frac{\alpha}{\sigma}$, substituting it back to the third one we have

$$\begin{aligned} 0 &= r_B B^* (K_B - B^*) - \theta \frac{\alpha}{\sigma} B^* \\ &= r_B B^* \left(\left(K_B - \frac{\theta \alpha}{\sigma r_B} \right) - B^* \right), \end{aligned}$$

which leads to two possibilities for B^* :

$$B_1^* = 0 \quad \text{and} \quad B_2^* = K_B - \frac{\theta \alpha}{\sigma r_B}.$$

Since $B(t)$ is nonnegative, B_2^* can only exist if $K_B - \frac{\theta \alpha}{\sigma r_B} \geq 0$, which is exactly the case when $\mathcal{F}_1 \geq 0$. From now on we suppose that $\mathcal{F}_1 > 0$, i.e. B_2^* is positive. We will discuss the case of equality later. Now we substitute the possible B components of the equilibria into the equation for C :

$$0 = \gamma - \mu C^* - \lambda B^* C^* = \gamma - (\mu + \lambda B^*) C^*,$$

thus $C^* = \frac{\gamma}{\mu + \lambda B^*}$, i.e.

$$\begin{aligned} C_1^* &= \frac{\gamma}{\mu + \lambda B_1^*} = \frac{\gamma}{\mu} \quad \text{and} \\ C_2^* &= \frac{\gamma}{\mu + \lambda B_2^*} = \frac{\gamma}{\mu + \lambda \left(K_B - \frac{\theta \alpha}{\sigma r_B} \right)}, \end{aligned}$$

where for the existence of $C_2^* > 0$ we also need $\mathcal{F}_1 > 0$ to stand. Lastly, we substitute the derived C^* values into the first equation:

$$\begin{aligned} 0 &= r_T T^* (K_T - T^*) - \delta C^* T^* \\ &= r_T T^* \left(\left(K_T - \frac{\delta}{r_T} C^* \right) - T^* \right), \end{aligned}$$

hence the possible T components of the equilibrium points are $T_1^* = 0$ and $T_2^* = K_T - \frac{\delta}{r_T} C^*$, i.e.

$$\begin{aligned} T_{2,1}^* &= K_T - \frac{\delta}{r_T} C_1^* = K_T - \frac{\delta \gamma}{\mu r_T} \quad \text{and} \\ T_{2,2}^* &= K_T - \frac{\delta}{r_T} C_2^* = K_T - \frac{\delta \gamma}{\left(\mu + \lambda \left(K_B - \frac{\theta \alpha}{\sigma r_B} \right) \right) r_T}. \end{aligned}$$




Fig. 2: Illustration of the five possible cases in Theorem 2.3, in the parameter plane of the two drug doses. The equilibria with bold are the stable ones discussed in Theorem 2.7. The equations of the (half) lines on the boundaries: vertical: $\mathcal{F}_1 = 0$, horizontal: $\mathcal{F}_2 = 0$, tilted on the left: $\mathcal{F}_3 = 0$. The figure was made using the parameters $r_T = 1.71$, $K_T = 1.61$, $\delta = 3.59$, $\mu = 1.78$, $\lambda = 2$, $r_B = 1.81$, $K_B = 1.81$, $\theta = 1.085$, $\sigma = 0.7$ and $\alpha, \gamma \in [0, 5]$.

Since $T(t)$ is nonnegative, T_2^* can only exist if $K_T - \frac{\delta}{r_T}C^* \geq 0$, but similarly to B_2^* , we also require the strict inequality, so T_2^* is positive. In the case of $T_{2,1}^*$, the condition $K_T - \frac{\delta}{r_T}C^* > 0$ stands iff $\mathcal{F}_2 > 0$, while in the case of $T_{2,2}^*$, it is equivalent to $\mathcal{F}_3 > 0$. For the existence of $T_{2,2}^*$, we also need $\mathcal{F}_1 > 0$, inherited from C_2^* .

We look separately at the cases when there is equality somewhere. If $\mathcal{F}_1 = 0$, then $B_1^* = B_2^* = 0$, i.e. the bacteria are not present. If $\mathcal{F}_2 = 0$ or $\mathcal{F}_3 = 0$, then $T_{2,1}^* = 0$ or $T_{2,2}^* = 0$, respectively, so in these situations the tumour is absent.

Hence, we obtain that if $\mathcal{F}_1 \leq 0$, then there is no positive B_2^* . Furthermore, if $\mathcal{F}_2 \leq 0$ also, then a positive T_2^* does not exist either, so the only possibly equilibrium is E_0 , otherwise E_0 and E_B exist. If the condition $\mathcal{F}_1 > 0$ stands, then there exist a positive B_2^* . In this case, similarly, if $\mathcal{F}_3 \leq 0$, then there is no positive T_2^* , so only the equilibrium points E_0 and E_T exist. Finally, if the condition $\mathcal{F}_3 > 0$ holds, then there are two possible

scenarios: if $\mathcal{F}_2 \leq 0$, then there is no positive T_2^* associated with $B = 0$, so only E_0, E_T and E_{coex} exist, while in the other case, all four types of equilibria exist.




Fig. 3: Derivation of the possible coordinates of the equilibria depending on the signs of the threshold parameters.

168

Proposition 2.4. *There are just these five possible scenarios, since in Figure 2, the three lines $\mathcal{F}_1 = 0$, $\mathcal{F}_2 = 0$ and $\mathcal{F}_3 = 0$ intersect in the same point.* □

Proof. From $\mathcal{F}_1 = 0$ we obtain that $\tilde{\alpha} = \frac{K_B \sigma r_B}{\theta}$, which substituted into $\mathcal{F}_3 = 0$ gives us

$$\begin{aligned}
 \mathcal{F}_3(\tilde{\alpha}) &= \frac{K_T \left(\mu + \lambda \left(K_B - \frac{\theta \tilde{\alpha}}{\sigma r_B} \right) \right) r_T}{\delta} - \gamma \\
 &= \frac{K_T \left(\mu + \lambda \left(K_B - \frac{\theta \frac{K_B \sigma r_B}{\theta}}{\sigma r_B} \right) \right) r_T}{\delta} - \gamma \\
 &= \frac{K_T (\mu + \lambda \cdot 0) r_T}{\delta} - \gamma = \frac{K_T \mu r_T}{\delta} - \gamma = \mathcal{F}_2(\tilde{\alpha}),
 \end{aligned}$$

i.e. the lines $\mathcal{F}_1 = 0$ and $\mathcal{F}_3 = 0$ intersect on the line $\mathcal{F}_2 = 0$. ■

So, we have seen that our system of equations (1) has four possible types of equilibria, and we have shown when each of them exists depending on the parameters of the equation system. In the following, we will determine the stability of these equilibria and prove the convergence of the solutions to the stable one. To achieve this, we use Theorem 1.3. from [20].

Theorem 2.5 ([20]). *Suppose f is locally Lipschitz continuous with respect to x , uniformly in t . Let $x(t)$ and $y(t)$ be two differentiable functions such that*

$$x(t_0) \leq y(t_0), \quad x'(t) - f(t, x(t)) \leq y'(t) - f(t, y(t)), \quad t \in [t_0, \tilde{t}).$$

Then we have $x(t) \leq y(t)$ for every $t \in [t_0, \tilde{t})$. Moreover, if $x(t) < y(t)$ for some t , this remains true for all later times. □

Based on this theorem, we can prove the following comparison principle, a key tool in our proofs.

Proposition 2.6. *Let f and g be locally Lipschitz continuous functions, and let $x(t)$ and $y(t)$ be two differentiable functions such that*

$$x'(t) = f(x(t)), \quad y'(t) = g(y(t)), \quad \text{and} \quad x(t_0) \leq y(t_0), \quad t \in [t_0, \tilde{t}).$$

Suppose that $f(x) \leq g(x)$ holds for every x , then $x(t) \leq y(t)$ for $t \in [t_0, \tilde{t})$. □

185 **Proof.** Since $x'(t) = f(x(t))$ and $y'(t) = g(y(t))$, we see that

$$x'(t) - f(x(t)) = 0 = y'(t) - g(y(t)).$$

186 Since $f(x) \leq g(x)$ holds for all x , it follows that

$$x'(t) - f(x(t)) = y'(t) - g(y(t)) \leq y'(t) - f(y(t)).$$

187 Then f , $x(t)$ and $y(t)$ satisfy the conditions of Theorem 2.5, from which we obtain $x(t) \leq y(t)$ for all $t \in [t_0, \tilde{t})$. ■

188 Let $\mathbb{R}^+$ and $\mathbb{R}_0^+$ denote the set of the positive and nonnegative real numbers, respectively.

189 **Theorem 2.7** (Convergence and stability). *In Case 1 the equilibrium E_{coex} , in Case 2 the equilibrium E_T , in*
 190 *Case 3 the equilibrium E_B , and in Case 4 the equilibrium E_0 is globally asymptotically stable in the sense that it*
 191 *is stable and attracts all solutions starting from $(T(t_0), C(t_0), B(t_0), A(t_0)) \in \mathbb{R}^+ \times \mathbb{R}_0^+ \times \mathbb{R}^+ \times \mathbb{R}_0^+$. Analogous*
 192 *results hold in the subspaces where either $T = 0$ or $B = 0$, according to Table 2.*

Domain \ Case	Case 1	Case 2	Case 3	Case 4
$\mathbb{R}^+ \times \mathbb{R}_0^+ \times \mathbb{R}^+ \times \mathbb{R}_0^+$	E_{coex} $= (T_{2,2}^*, C_2^*, B_2^*, A^*)$	E_T $= (0, C_2^*, B_2^*, A^*)$	E_B $= (T_{2,1}^*, C_1^*, 0, A^*)$	E_0 $= (0, C_1^*, 0, A^*)$
$\{0\} \times \mathbb{R}_0^+ \times \mathbb{R}^+ \times \mathbb{R}_0^+$	E_T $= (0, C_2^*, B_2^*, A^*)$	E_T $= (0, C_2^*, B_2^*, A^*)$	E_0 $= (0, C_1^*, 0, A^*)$	E_0 $= (0, C_1^*, 0, A^*)$
$\mathbb{R}^+ \times \mathbb{R}_0^+ \times \{0\} \times \mathbb{R}_0^+$	a) E_B $= (T_{2,1}^*, C_1^*, 0, A^*)$	E_0 $= (0, C_1^*, 0, A^*)$	E_B $= (T_{2,1}^*, C_1^*, 0, A^*)$	E_0 $= (0, C_1^*, 0, A^*)$
	b) E_0 $= (0, C_1^*, 0, A^*)$			
$\{0\} \times \mathbb{R}_0^+ \times \{0\} \times \mathbb{R}_0^+$	E_0 $= (0, C_1^*, 0, A^*)$	E_0 $= (0, C_1^*, 0, A^*)$	E_0 $= (0, C_1^*, 0, A^*)$	E_0 $= (0, C_1^*, 0, A^*)$

Table 2: Possible stable equilibria of the equation system (1).

193 □

194 **Proof.** It is easy to see that for $T(t_0) = 0$, the solution $T(t) \equiv 0$, and for $B(t_0) = 0$, the solution $B(t) \equiv 0$.
 195 This implies that these two functions will always take the value 0, no matter how the parameters of the system
 196 of equations are chosen, so they cannot converge to positive equilibrium situations. If $B(t_0) = 0$, we also have
 197 to take into consideration the sign of $\mathcal{F}_2$ in Cases 1 and 2. Because of $B(t) \equiv 0$, the positive equilibrium $T_{2,2}^*$
 198 cannot exist, but for the existence of a positive $T_{2,1}^*$ we need the condition $\mathcal{F}_2 > 0$, which we do not have here.
 199 In Case 1, we already did a separation (Case 1 a) and b)), and in Case 2, $\mathcal{F}_1 > 0$ and $\mathcal{F}_3 \leq 0$ imply $\mathcal{F}_2 \leq 0$, so
 200 only $T_1^* = 0$ exists.

201 Since in the case of $T(t_0) = 0$ or $B(t_0) = 0$ the phenomenon under consideration would not occur (i.e.
 202 we would not have bacteria sabotaging the cancer treatment), we can assume that $T(t_0), B(t_0) \in \mathbb{R}^+$ and
 203 $C(t_0), A(t_0) \in \mathbb{R}_0^+$, i.e. we are working on the domain $D = \mathbb{R}^+ \times \mathbb{R}_0^+ \times \mathbb{R}^+ \times \mathbb{R}_0^+$, which is the most interesting
 204 case. The other scenarios listed in Table 2 can also be proved.

205 Consider first the equation $A'(t) = \alpha - \sigma A(t)$ for the function $A(t)$, which is an inhomogeneous linear
 206 equation, so it has the solution

$$A(t) = \frac{(\sigma A(t_0) - \alpha) e^{\sigma(t_0-t)} + \alpha}{\sigma},$$

207 from which we get

$$\lim_{t \rightarrow \infty} A(t) = \frac{\alpha}{\sigma}.$$

208 From that we can see the A component of the equilibrium point: $A^* = \frac{\alpha}{\sigma}$, and that $A(t)$ converges to A^* , e.g.
 209 $\forall \varepsilon_1 > 0 \exists t_1 :$

$$\text{if } t > t_1, \text{ then } \frac{\alpha}{\sigma} - \varepsilon_1 < A(t) < \frac{\alpha}{\sigma} + \varepsilon_1.$$

210 Next we look at the equation $B'(t) = r_B B(t) (K_B - B(t)) - \theta B(t) A(t)$ which describes the size of the
 211 bacteria population. Let

$$\begin{aligned}\underline{B}'_\varepsilon(t) &= r_B \underline{B}_\varepsilon(t) (K_B - \underline{B}_\varepsilon(t)) - \theta \underline{B}_\varepsilon(t) \left(\frac{\alpha}{\sigma} + \varepsilon \right) \\ &= r_B \underline{B}_\varepsilon(t) \left(\left(K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} + \varepsilon \right) \right) - \underline{B}_\varepsilon(t) \right)\end{aligned}$$

212 and

$$\begin{aligned}\overline{B}'_\varepsilon(t) &= r_B \overline{B}_\varepsilon(t) (K_B - \overline{B}_\varepsilon(t)) - \theta \overline{B}_\varepsilon(t) \left(\frac{\alpha}{\sigma} - \varepsilon \right) \\ &= r_B \overline{B}_\varepsilon(t) \left(\left(K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} - \varepsilon \right) \right) - \overline{B}_\varepsilon(t) \right).\end{aligned}$$

213 The equations for $\underline{B}_\varepsilon(t)$ and $\overline{B}_\varepsilon(t)$ will still be logistic if $K_B > \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} + \varepsilon \right)$, so the solutions are:

$$\begin{aligned}\underline{B}_\varepsilon(t) &= \frac{\underline{B}_\varepsilon(t_0) \left(K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} + \varepsilon \right) \right)}{\underline{B}_\varepsilon(t_0) + \left(\left(K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} + \varepsilon \right) \right) - \underline{B}_\varepsilon(t_0) \right) e^{\left(K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} + \varepsilon \right) \right) r_B (t_0 - t)}}, \\ \overline{B}_\varepsilon(t) &= \frac{\overline{B}_\varepsilon(t_0) \left(K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} - \varepsilon \right) \right)}{\overline{B}_\varepsilon(t_0) + \left(\left(K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} - \varepsilon \right) \right) - \overline{B}_\varepsilon(t_0) \right) e^{\left(K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} - \varepsilon \right) \right) r_B (t_0 - t)}}.\end{aligned}$$

214 Since for $t > t_1$ the inequalities $\underline{B}'_{\varepsilon_1}(t) \leq B'(t) \leq \overline{B}'_{\varepsilon_1}(t)$ stand, we get $\underline{B}_{\varepsilon_1}(t) \leq B(t) \leq \overline{B}_{\varepsilon_1}(t)$ from Proposi-
 215 tion 2.6 if we set $\underline{B}_{\varepsilon_1}(t_0) = \overline{B}_{\varepsilon_1}(t_0) = B(t_0)$. Note that

$$\lim_{t \rightarrow \infty} \underline{B}_{\varepsilon_1}(t) = K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} + \varepsilon_1 \right),$$

216 and

$$\lim_{t \rightarrow \infty} \overline{B}_{\varepsilon_1}(t) = K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} - \varepsilon_1 \right).$$

217 This implies that

$$K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} + \varepsilon_1 \right) \leq \liminf_{t \rightarrow \infty} B(t) \leq \limsup_{t \rightarrow \infty} B(t) \leq K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} - \varepsilon_1 \right).$$

218 Since these inequalities hold for all $\varepsilon_1 > 0$, we get that if $K_B - \frac{\theta\alpha}{\sigma r_B} > 0$ e.g. $\mathcal{F}_1 > 0$ stands, then

$$\lim_{t \rightarrow \infty} B(t) = K_B - \frac{\theta\alpha}{\sigma r_B} = B_2^*,$$

219 otherwise

$$\lim_{t \rightarrow \infty} B(t) = 0 = B_1^*,$$

220 so $B(t)$ converges to one of the B components of the equilibria characterized in Table 2.

221 Next we consider the equation describing the change of the concentration of the chemotherapy drug as
 222 follows: $C'(t) = \gamma - \mu C(t) - \lambda C(t) B(t)$. There are two cases based on which limit point of $B(t)$ we use.

223 1. If $B(t) \rightarrow B_1^* = 0$, e.g. $\forall \varepsilon_2 > 0 \exists t_2$:

$$\text{if } t > t_2, \text{ then } -\varepsilon_2 < B(t) < \varepsilon_2.$$

224 Let

$$\underline{C}'_\varepsilon(t) = \gamma - \mu \underline{C}_\varepsilon(t) - \lambda \underline{C}_\varepsilon(t) \varepsilon = \gamma - (\mu + \lambda \varepsilon) \underline{C}_\varepsilon(t)$$

225 and

$$\overline{C}'_\varepsilon(t) = \gamma - \mu \overline{C}_\varepsilon(t) + \lambda \overline{C}_\varepsilon(t) \varepsilon = \gamma - (\mu - \lambda \varepsilon) \overline{C}_\varepsilon(t)$$

The equations for $\underline{C}_\varepsilon(t)$ and $\overline{C}_\varepsilon(t)$ will still be inhomogeneous linear equations if $\mu - \lambda\varepsilon > 0$, so their solutions are:

$$\underline{C}_\varepsilon(t) = \frac{((\mu + \lambda\varepsilon)\underline{C}_\varepsilon(t_0) - \gamma)e^{(\mu + \lambda\varepsilon)(t_0 - t)} + \gamma}{\mu + \lambda\varepsilon},$$

$$\overline{C}_\varepsilon(t) = \frac{((\mu - \lambda\varepsilon)\overline{C}_\varepsilon(t_0) - \gamma)e^{(\mu - \lambda\varepsilon)(t_0 - t)} + \gamma}{\mu - \lambda\varepsilon}.$$

Since the inequalities $\underline{C}'_{\varepsilon_2}(t) \leq C'(t) \leq \overline{C}'_{\varepsilon_2}(t)$ stand when $t > t_2$, we get $\underline{C}_{\varepsilon_2}(t) \leq C(t) \leq \overline{C}_{\varepsilon_2}(t)$ from Proposition 2.6 if we set $\underline{C}_{\varepsilon_2}(t_0) = \overline{C}_{\varepsilon_2}(t_0) = C(t_0)$. Note that

$$\lim_{t \rightarrow \infty} \underline{C}_{\varepsilon_2}(t) = \frac{\gamma}{\mu + \lambda\varepsilon_2},$$

and

$$\lim_{t \rightarrow \infty} \overline{C}_{\varepsilon_2}(t) = \frac{\gamma}{\mu - \lambda\varepsilon_2}.$$

Since $\varepsilon_2 > 0$ was arbitrary, thus

$$\lim_{t \rightarrow \infty} C(t) = \frac{\gamma}{\mu} = C_1^*.$$

2. If $B(t) \rightarrow B_2^* = K_B - \frac{\theta\alpha}{\sigma r_B}$, e.g. $\forall \varepsilon_3 > 0 \exists t_3 :$

$$\text{if } t > t_3, \text{ then } K_B - \frac{\theta\alpha}{\sigma r_B} - \varepsilon_3 < B(t) < K_B - \frac{\theta\alpha}{\sigma r_B} + \varepsilon_3.$$

Similarly to the previous case, now we use these boundaries and the functions $\underline{C}_{\varepsilon_3}(t)$ and $\overline{C}_{\varepsilon_3}(t)$. Then applying Proposition 2.6. again, we get that in this case

$$\lim_{t \rightarrow \infty} C(t) = \frac{\gamma}{\mu + \lambda \left(K_B - \frac{\theta\alpha}{\sigma r_B} \right)} = C_2^*.$$

So we can see that $C(t)$ also converges to one of C components of the equilibria in Table 2.

Last we consider the equation $T'(t) = r_T T(t) (K_T - T(t)) - \delta T(t) C(t)$ describing the change of the tumour size. In a completely analogous way to the calculations done earlier for $B(t)$, we obtain the following.

1. In the case of $C(t) \rightarrow C_1^* = \frac{\gamma}{\mu}$, if $K_T - \frac{\delta\gamma}{\mu r_T} > 0$ e.g. $\mathcal{F}_2 > 0$, then

$$\lim_{t \rightarrow \infty} T(t) = K_T - \frac{\delta\gamma}{\mu r_T} = T_{2,1}^*,$$

otherwise

$$\lim_{t \rightarrow \infty} T(t) = 0 = T_1^*.$$

2. In the case of $C(t) \rightarrow C_2^* = \frac{\gamma}{\mu + \lambda \left(K_B - \frac{\theta\alpha}{\sigma r_B} \right)}$, if $\mathcal{F}_3 > 0$, then

$$\lim_{t \rightarrow \infty} T(t) = K_T - \frac{\delta\gamma}{\left(\mu + \lambda \left(K_B - \frac{\theta\alpha}{\sigma r_B} \right) \right) r_T} = T_{2,2}^*,$$

otherwise

$$\lim_{t \rightarrow \infty} T(t) = 0 = T_1^*.$$

So $T(t)$ also converges to one of the T components of the equilibria in Table 2.

Thus, we obtain that all four functions converge to the corresponding components of an equilibrium characterized in Table 2., i.e. the solutions converge. In the following, we investigate the global asymptotic stability of the equilibrium situations.

The stability of an equilibrium point is shown using $\varepsilon - \delta$ neighborhoods. Since we have a final dimensional system, we can show that componentwise, and for that, we will use the pinching functions again. Here we

investigate the stability of $B_2^* = K_B - \frac{\theta\alpha}{\sigma r_B}$, the rest can be proved in a similar way. Let $\varepsilon > 0$ be given and let $\varepsilon' = \varepsilon \frac{r_B}{\theta}$, then

$$\lim_{t \rightarrow \infty} \underline{B}_{\varepsilon'}(t) = K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} + \varepsilon' \right) = B_2^* - \varepsilon,$$

and

$$\lim_{t \rightarrow \infty} \overline{B}_{\varepsilon'}(t) = K_B - \frac{\theta}{r_B} \left(\frac{\alpha}{\sigma} - \varepsilon' \right) = B_2^* + \varepsilon.$$

Let $0 < \delta < \varepsilon$ be arbitrary and $B(t_0) \in (B_2^* - \delta, B_2^* + \delta)$. Now consider the functions $\underline{B}_{\varepsilon'}(t)$ and $\overline{B}_{\varepsilon'}(t)$ starting from the points $B_2^* - \delta$ and $B_2^* + \delta$, respectively, i.e.

$$\underline{B}_{\varepsilon'}(t_0) = B_2^* - \delta \text{ and } \overline{B}_{\varepsilon'}(t_0) = B_2^* + \delta.$$

We know that the positive equilibrium of a logistic equation is asymptotically stable, moreover, the solution converges to it monotonically. This means that the $\underline{B}_{\varepsilon'}(t)$ function starting from the point $B_2^* - \delta$ will converge to $B_2^* - \varepsilon$ in a monotonically decreasing way, and the function $\overline{B}_{\varepsilon'}(t)$ starting from the point $B_2^* + \delta$ will converge to $B_2^* + \varepsilon$ in a monotonically increasing way. Thus both of these functions stay in the interval $(B_2^* - \varepsilon, B_2^* + \varepsilon)$. Since the conditions of Proposition 2.6 hold, we get that $\underline{B}_{\varepsilon'}(t) \leq B(t) \leq \overline{B}_{\varepsilon'}(t)$, so $B(t)$ also stays in the interval $(B_2^* - \varepsilon, B_2^* + \varepsilon)$, which we wanted.

In the remaining cases, the stability of a positive component of an equilibrium can be derived from analogous calculations. If we consider a 0 component, e.g. B_1^* , then we use $\overline{B}_{\varepsilon}(t)$ and the constant 0 function as the pinching functions. Thus the previously characterized equilibria of the system are all globally asymptotically stable. ■

Remark 2.8 (Bifurcations). A transcritical bifurcation is the situation when, as a parameter varies, a stable and an unstable equilibrium come together and exchange stability at the critical point (in higher dimensional systems, this happens in some subspace), i.e., there is a stable and an unstable equilibrium both before and after the bifurcation, but their role is reversed ([21]).

In our model, transcritical bifurcations are observed at the points where the sign of a threshold parameters changes. Typically, the positive component of an equilibrium becomes zero at the critical point and changes sign, but then it moves outside of the biologically meaningful domain, so we only track it until reaching 0. Since we can explicitly describe the equilibria and their stability, we can also identify these bifurcations: both the meeting point of two equilibria, as well as how their stability changes. Overall we find the following bifurcations.

- Between Case 1 a) and Case 1 b), i.e. when the sign of $\mathcal{F}_2$ changes. If we increase γ , then the positive $T_{2,1}^*$ will decrease, and eventually become 0. So when $\mathcal{F}_2$ becomes nonpositive, then E_B merges into E_0 .
- Between Case 1 b) and Case 2, i.e. when the sign of $\mathcal{F}_3$ changes. If we increase γ , then the positive $T_{2,2}^*$ will decrease, and eventually become 0. So when $\mathcal{F}_3$ becomes nonpositive, then E_{coex} merges into E_T . The same happens if we increase α instead of γ .
- Between Case 3 and Case 4, i.e. when the sign of $\mathcal{F}_2$ changes. If we increase γ , then the positive $T_{2,1}^*$ will decrease, and eventually become 0. So when $\mathcal{F}_2$ becomes nonpositive, then E_B merges into E_0 .
- Between Case 1 a) and Case 3, i.e. when the sign of $\mathcal{F}_1$ changes. If we increase α , then the positive B_2^* will decrease, and eventually become 0. In this case, two transcritical bifurcations happen simultaneously: when $\mathcal{F}_1$ becomes nonpositive, then E_{coex} and E_T merge into E_B and E_0 , respectively.
- Between Case 2 and Case 4, i.e. when the sign of $\mathcal{F}_1$ changes. If we increase α , then the positive B_2^* will decrease, and eventually become 0. So when $\mathcal{F}_1$ becomes nonpositive, then E_T merges into E_0 .

□

3 Simulations and data fitting

3.1 Simulations

We derived that the treatment has four possible outcomes characterized by the signs of the threshold parameters. These outcomes are illustrated in Figures 4-7., which were made with the following parameter values: $r_T = 0.4$, $K_T = 2.62$, $\delta = 1.15$, $\mu = 2.6$, $\lambda = 2$, $r_B = 1.81$, $K_B = 1.81$, $\theta = 1.085$, $\sigma = 0.7$, and we have $\alpha \in \{1.39, 3.06\}$ and $\gamma \in \{1.24, 3.85\}$. In the colored squares on the left side of the figures ($\alpha \in [0, 6]$, $\gamma \in [0, 6]$) the red points denote the current (α, γ) values for which the simulation was run.




Fig. 4: Both the bacteria population and the tumour survive. Here $\alpha = 1.39$ and $\gamma = 1.24$.




Fig. 5: The bacteria survive, but the tumour dies out. Here $\alpha = 1.39$ and $\gamma = 3.85$.




Fig. 6: The tumour survives, but the bacteria die out. Here $\alpha = 3.06$ and $\gamma = 1.24$.




Fig. 7: Both the tumour and the bacteria population die out. Here $\alpha = 3.06$ and $\gamma = 3.85$.

3.2 Data fitting

In [17], colon cancer cells were implanted under the skin of mice and their growth was monitored, and we use this to test a specified version of our model. Tumour growth was measured over a nine-days interval, while the tumour was treated with Gemcitabine on the fourth and ninth days, as described in the figure caption, and the relative tumour volume (proportional to the first day volume) was plotted as a function of time (Figure 1.B in the article). However, their image description and the measured data is not completely in sync, hence we will fit the timing of the second dose of chemotherapy to the observed data. In this experiment no additional antibiotic treatment was given, hence we omit the A -equation, and consider

$$\begin{cases} T'(t) = r_T T(t) (K_T - T(t)) - \delta T(t) C(t) \\ C'(t) = \gamma - \mu C(t) - \lambda C(t) B(t) \\ B'(t) = r_B B(t) (K_B - B(t).) \end{cases}$$

For convenience, we rewrite the first equation as

$$T'(t) = rT(t) \left(1 - \frac{T(t)}{K_T}\right) - \delta T(t) C(t),$$

where $r = r_T K_T$. During the experiment, which was a relatively short time period, the bacteria population size is assumed to be constant B_0 . As in the experiment, the chemotherapy drug will not be administered continuously, but will be given at times t_1 and t_2 . Thus, except for these two time points, the change of the chemotherapy drug will be described with the equation

$$C'(t) = -(\mu + \lambda B_0) C(t) = -\varphi B_0 C(t), \text{ where } \varphi = \frac{\mu + \lambda B_0}{B_0},$$

and then we can think of φ as a kind of combined rate of decay. Having these modifications, we consider our new system of impulsive differential equations:

$$\begin{cases} T'(t) = rT(t) \left(1 - \frac{T(t)}{K_T}\right) - \delta T(t) C(t) \\ C'(t) = -\varphi B_0 C(t) \\ B'(t) = 0 \end{cases}, \text{ if } t \notin \{t_1, t_2\}, \quad (2)$$

$$\begin{cases} T(t^+) = T(t^-) \\ C(t^+) = C(t^-) + p \\ B(t^+) = B(t^-) \end{cases}, \text{ if } t \in \{t_1, t_2\},$$

where t_i^+ is the time at which the p dose of chemotherapy drug is administered, and t_i^- is the last moment immediately before administration for $i = 1, 2$.

The parameters of the system of equations (2) were optimized using the FindFit procedure of Wolfram Mathematica software (version 12.2), based on the method of least squares. Since it is not known at which time of day the tumour volume was measured or the chemotherapy drug was administered on a given day, the time of drug administration is also a parameter to be optimized. Thus, $r, K_T, \delta, \varphi, t_1, t_2$ are the parameters to be estimated, and the following constraints were imposed in order to obtain a good and feasible fit:

- all parameters are positive,
- the constant amount of the bacteria is $B_0 = 22$ units (this can be scaled by φ , hence this value is actually irrelevant),
- to match the possible dosage times in the experiment, we restrict $3.8 < t_1 < 4.5$ and $6.8 < t_2 < 7.5$.
- for the tumour, the capacity is $K_T > 400$ units,
- the dose of the chemotherapy drug is $p = 45$ units (its effect is scaled by δ hence the exact value is irrelevant).

In the best fit, the least squared error was 24.7869. The data series and the graph of the best fit are shown in Picture 8.




Fig. 8: The best fit of the function $T(t)$ to the measured data. The optimized parameter values are: $r = 0.6867$, $K_T = 51831.9$, $\delta = 0.06312$, $\varphi = 0.1325$, and the precise times of the chemotherapeutic treatment administration are $t_1 = 3.8$, $t_2 = 7.5$.

4 Generalizations and robustness

4.1 More universal growth rate

When setting up the equation system (1), we made several assumptions, one of which was that the growth of the tumour and the bacteria population was modelled by a logistic equation. Instead, however, we can use more general functions with some simple – and biologically realistic – assumptions that ensure that our previous results remain true.

Let $f(T(t))$ be the growth rate at time t for the tumour size $T(t)$, then our equation describing the tumour is

$$T'(t) = f(T(t))T(t).$$

We need f to be a continuous, monotonically decreasing function, but this is a natural assumption, since the larger the tumour, the less nutrients and resources a tumour cell has. Also, we assume that f has a positive root, T^0 . Then $f(T^0)$ will be the nontrivial equilibrium state of the equation, since $T'(t) = f(T^0)T(t) = 0$. For $T(t) < T^0$, $f(T(t))$ is positive, so the tumour increases, while for $T(t) > T^0$, $f(T(t))$ is negative, so the tumour shrinks, thus $f(T(t))$ indeed tends to $f(T^0)$. For the bacteria, a similar condition can be used to define a general growth rate $g(B(t))$. Then our system of equations is modified to the following form:

$$\begin{cases} T'(t) = f(T(t))T(t) - \delta T(t)C(t) \\ C'(t) = \gamma - \mu C(t) - \lambda C(t)B(t) \\ B'(t) = g(B(t))B(t) - \theta B(t)A(t) \\ A'(t) = \alpha - \sigma A(t). \end{cases} \quad (3)$$

As in the case of the equation system (1), it can be seen that for nonnegative initial values, the solutions of the equation system (3) will also be nonnegative and bounded. The statements of Theorem 2.3. and Theorem 2.7. hold, i.e., in the case of more general functions f and g , equilibrium points exist and the solutions always converge to one of these. The proofs can be carried out in almost the same way in both cases. In the proof of Theorem 2.3. we exploit the fact that f and g are monotonically decreasing and both of them have a positive root. Consider, for example, the step when we substitute back the C component of the equilibrium (C^*) into our equation for the tumour:

$$T'(t) = f(T(t))T(t) - \delta C^*T(t) = T(t) (f(T(t)) - \delta C^*),$$

then again there can be at most one positive equilibrium, since $f(T(t)) - \delta C^*$ also has one zero due to the conditions on f . To prove convergence, as in the proof of Theorem 2.7., one can use the previous comparison

principle (we can squeeze the function between $\pm\varepsilon$ versions) and Proposition 2.6 due to the continuity of f and g .

Examples for possible f (and similarly for g) functions:

- logistic equation: $f(T(t)) = r_T (K_T - T(t))$,
- Richards's growth model, which is a generalization of the logistic one: $f(T(t)) = r_T K_T \left(1 - \left(\frac{T(t)}{K_T}\right)^q\right)$, where $q > 0$.
- Gompertz-function: an other commonly used growth model, where the growth of the tumour is described by the function $T'(t) = r_T \ln\left(\frac{K_T}{T(t)}\right) T(t)$, from which we get $f(T(t)) = r_T \ln\left(\frac{K_T}{T(t)}\right)$.

4.2 Pharmacokinetics

During setting up the equation system (1), we assumed that the antibiotic drug kills bacteria in direct proportion to its concentration. In reality, however, this is not necessarily a linear relationship, and the dose-response curves often given by Hill functions:

$$B'(t) = r_B B(t) (K_B - B(t)) - H(A(t)) B(t),$$

where $H(x) = \frac{ax^n}{b+x^n}$ is a Hill-function. Similarly, we can replace the $-\delta T(t)C(t)$ term for cytotoxic effect by the Hill-type term

$$-\delta T(t) \frac{(C(t))^{n_1}}{m + (C(t))^{n_2}}.$$

In this case, however, using a mass action term gave a better fit to data for the experiment described in Section 3.2.

4.3 Bacteria population dynamics

In our model we implicitly assumed that although the bacteria metabolize the chemotherapy drug, it does not substantially influence the growth rate of the bacteria population. We found no evidence for this in [17], but we can not exclude this possibility. In this case, an additional term appears in the equation describing the change in the enhanced growth rate of the bacteria population:

$$B'(t) = (r_B + \eta C(t)) B(t) (K_B - B(t)) - \theta B(t) A(t),$$

where it is assumed that the bacteria benefit in proportion to the amount of chemotherapeutic drug at a given time, and $\eta > 0$ is the parameter describing the extent of this benefit.

4.4 Impulsive drug doses

As we have done in Section 3.2., instead of giving a continuous dose of medicines, we can prescribe periodically taken medication. Then we write the $\gamma(t)$ and $\alpha(t)$ functions in the following forms:

$$\gamma(t) = \sum_{i=1}^{n_C} p_i^C \delta(t - t_i^C) \quad \text{and} \quad \alpha(t) = \sum_{k=1}^{n_A} p_k^A \delta(t - t_k^A),$$

where $\delta(t)$ is the Dirac-delta, t_i^C and t_k^A are the times when the patient takes the chemotherapy drug and the antibiotics, respectively, and p_i^C and p_k^A are the given doses corresponding to those times. This approach requires further consideration, for example, optimizing the time intervals between subsequent medications. By numerical simulations, we can confirm that similarly to the continuous dosage case, the four possible outcomes described earlier are achievable. The similar dynamical behaviour of the continuous and impulsive treatment models is highlighted by two examples shown in Figure 9.




Fig. 9: Comparison of treatment outcomes using continuous and impulsive drug dosage. In the impulsive case, the same amount of drug is given in each time unit, but all at once at integer time points. The left figure depicts a situation when the bacteria population persists, but the tumour is eradicated, while in the right figure the tumour remains despite the bacteria die out. These scenarios correspond to Case 2 and Case 3, respectively, and the plots show the same dynamics for the continuous and the impulsive treatment.

5 Conclusions

Our work was motivated by a recent biological discovery that certain bacteria can metabolize anti-cancer chemotherapy drugs. Therefore, if the bacteria are present in a treated patient, they have the potential to sabotage the treatment, as the metabolized drug cannot exert its anti-tumour effect. In the presence of sabotaging bacteria, it may be recommended to supplement the treatment with antibiotics. To describe this phenomenon, we constructed a mathematical model consisting of four differential equations was constructed, following the variation of the size of the tumour, the concentration of the chemotherapeutic drug, the size of the bacteria population and the concentration of the antibiotic drug. The model expresses the interactions between these components.

We introduced three compound threshold parameters and shown that their signs characterize the existence and stability of the equilibria. We found four possible outcomes, which we explicitly described: both the tumour and the bacterial population die, the tumour dies but the bacteria survive, the tumour survives even though the bacterial population dies, or neither the tumour nor the bacterial population dies. These outcomes are guaranteed by the global asymptotic stability of the corresponding equilibrium. The proof is based on the iterative application of a comparison principle. We highlighted these possibilities in a bifurcation diagram, and described the types of bifurcations in the transitions between the different cases, as single and simultaneous transcritical bifurcations.

Our model was able to reproduce a biological experiment, where rectal cancer tumour was grown in mice and then treated with the chemotherapeutic drug Gemcitabine. The experiment compared the tumour growth of bacteria in mice where the bacteria that metabolize the drug were present or absent. Since the drug was administered at specific time points, the model equations were augmented with corresponding impulsive terms. Parameter estimation was performed using data extracted from [17] and a relatively good fit was obtained. Our model fitting was able to estimate the precise timing of the treatments that was not reported in the biological publication.

We investigated the robustness of our findings by considering possible generalizations of the model and showed how the results extend to such situations. Our model has limitations: we assume homogeneous spatial distribution and a well-mixed system of tumour cells, bacteria and drugs. Potential future research can incorporate this and other important biological factors. In particular, a promising future research direction is the application of a control theory approach to optimize the combined treatment of chemotherapy and antibiotics when the saboteur bacteria are present. The model we presented here contributes to the better understanding of the role of intratumour bacteria in the success or failure of cancer treatment.

410 CRediT authorship contribution statement

411 Anna Geretovszky: Conceptualization, Formal analysis, Methodology, Visualization, Writing - original draft,
412 Writing - review & editing.

413 Gergely Röst: Conceptualization, Formal analysis, Funding acquisition, Methodology, Supervision, Writing -
414 review & editing.

415 Declaration of Competing Interest

416 The authors declare that they have no known competing financial interests or personal relationships that could
417 have appeared to influence the work reported in this paper.

418 Acknowledgement

419 AG was supported by the ÚNKP-23-2 - New National Excellence Program of the Ministry for Culture and
420 Innovation from the source of the National Research, Development and Innovation Fund (grant number ÚNKP-
421 23-2-SZTE-586), and TKP-2021-EGA-05, 2022-2.1.1-NL-2022-00005. GR was supported by RRF-2.3.1-21-2022-
422 00006 and KKP 129877.

References

- [1] Saloni Dattani et al. “Cancer”. In: *Our World in Data* (2024).
- [2] *International Agency for Research on Cancer: Global Cancer Observatory*. URL: <https://gco.iarc.fr/en> (visited on 01/05/2025).
- [3] Global Burden of Disease Collaborative Network. *Global Burden of Disease Study 2021 - with minor processing by Our World in Data*. Institute for Health Metrics and Evaluation (IHME), 2024.
- [4] Yang Kuang, John D. Nagy, and Steffen E. Eikenberry. *Introduction to Mathematical Oncology*. 1st ed. Chapman & Hall/CRC, 2016.
- [5] Russel C. Rockne et al. “The 2019 mathematical oncology roadmap”. In: *Physical Biology* 16.4 (2019). DOI: 10.1088/1478-3975/ab1a09.
- [6] Jacob G Scott et al. “Mathematical modeling of the metastatic process”. In: *Experimental Metastasis: Modeling and Analysis* (2013), pp. 189–208. DOI: 10.1007/978-94-007-7835-1_9.
- [7] Heiko Enderling and Mark Chaplain. “Mathematical Modeling of Tumor Growth and Treatment”. In: *Current Pharmaceutical Design* 20 (2013). DOI: 10.2174/1381612819666131125150434.
- [8] Angela M. Jarrett, Ernesto A.B.F. Lima, et al. “Mathematical Models of Tumor Cell Proliferation: A Review of the Literature”. In: *Expert Review of Anticancer Therapy* 8.12 (2018), pp. 1271–1286. DOI: 10.1080/14737140.2018.1527689.
- [9] Mary K. Anderson and Lauri Matey. “Overview of Cancer and Cancer Treatment”. In: *Chemotherapy and Immunotherapy Guidelines and Recommendations for Practice*. Ed. by MiKaela Olsen, Kristine LeFebvre, and Kelly Brassil. 1st ed. Oncology Nursing Society, 2018. Chap. 4.
- [10] Vikas Malhotra and Michael C. Perry. “Classical Chemotherapy: Mechanisms, Toxicities and the Therapeutic Window”. In: *Cancer Biology & Therapy* 2 (2003). DOI: 10.4161/cbt.199.
- [11] Khalid O. Alfarouk et al. “Resistance to Cancer Chemotherapy: Failure in Drug Response from ADME to P-gp”. In: *Cancer Cell International* 15.71 (2015). DOI: 10.1186/s12935-015-0221-1.
- [12] Marco Gerlinger and Charles Swanton. “How Darwinian models inform therapeutic failure initiated by clonal heterogeneity in cancer medicine”. In: *British Journal of Cancer* 103.8 (2010), pp. 1139–1143. DOI: 10.1038/sj.bjc.6605912.
- [13] Angela Oliveira Pisco, Amy Brock, Joseph Zhou, et al. “Non-Darwinian dynamics in therapy-induced cancer drug resistance”. In: *Nature Communications* 4 (2013). DOI: 10.1038/ncomms3467.
- [14] Arturo Álvarez-Arenas, Ana Podolski-Renic, Juan Belmonte-Beitia, et al. “Interplay of Darwinian Selection, Lamarckian Induction and Microvesicle Transfer on Drug Resistance in Cancer”. In: *Scientific Reports* 9.1 (2019). DOI: 10.1038/s41598-019-45863-z.
- [15] Attila Dénes, Sadegh Marzban, and Gergely Röst. “Global analysis of a cancer model with drug resistance due to Lamarckian induction and microvesicle transfer”. In: *Journal of Theoretical Biology* 527 (2021). DOI: 10.1016/j.jtbi.2021.110812.
- [16] Anyuye Yin et al. “A Review of Mathematical Models for Tumor Dynamics and Treatment Resistance Evolution of Solid Tumors”. In: *CPT: Pharmacometrics & Systems Pharmacology* 8.10 (2019), pp. 720–737. DOI: 10.1002/psp4.12450.
- [17] Leore T. Geller et al. “Potential role of intratumor bacteria in mediating tumor resistance to the chemotherapeutic drug gemcitabine”. In: *Science* 357.63568 (2017), pp. 1156–1160. DOI: 10.1126/science.aah5043.
- [18] Elizabeth A. Sarapata and Lisette de Pillis. “A Comparison and Catalog of Intrinsic Tumor Growth Models”. In: *Bulletin of Mathematical Biology* 76.8 (2014), pp. 2010–2024. DOI: 10.1007/s11538-014-9986-y.
- [19] Horst R. Thieme. *Mathematics in Population Biology*. Princeton University Press, 2003.
- [20] Gerald Teschl. *Ordinary Differential Equations and Dynamical Systems*. American Mathematical Society, 2012.
- [21] Steven H. Strogatz. *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemistry, and Engineering*. Westview Press, 1994.

Declaration of interests

☒The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: